

ight 9.0in
isp

Analisi matematica 1, Area of the Ingegneria of the Informazione
Esercizi in preparazione of the prova scritta
a cura of the docenti of “Analisi Matematica 1”
Anno Accademico 2021-2022

Indice

1 Temi d'esame da 2h	4
2 Temi d'esame da 1h30m for modalità telematica	8
3 Esercizi 2h30m	14
4 Altri esercizi diallenamento	18
5 Temi d'esame of the quattro ultimi anni accademici (solutions in fondo alla dispensa)	20
6 Esercizi scelti dai temi d'esame di anni passati	62
7 Ulteriori esercizi (a cura di C. Sartori)	67
8 Soluzioni of the Temi 1 of the prove scritte of the quattro anni precedenti	70

NOTA: both \ln and \log indicano the logarithm in base e .

Buon lavoro!

1 Temi d'esame da 2h

Appello of the 17.01.2022

THEME 1

Exercise 1 [10 punti] Given the function

$$f(x) = \arctan\left(\frac{|x+1|}{x^2+4}\right),$$

- (i) find the domain, study the sign, compute the limits at the extremes of the domain;
- (ii) study the derivability of f sul suo domain, compute the first derivative, study the monotonicity intervals and points of absolute/relative maximum or minimum;
- (iii) draw the graph.

Exercise 2 [7 punti] Determine the solutions in \mathbb{C} of the equation

$$\left(\frac{z}{i}\right)^3 = -8.$$

Exercise 3 [7 punti]

Study, by utilizing sviluppi of Mac Laurin applicati alla sequenze

$$a_n = \frac{1}{n} - \sin\left(\frac{1}{n}\right) - \alpha \log\left(1 + \frac{1}{n^3}\right),$$

the convergence of the series $\sum_{n=1}^{\infty} n^2 a_n$ for every $\alpha \in \mathbb{R}$.

Exercise 4 [8 punti]

By making use the definizione (and the metodo of sostituzione), compute the integral generalizzato

$$\int_0^{+\infty} \frac{\arctan t}{(1+t^2)(\arctan^2 t + 8 \arctan t + 17)} dt.$$

: Discutere, for all values of the parameter $\alpha \in \mathbb{R}$, the convergence of the generalized integral

$$\int_0^{+\infty} \frac{\arctan t}{(1+t^2)^{2\alpha}(\arctan^2 t + 8 \arctan t + 17)} dt.$$

NB: con log si indica the logarithm in base e .

Tempo a disposizione: 2 ore.

Appello of the 07.02.2022

THEME 1

Exercise 1 [10 punti] Given the function

$$f(x) = \log(|x| - x^2 + 2),$$

- (i) determine the domain; determine the simmetria and the sign; compute the limits and asymptotes at the extremes of the domain;
- (ii) study the derivability and calcolare the first derivative ; study the monotonicity intervals individuando the points of maximum and of minimum both relative and absolute ;
- (iii) draw the graph.

Exercise 2 [7 punti] Determine the insieme A of the numeri complessi $z \in \mathbb{C}$ tali che

$$\frac{|z + i\operatorname{Im}(z)|^2}{|z|^2 + \operatorname{Re}(z)^2} \geq 1$$

and disegnarlo in the Gauss plane .

Exercise 3 [7 punti]

Study the convergence of the series

$$\sum_{n=1}^{\infty} n \left\{ \alpha \sinh \left(\frac{1}{n^2} \right) + \log \left[\cosh \left(\frac{1}{n} \right) \right] \right\}$$

as $\alpha \in \mathbb{R}$.

Exercise 4 [8 punti]

By making use the integration by parts, compute

$$\int \arctan \left(\frac{2}{x} \right) dx.$$

. Study the convergence of the integral improprio

$$\int_0^{+\infty} \arctan \left(\frac{x^3 + 1}{x^\alpha} \right) dx$$

as $\alpha > 0$.

Tempo a disposizione: 2 ore.

Appello of the 01.07.2022

THEME 1

Exercise 1 [9 punti] Consider the function

$$f(x) = |x - 2| e^{\frac{1}{(x-2)^2}}.$$

- (i) determine the domain of f and the sign of f ;
- (ii) compute the main limits of f ;
- (iii) compute the derivative of f , discuss the monotonicity of f and determine the infimum and the supremum of f and relative and absolute minimum and maximum points;
- (iv) compute asymptotes of $f^{(*)}$;
- (v) draw a qualitative graph of f .

(*) this question it is 1 point .

Exercise 2 [8 punti] Determine in algebraic form the solutions in \mathbb{C} of the equation

$$z^4 + (-2 - 2i)z^2 + 4i = 0.$$

Exercise 3 [7 punti]

- (i) Determine, as $\alpha \in \mathbb{R}$, the limit

$$\lim_{x \rightarrow 0^+} \frac{(1+x)^{\alpha x} - 1}{x^2}.$$

Exercise 4 [8 punti] (i) Compute the seguente indefinite integral

$$\int \frac{\sqrt{t}}{1+t} dt.$$

- (ii) Discutere the convergence of the generalized integral

$$\int_0^{+\infty} \frac{\sqrt{t}}{1+t^\alpha} dt$$

as $\alpha \in \mathbb{R}$.

NB: con log si indica the logarithm in base e .

Tempo a disposizione: 2 ore.

Appello of the 12.09.2022

THEME 1

Exercise 1 [8 punti] Given the function

$$f(x) = \arctan\left(\frac{1}{\sin x}\right),$$

- (i) find the domain, study the periodicity and the simmetria, calcolarne the sign, compute the limits at the extremes of the domain;
- (ii) study the derivability of f sul suo domain, compute the first derivative, find the monotonicity intervals and the points of minimum and of maximum, both relative and absolute , and infimum and superiore;
- (iii) draw the graph of f .

Exercise 2 [8 punti] Find the solutions $z \in \mathbb{C}$ of the inequality

$$\left| \frac{z-i}{z-1} \right| \geq 1$$

and le si segni on Gauss plane .

Exercise 3 [8 punti] Compute the limit

$$\lim_{x \rightarrow 0^+} \frac{\sin x - \alpha x + \frac{1}{6}\alpha x^3}{\arctan(x^2 + 4x^3)}$$

for every $\alpha \in \mathbb{R}$.

Exercise 4 [8 punti] (a) Compute the integral definito:

$$\int_{\log 4}^{\log 6} \frac{e^x}{(e^x - 2)(e^x - 1)} dx$$

(b) Al variare of $\alpha \in \mathbb{R}$ Study the convergence of

$$\int_{\log 4}^{+\infty} \frac{e^x}{(e^x - 2)^\alpha (e^x - 1)} dx.$$

NB: con log si inca the logarithm in base e .

Tempo a sposizione: 2 ore.

2 Temi d'esame da 1h30m for modalità telematica

Appello of the 06.07.2020 - Modalità telematica (causa COVID)

THEME 1

Exercise 1 [6 punti] Consider the function

$$f(x) = |(x+3)\log(x+3)|, \quad x \in D =]-3, +\infty[.$$

(i) Compute

$$\lim_{x \rightarrow -3^+} f(x), \quad \lim_{x \rightarrow +\infty} f(x).$$

(ii) Compute the first derivative of the function f , study the monotonicity intervals and draw the graph of f .

Exercise 2 [6 punti] Find the solutions of the equation

$$z^3 = 8i$$

in algebraic and exponential form (or trigonometric), and draw them on the Gauss plane .

Exercise 3 [6 punti] Study the convergence of the series

$$\sum_{n=1}^{\infty} \frac{(1+n^2) \log n}{n^4}.$$

Exercise 4 [6 punti] Compute

$$\int_0^{+\infty} e^{-\sqrt{2x}} dx.$$

Exercise 5 [6 punti] Compute the limit

$$\lim_{x \rightarrow +\infty} x^{\frac{4}{3}} \left(\sqrt[3]{x+2} - \sqrt[6]{x^2-1} \right)^2.$$

Tempo a disposizione: 1 ore and 30 minuti.

Appello of the 14.09.2020 - Modalità telematica (causa COVID)

THEME 1

Exercise 1 [6 punti] Consider the function

$$f(x) = \arctan\left(\frac{x+1}{x-1}\right), \quad x \in (1, \infty).$$

- (i) Inviduarne the asymptotes.
- (ii) If ne determini the monotonicity .

Exercise 2 [6 punti] Consider the complex number $z = \sqrt{3} - i$.

- (i) Scriverlo in exponential form .
- (ii) Compute the real part of z^6 .

Exercise 3 [6 punti] Establish the convergence semplice and assoluta of the series

$$\sum_{n=1}^{\infty} (-1)^n \frac{n}{n^2 + 1}.$$

Exercise 4 [6 punti] Compute the limit

$$\lim_{x \rightarrow 0^+} \frac{\log(1 + \sinh x) - \sin x}{x^2}.$$

Exercise 5 [6 punti] Consider the generalized integral

$$\int_1^{\infty} \log\left(\frac{x^{\alpha}}{x^{\alpha} + 1}\right) dx.$$

- (i) Compute the integral for $\alpha = 2$.
 - (ii) Establish for which $\alpha \in [0, \infty)$ it converges.
- Tempo a sposizione: 1 ore and 30 minuti.

Appello of the 18.01.2021 - Modalità telematica (causa COVID)

THEME 1

Exercise 1 [8 punti] Consider the function

$$f(x) = \arctan\left(\frac{x}{x^2 + x + 1}\right);$$

- (i) inviduarne the domain, stuarne the sign, compute the limits at the extremes of the domain;
- (ii) calcolarne the first derivative, study the monotonicity intervals inviduando the punti estremanti;
- (iii) draw the graph of f .

Exercise 2 [8 punti] Find in \mathbb{C} the solutions of the equation

$$z^4 + (-1 + i)z^2 - i = 0.$$

Suggerimento: sostituire $w = z^2$.

Exercise 3 [8 punti]

(i) Compute

$$\lim_{n \rightarrow \infty} \frac{n^{2n}}{(n+1)^{2n}}.$$

(ii) Study the convergence of the series

$$\sum_{n=1}^{\infty} \frac{(2n)!}{n^{2n}}.$$

Exercise 4 [8 punti] Per $\alpha \in \mathbb{R}$, si consideri

$$f_{\alpha}(x) = \frac{1}{\sinh x + x^{\alpha}}.$$

(a) Study as $\alpha \in \mathbb{R}$ the convergence

$$\int_0^{\log 2} f_{\alpha}(x) dx.$$

(b) Compute

$$\int_0^{\log 2} f_0(x) dx.$$

Tempo a disposizione: 1 ore and 30 minuti.

Appello of the 08.02.2021 - Modalità telematica (causa COVID)

THEME 1

Exercise 1 [8 punti] Consider the function

$$f(x) = \sqrt{\frac{|x|}{x^2 + 1}}.$$

- (i) Determine the domain of f , study the sign and the simmetria of f and compute limits and asymptotes at the extremes of the domain;
- (ii) Study the derivability of f and compute the first derivative, study the monotonicity intervals individuando the points of maximum/ absolute minimum/relativo;
- (iii) draw the graph of f .

Exercise 2 [8 punti] Find the complex solutions of the equation

$$\frac{8}{z^3} = \frac{1+i}{1-i},$$

in algebraic and exponential form (or trigonometric), and draw them on the Gauss plane .

Exercise 3 [8 punti]

(i) Compute

$$\int \log(t+1) dt.$$

(ii) Dedurre the value of

$$\int_0^1 \frac{\log(\sqrt{x}+1)}{\sqrt{x}} dx.$$

Exercise 4 [8 punti](i) Individuare as $\alpha \in \mathbb{R}$ the order diinfinitesimal l of

$$n (\cos(1/n) - 1) + \frac{\alpha}{n}$$

(ii) Study as $\alpha \in \mathbb{R}$ the convergence of

$$\sum_{n=1}^{+\infty} \left| n (\cos(1/n) - 1) + \frac{\alpha}{n} \right|.$$

Tempo a disposizione: 1 ore and 30 minuti.

Appello of the 05.07.2021 - Modalità telematica (causa COVID)**THEME 1****Exercise 1 [8 punti]** Consider the function

$$f(x) = \log \left(1 + \sqrt{1 - x^2} \right).$$

- (i) Determine the domain of f , study the sign and the simmetria of f and compute the limits at the extremes of the domain;
- (ii) Study the derivability of f and compute the first derivative, study the monotonicity intervals and find the points of absolute /relative maximum and minimum ;
- (iii) draw the graph of f .

Exercise 2 [8 punti] Find the complex solutions of the equation

$$\operatorname{Im}(z^2) + |z|^2 \operatorname{Re}\left(\frac{1}{z}\right) = 0,$$

and draw them on the Gauss plane .

Exercise 3 [8 punti]

Sia

$$f_\alpha(x) := \frac{\arctan x}{1 + x^{2\alpha}}.$$

(i) Compute

$$\int f_1(x) dx = \int \arctan x \left(\frac{1}{1 + x^2} \right) dx.$$

(ii) Study as $\alpha \in [0, \infty)$ the convergence of

$$\int_1^{+\infty} f_\alpha(x) dx.$$

Exercise 4 [8 punti]

(i) Compute as $\alpha \in \mathbb{R}$ the limit

$$\lim_{n \rightarrow \infty} \frac{2 \log[\cos(1/n)] + \alpha[\sin(1/n)]^2}{(1/n)^2}.$$

(ii) Dedurre the comportamento of the series

$$\sum_{n=1}^{\infty} \{2 \log[\cos(1/n)] + [\sin(1/n)]^2\}.$$

Tempo a disposizione: 1 ore and 30 minuti.

Appello of the 13.09.2021 - Modalità telematica (causa COVID)

THEME 1

Exercise 1 [8 punti] Consider the function

$$f(x) = \frac{|\sin x|}{1 - 2\cos x} .$$

- (i) Find the domain; study the periodicity , the sign and the simmetria of f ;
- (ii) study the derivability and calcolarne the first derivative ; study the monotonicity intervals and find the points of maximum/ absolute minimum/relativo;
- (iii) draw the graph.

Exercise 2 [8 punti] Find the solutions $z \in \mathbb{C}$ of the inequality

$$\left| \frac{z+1}{z} \right| \geq 1$$

and draw them on the Gauss plane .

Exercise 3 [8 punti] Study the convergence of the series

$$\sum_{n=1}^{\infty} n^{\alpha} \left(\frac{1}{n} - \sin \frac{1}{n} \right)$$

as $\alpha \in \mathbb{R}$.

Exercise 4 [8 punti]

Compute the integral

$$\int_{-1}^0 \frac{x}{x^2 + 2x + 2} dx.$$

Tempo a disposizione: 1 ore and 30 minuti.

3 Esercizi 2h30m

Traccia 1

1) Sia

$$f(x) = \frac{|x - 1| - 2}{x^2 + 1}, \quad x \in [-2, 4].$$

Studiarne the sign, the derivability, the monotonicity and the massimi and minimi locali and absolute .

2) Compute

$$\lim_{x \rightarrow 0} \frac{\sin x^3}{x(1 - \cos x) + x^4}.$$

3) Compute le radici terze of $-27i$.

4) (a) Compute

$$\int_0^{\frac{\pi^2}{4}} \sqrt{x} \sin \sqrt{x} dx.$$

(b*) Determine for which $\alpha \in \mathbb{R}$ the seguente integral converges:

$$\int_0^{\sqrt{\frac{\pi}{2}}} x^\alpha \sin \sqrt{x} dx.$$

5*) Discutere, for all values of the parameter $\alpha \in \mathbb{R}$, the convergence of the series

$$\sum_{n=1}^{\infty} \log \left(n(e^{\frac{1}{n}} - 1) - \frac{\alpha}{n} \right).$$

Tempo a disposizione: 2 ore and 30 minuti. **Si consiglia disvolgere for primi the esercizi senza the asterisco.**

Ogni affermazione deve essere adeguatamente giustificata.

Traccia 2

1) Sia

$$f(x) = \log(|x - 1| + 1) - \log x, \quad x \in]0, 2].$$

Simplify it and study the derivability, the monotonicity and the massimi and minimi locali and absolute .

2) Compute

$$\lim_{x \rightarrow 0} \frac{1 - \cos x^2}{x(x - \sin x) + x^3 \sin^2 x}.$$

3) Simplify the expression

$$\frac{\overline{(1+i)}^2}{(1-i)^2 \left(\frac{-1}{i} + \sqrt{3}\right)}$$

writing the result in algebraic form and in trigonometric form .

4) (a) Compute

$$\int_{\log \pi}^{2 \log \pi} e^{2x} \cos e^x dx.$$

(b*) Study the convergence of the generalized integral

$$\int_{-\infty}^{\log \pi} e^{\alpha x} \cos e^x dx$$

as $\alpha \in \mathbb{R}$.

5*) Sia

$$f(x) = \int_{\sqrt{\frac{\pi}{2}}}^x \sin(t^2) dt.$$

(a) Computethe Taylor expansion of f diorder 2 at the point $x_0 = \sqrt{\frac{\pi}{2}}$ (letting the value of $f(\sqrt{\frac{\pi}{2}})$ as known);

(b) study the monotonicity and the convexity and the concavity of f in the interval $[-1, 2]$;

(c) compute , as $\lambda \in \mathbb{R}$, the numero of the solutions of the equation $f(x) = \lambda$ contained in the interval $[-1, 2]$.

Tempo a disposizione: 2 ore and 30 minuti. **Si consiglia disvolgere for primi the esercizi senza the asterisco.**

Ogni affermazione deve essere adeguatamente giustificata.

Traccia 3

1) Sia

$$f(x) = \arctan|x^2 - 1|, \quad x \in [-1, 2].$$

Studiarne the derivability, the monotonicity and the massimi and minimi locali and absolute .

2) Compute

$$\lim_{x \rightarrow 0} \frac{(1 - \cosh x)^2}{\sin x(x - \arctan x) + x^3 \sinh^2 x}.$$

3) Solve the equation

$$(z^2 + 2i)(z^3 + 8) = 0$$

writing the result in algebraic form and in trigonometric form .

4) (a) Compute

$$\int_{\log 3}^1 \frac{e^x}{e^{2x} - 3e^x + 2} dx.$$

(b*) Study the convergence of the generalized integral

$$\int_{\log 2}^{+\infty} \frac{e^x}{(e^{2x} - 3e^x + 2)^\alpha} dx$$

as $\alpha \in \mathbb{R}$.

5*) Sia

$$f(x) = \begin{cases} \alpha(\arctan \sin x + 1) & \text{for } x \leq 0 \\ e^{x+1} & \text{for } x > 0. \end{cases}$$

(a) Determine all the $\alpha \in \mathbb{R}$ such that the graph of f admits a tangent line in $(0, f(0))$ and compute it for tali α ;

(b) discuss , as $\lambda \in \mathbb{R}$, the numero of the solutions of the equation $f(x) = e + \lambda x$ contained in the interval $[0, 2]$.

Tempo a disposizione: 2 ore and 30 minuti. **Si consiglia disvolgere for primi the esercizi senza the asterisco.**

Ogni affermazione deve essere adeguatamente giustificata.

Traccia 4

1) [6 punti] Study the absolute convergence and the convergence of the series

$$\sum_{n=1}^{\infty} \frac{(\sin x)^n}{n}$$

for all values of the parameter $x \in [0, 2\pi[$.

2) [4 punti] Compute

$$\lim_{x \rightarrow 0^+} \frac{e^{-\frac{1}{x}}}{\arctan x}.$$

3) [4 punti] Solve the inequality

$$|e^{i \operatorname{Re} z} (\bar{z} - i)| \leq 1$$

and draw the solutions in the Gauss plane .

4) [4 + 4 punti]

(a) Compute

$$\int_0^2 \sqrt{4 - x^2} dx \quad (\text{eseguire a sostituzione iperbolica}).$$

(b*) Study the convergence of the generalized integral

$$\int_0^2 (4 - x^2)^\alpha dx$$

as $\alpha \in \mathbb{R}$.

5*) [7 punti] Consider the function

$$f(x) = |1 - x| e^{\arctan(4/x)}.$$

- 1) Determine the domain, compute the main limits of f and determine the asymptotes.
 - 2) Compute f' nei punti where is possibile and determine the monotonicity intervals and the points of extreme of f .
 - 3) Draw a graph of f .
-

Tempo a disposizione: 2 ore and 30 minuti. **Si consiglia disvolgere for primi the esercizi senza the asterisco.**

Ogni affermazione deve essere adeguatamente giustificata.

4 Altri esercizi di allenamento

Limits .

- 1) Compute the limits

$$\lim_{x \rightarrow 2} \frac{\sqrt{x} - 2}{x - 2}, \quad \lim_{x \rightarrow -\infty} x \log \frac{1-x}{3-x}, \quad \lim \left(\frac{x^2}{x^2 - 2} \right)^{x^2}.$$

- 2) Compute the limits

$$\lim_{n \rightarrow \infty} \frac{2^n - n! \sin \frac{1}{n} - n}{2^{n-1} + (n-1)!}, \quad \lim_{n \rightarrow \infty} \frac{e^n - n \sin n}{n! - 2^n}.$$

Series.

- 1) Determine the character of the series

$$\sum_{n=1}^{\infty} 2^{\frac{1}{n}}, \quad \sum_{n=1}^{\infty} \frac{1 - \cos n}{n^2}, \quad \sum_{n=1}^{\infty} (-1)^n \frac{n}{n^2 + n - 1}$$

- 2) Study the convergence and the absolute convergence for all values of the parameter $x \in \mathbb{R}$ of the series

$$\sum_{n=1}^{\infty} \left(1 - \cos \frac{x^n}{n} \right), \quad \sum_{n=1}^{\infty} \arctan \frac{x^n}{n}.$$

Funzioni.

- 1) Discutere la derivabilità e calcolare le derivate prime e seconde delle funzioni

$$f_1(x) = \log \frac{1}{\cos x}, \quad f_2(x) = \log |\sin x - \frac{1}{2}|$$

in il loro dominio.

- 2) Verificare l'identità

$$\arcsin x = \arctan \frac{x}{\sqrt{1-x^2}}, \quad x \in]-1, 1[.$$

- 3) Studiare la monotonicità e determinare i punti di massimo e minimo relativo e assoluto di

$$f_1(x) = \sin x - x \cos x, \quad f_2(x) = \sqrt{x} - \sqrt{x-1} \text{ (per } x \in [0, 1]), \quad \arctan \left| x - \frac{1}{x} \right| \text{ (per } x \neq 0).$$

- 4) Studiare il numero di soluzioni dell'equazione

$$x - \log |x| = \alpha \quad (x \neq 0)$$

per $\alpha \in \mathbb{R}$.

Integrali.

- 1) Calcolare il polinomio di Taylor di ordine 3 con centro $x_0 = 1$ delle funzioni

$$F_1(x) = \int_1^x \frac{e^t}{t} dt \quad F_2(x) = \int_1^x \frac{\log t}{t} dt$$

and dire if in the interval $[1, 2]$ they are invertibili.

2) Compute the integral i

$$\int x \log^2 x \, dx, \quad \int_0^1 \frac{x^2 - 4}{x^2 + 5x + 4} \, dx, \quad \int \frac{x}{\cos^2 x} \, dx, \quad \int_1^{+\infty} \frac{dx}{x^2 + x}, \quad \int_1^2 \frac{1}{\sqrt{x^2 - 1}} \, dx.$$

3) Study the convergence degli integral i

$$\int_0^{+\infty} \frac{x^\alpha}{\sqrt{e^x - 1}} \, dx, \quad \int_0^{+\infty} \frac{dx}{x^\alpha + x^2}$$

as $\alpha \in \mathbb{R}$.

5 Temi d'esame of the quattro ultimi anni accademici (solutions in fondo alla dispensa)

Appello of the 23.01.2017

THEME 1

Exercise 1 [6 punti] Compute the integral

$$\int_{\log 3}^2 \frac{e^x}{e^{2x} - 4} dx$$

Exercise 2 [6 punti] Solve the inequality

$$|2z^2 - 2\bar{z}^2| < 3$$

and draw the solutions on Gauss plane .

Exercise 3 [6 punti] Study the convergence of the series

$$\sum_{n=1}^{+\infty} n^2 \left(\cos \frac{1}{n} - 1 + \sin \frac{1}{2n^\alpha} \right)$$

for all values of the parameter $\alpha > 0$.

Exercise 4 [8 punti] Consider the function

$$f(x) := \arcsin \frac{|x| - 4}{x^2 + 2}.$$

- i) Determine the domain D of f , its simmetries and study the sign; determine the limits of f at the extremes of D ;
- ii) study the derivability, compute the derivative and study the monotonicity of f ; determine the points of extreme relative and absolute and compute the main limits of f' ;
- iii) draw a qualitative graph of f .

Exercise 5 [6 punti] Study the convergence of the generalized integral

$$\int_0^{+\infty} \frac{\arctan x |\arctan(x-1)|}{|1-x^2|^\alpha (\sinh \sqrt{x})^\beta} dx$$

as $\alpha, \beta \in \mathbb{R}$.

Exercise . Sia I a interval chiuso and limitato and sia $f : I \rightarrow \mathbb{R}$ a function continuous and tale che $f(x) \in I$ for every $x \in I$. Dimostrare that esiste almeno a $x \in I$ tale che $f(x) = x$.

THEME 2

Exercise 1 [6 punti] Compute the integral

$$\int_0^1 \frac{e^x}{e^{2x} + 4e^x + 5} dx$$

Exercise 2 [6 punti] Solve the inequality

$$|4\bar{z}^2 - 4z^2| < 5$$

and draw the solutions on Gauss plane .

Exercise 3 [6 punti] Study the convergence of the series

$$\sum_{n=1}^{+\infty} n^2 (2 - e^{1/2n^\alpha} - \cos(1/n))$$

for all values of the parameter $\alpha > 0$.

Exercise 4 [8 punti] Consider the function

$$f(x) := \arcsin \frac{4 - |x|}{1 + 2x^2}.$$

- i) Determine the domain D of f , its simmetries and study the sign; determine the limits of f at the extremes of D ;
- ii) study the derivability, compute the derivative and study the monotonicity of f ; determine the points of extreme relative and absolute and compute the main limits of f' ;
- iii) draw a qualitative graph of f .

Exercise 5 [6 punti] Study the convergence of the generalized integral

$$\int_0^{+\infty} \frac{|\arctan(x-2)| \arctan x}{|x^2 - 4|^\alpha (\sinh \sqrt[3]{x})^\beta} dx$$

as $\alpha, \beta \in \mathbb{R}$.

THEME 3

Exercise 1 [6 punti] Compute the integral

$$\int_{\log 4}^3 \frac{e^x}{e^{2x} - 9} dx$$

Exercise 2 [6 punti] Solve the inequality

$$|3z^2 - 3\bar{z}^2| < 2$$

and draw the solutions on Gauss plane .

Exercise 3 [6 punti] Study the convergence of the series

$$\sum_{n=1}^{+\infty} n^2 (\cosh(1/n^\alpha) + \cos(1/n) - 2)$$

for all values of the parameter $\alpha > 0$.

Exercise 4 [8 punti] Consider the function

$$f(x) := \arcsin \frac{|x| - 4}{2x^2 + 3}.$$

- i) Determine the domain D of f , its symmetries and study the sign; determine the limits of f at the extremes of D ;
- ii) study the derivability, compute the derivative and study the monotonicity of f ; determine the points of extreme relative and absolute and compute the main limits of f' ;
- iii) draw a qualitative graph of f .

Exercise 5 [6 punti] Study the convergence of the generalized integral

$$\int_0^{+\infty} \frac{|\arctan(3-x)| \arctan x}{|9-x^2|^\alpha (\cosh \sqrt{x}-1)^\beta} dx$$

as $\alpha, \beta \in \mathbb{R}$.

THEME 4

Exercise 1 [6 punti] Compute the integral

$$\int_0^1 \frac{e^x}{e^{2x} - 4e^x + 5} dx$$

Exercise 2 [6 punti] Solve the inequality

$$|9\bar{z}^2 - 9z^2| < 2$$

and draw the solutions on Gauss plane .

Exercise 3 [6 punti] Study the convergence of the series

$$\sum_{n=1}^{+\infty} n^2 (e^{1/n^2} - \tan 1/n^\alpha - 1)$$

for all values of the parameter $\alpha > 0$.

Exercise 4 [8 punti] Consider the function

$$f(x) := \arcsin \frac{4 - 4|x|}{5x^2 + 3}.$$

- i) Determine the domain D of f , its simmetries and study the sign; determine the limits of f at the extremes of D ;
- ii) study the derivability, compute the derivative and study the monotonicity of f ; determine the points of extreme relative and absolute and compute the main limits of f' ;
- iii) draw a qualitative graph of f .

Exercise 5 [6 punti] Study the convergence of the generalized integral

$$\int_0^{+\infty} \frac{\arctan x |\arctan(1-2x)|}{|1-4x^2|^\alpha (\cosh x - 1)^\beta} dx$$

as $\alpha, \beta \in \mathbb{R}$.

Appello of the 13.02.2017

THEME 1

Exercise 1 [8 punti] Consider the function

$$f(x) := \log |x^2 - 2x - 3|.$$

- i) Determine the domain D and study the sign of f ; determine the limits of f at the extremes of D and the asymptotes;
- ii) study the derivability, compute the derivata, study the monotonicity and determine the points of extreme relative and absolute of f ;
- (iii) compute f'' and study the concavity and the convexity of f ;
- (iv) draw a qualitative graph of f .

Exercise 2 [5 punti] Study the convergence of the series

$$\sum_{n=1}^{+\infty} \frac{1}{2^n} \frac{n^n}{n!}.$$

Exercise 3 [4 punti] Given

$$f(z) = \frac{2+iz}{iz+1},$$

determine the domain and determine all the $z \in \mathbb{C}$ tali che $f(z) = z$. Express tutte the solutions in algebraic form.

Exercise 4 [7 punti] Compute the limit

$$\lim_{x \rightarrow 0^+} \frac{\arctan x - \sin x + x^{\frac{10}{3}} \log x}{x^\alpha (1 - \cos^2 x)}$$

as $\alpha > 0$.

Exercise 5 [8 punti] Study the convergence of the generalized integral

$$\int_2^{+\infty} \frac{1}{x^\alpha \sqrt{x-2}} dx$$

as $\alpha \in \mathbb{R}$ and calcolarlo for $\alpha = 1$.

THEME 2

Exercise 1 [8 punti] Consider the function

$$f(x) := \log |x^2 + x - 6|.$$

- i) Determine the domain D and study the sign of f ; determine the limits of f at the extremes of D and the asymptotes;
- ii) study the derivability, compute the derivata, study the monotonicity and determine relative and absolute extreme points of f ;
- (iii) compute f'' and study the concavity and the convexity of f ;
- (iv) draw a qualitative graph of f .

Exercise 2 [5 punti] Study the convergence of the series

$$\sum_{n=1}^{+\infty} \left(\frac{2}{3}\right)^n \frac{n^n}{n!}.$$

Exercise 3 [4 punti] Given

$$f(z) = \frac{-1 - 2iz}{iz - 1},$$

determine the domain and determine all the $z \in \mathbb{C}$ tali che $f(z) = 2z$. Express tutte the solutions in algebraic form.

Exercise 4 [7 punti] Compute the limit

$$\lim_{x \rightarrow 0^+} \frac{\arctan x - \sinh x + x^{\frac{11}{2}} \log x}{x^\alpha (1 - \cosh^2 x)}$$

as $\alpha > 0$.

Exercise 5 [8 punti] Study the convergence of the generalized integral

$$\int_3^{+\infty} \frac{1}{x^\alpha \sqrt{x-3}} dx$$

as $\alpha \in \mathbb{R}$ and calcolarlo for $\alpha = 1$.

THEME 3

Exercise 1 [8 punti] Consider the function

$$f(x) := \log |x^2 - 2x - 8|.$$

- i) Determine the domain D and study the sign of f ; determine the limits of f at the extremes of D and the asymptotes;
- ii) study the derivability, compute the derivata, study the monotonicity and determine the points of extreme relative and absolute of f ;

- (iii) compute f'' and study the concavity and the convexity of f ;
- (iv) draw a qualitative graph of f .

Exercise 2 [5 punti] Study the convergence of the series

$$\sum_{n=1}^{+\infty} \frac{1}{3^n} \frac{n^n}{n!}.$$

Exercise 3 [4 punti] Given

$$f(z) = \frac{-2 + 3iz}{2iz - 3},$$

determine the domain and determine all the $z \in \mathbb{C}$ tali che $f(z) = -z$. Express tutte the solutions in algebraic form.

Exercise 4 [7 punti] Compute the limit

$$\lim_{x \rightarrow 0^+} \frac{x^{\frac{9}{2}} \log x - \tan x + \sin x}{x^\alpha (1 - \cosh^2 x)}$$

as $\alpha > 0$.

Exercise 5 [8 punti] Study the convergence of the generalized integral

$$\int_4^{+\infty} \frac{1}{x^\alpha \sqrt{x-4}} dx$$

as $\alpha \in \mathbb{R}$ and calcolarlo for $\alpha = 1$.

THEME 4

Exercise 1 [8 punti] Consider the function

$$f(x) := \log |x^2 + 3x - 4|.$$

- i) Determine the domain D and study the sign of f ; determine the limits of f at the extremes of D and the asymptotes;
- ii) study the derivability, compute the derivata, study the monotonicity and determine the points of extreme relative and absolute of f ;
- (iii) compute f'' and study the concavity and the convexity of f ;
- (iv) draw a qualitative graph of f .

Exercise 2 [5 punti] Study the convergence of the series

$$\sum_{n=1}^{+\infty} \left(\frac{2}{7}\right)^n \frac{n^n}{n!}.$$

Exercise 3 [4 punti] Given

$$f(z) = \frac{1 - 4iz}{iz + 4},$$

determine the domain and determine all the $z \in \mathbb{C}$ tali che $f(z) = z$. Express tutte the solutions in algebraic form.

Exercise 4 [7 punti] Compute the limit

$$\lim_{x \rightarrow 0^+} \frac{\sinh x - \tan x - x^{\frac{15}{4}} \log x}{x^\alpha (1 - \cos^2 x)}$$

as $\alpha > 0$.

Exercise 5 [8 punti] Study the convergence of the generalized integral

$$\int_5^{+\infty} \frac{1}{x^\alpha \sqrt{x-5}} dx$$

as $\alpha \in \mathbb{R}$ and calcolarlo for $\alpha = 1$.

Appello of the 10.07.2017

THEME 1

Exercise 1 [8 punti] Consider the function

$$f(x) := \log |e^{2x} - 4|.$$

- i) Determine the domain D and study the sign of f ; determine the limits of f at the extremes of D and the asymptotes;
- ii) study the derivability, compute the derivata, study the monotonicity and determine the points of extreme relative and absolute of f ;
- iii) compute f'' and study the concavity and the convexity of f ;
- iv) draw a qualitative graph of f .

Exercise 2 [5 punti] Draw in the Gauss plane the insieme

$$S := \left\{ z \in \mathbb{C} : \operatorname{Re} \frac{z-1}{z-i} \geq 0, |z+1-i| \leq 1 \right\}.$$

Exercise 3 [5 punti] Compute the integral

$$\int e^{2x} \arctan(3e^x) dx.$$

Exercise 4 [7 punti] Compute the limit

$$\lim_{x \rightarrow 0^+} \frac{\arctan \sin x - \sinh x}{x^\alpha (1 - \cos^2 x)}$$

for all values of the parameter $\alpha > 0$.

Exercise 5 [7 punti] Study the convergence semplice and assoluta di

$$\sum_{n=2}^{+\infty} \frac{(1-e^a)^n}{n + \sqrt{n}}$$

as $a \in \mathbb{R}$.

THEME 2

Exercise 1 [8 punti] Consider the function

$$f(x) := \log |e^{-3x} - 9|.$$

- i) Determine the domain D and study the sign of f ; determine the limits of f at the extremes of D and the asymptotes;
- ii) study the derivability, compute the derivata, study the monotonicity and determine the points of extreme relative and absolute of f ;
- iii) compute f'' and study the concavity and the convexity of f ;
- iv) draw a qualitative graph of f .

Exercise 2 [5 punti] Draw in the Gauss plane the insieme

$$S := \left\{ z \in \mathbb{C} : \operatorname{Re} \frac{z+1}{z-i} > 0, |z-1-i| \leq 1 \right\}.$$

Exercise 3 [5 punti] Compute the integral

$$\int e^{2x} \arctan(2e^x) dx.$$

Exercise 4 [7 punti] Compute the limit

$$\lim_{x \rightarrow 0^+} \frac{\sin \arctan x - \sinh x}{x^\alpha (1 - \cosh^2 x)}$$

for all values of the parameter $\alpha > 0$.

Exercise 5 [7 punti] Study the convergence semplice and assoluta of

$$\sum_{n=2}^{+\infty} \frac{(1-2^a)^n}{n + \log n}$$

as $a \in \mathbb{R}$.

Appello of the 18.09.2017

THEME 1

Exercise 1 [8 punti] Consider the function

$$f(x) := \frac{3x}{\log |2x|}.$$

- i) Determine the domain D and study le simmetries and the sign of f ; determine the limits of f at the extremes of D , the prolungabilità of f and the asymptotes;

- ii) study the derivability, compute the derivative and its main limits , study the monotonicity e determine the points of extreme relative and absolute of f ;
- iii) compute f'' and study the concavity and the convexity of f ;
- iv) draw a qualitative graph of f .

Exercise 2 [5 punti] Given the polynomial

$$z^4 + z^3 + 8thez + 8i$$

determine prima a Root Test intera and le other roots , writing them in algebraic form.

Exercise 3 [5 punti] Study the convergence of the series

$$\sum_{n=1}^{\infty} \left(1 + \frac{3x}{n}\right)^{n^2}$$

as $x \in \mathbb{R}$.

Exercise 4 [7 punti] Compute, for all values of the real parameter α , the limit

$$\lim_{x \rightarrow 0^+} \frac{\cosh(\alpha x) - e^{x^2} + x \log(\cos x)}{x - \sin x + e^{-1/x^2}}.$$

Exercise 5 [7 punti] Study the convergence of the generalized integral

$$\int_0^{+\infty} xe^{ax} (2 + \cos x) dx$$

as $a \in \mathbb{R}$. Compute

$$\int_0^{+\infty} xe^{-x} \cos x dx$$

(sugg.: compute preliminarly a primitive of $e^{-x} \cos x$).

THEME 2

Exercise 1 [8 punti] Consider the function

$$f(x) := \frac{2x}{\log|3x|}.$$

- i) Determine the domain D and study le simmetries and the sign of f ; determine the limits of f at the extremes of D , the prolungabilità of f and the asymptotes;
- ii) study the derivability, compute the derivative and its main limits , study the monotonicity e determine the points of extreme relative and absolute of f ;
- iii) compute f'' and study the concavity and the convexity of f ;
- iv) draw a qualitative graph of f .

Exercise 2 [5 punti] Given the polynomial

$$z^4 - z^3 - 27thez + 27i$$

determine prima a Root Test intera and le other roots , writing them in algebraic form.

Exercise 3 [5 punti] Study the convergence of the series

$$\sum_{n=1}^{\infty} \left(1 - \frac{2x}{n}\right)^{n^2}$$

as $x \in \mathbb{R}$.

Exercise 4 [7 punti] Compute, for all values of the real parameter α , the limit

$$\lim_{x \rightarrow 0^+} \frac{\cos x - e^{\alpha x^2} + x \log(\cosh x)}{x - \sinh x + e^{-1/x^2}}.$$

Exercise 5 [7 punti] Study the convergence of the generalized integral

$$\int_0^{+\infty} xe^{ax} (2 - \sin x) dx$$

as $a \in \mathbb{R}$. Compute

$$\int_0^{+\infty} xe^{-x} \sin x dx$$

(sugg.: compute preliminarily a primitive of $e^{-x} \sin x$).

Appello of the 29.01.2018

THEME 1

Exercise 1 [6 punti] Consider the function

$$f(x) := \log \frac{|x^2 - 5|}{x + 1}.$$

- i) Determine the domain D of f , its simmetries and study the sign; determine the limits of f at the extremes of D and the asymptotes;
- ii) study the derivability, compute the derivative and study the monotonicity of f and determine the points of extreme relative and absolute ;the study of the second derivative may be skipped
- iii) draw a qualitative graph of f .

Exercise 2 [6 punti] Consider the sequence

$$a_n = \frac{(-1)^n e^{2n} \sin \frac{1}{n}}{(n-1)!}, \quad n \in \mathbb{N}, n \geq 2.$$

- a) Compute $\lim_{n \rightarrow \infty} a_n$;
- b) study the absolute convergence and the convergence semplice of the series $\sum_{n=2}^{\infty} a_n$.

Exercise 3 [5 punti] Sia $f(z) = z^2 + \bar{z}|z|$. Solve the equation

$$zf(z) = |z|^3 - 8i,$$

writing the solutions in algebraic form and disegnandole in the Gauss plane .

Exercise 4 [7 punti] Compute the limit

$$\lim_{x \rightarrow +\infty} \frac{\log(x+3) - \log(x+1) - \sin \frac{2}{x}}{\cos \sin \frac{1}{2x} - e^{\frac{\alpha}{x^2}} - e^{-x}}$$

as $\alpha \in \mathbb{R}$.

Exercise 5 [8 punti] a) Study the convergence of the generalized integral

$$\int_{\sqrt{2}}^{+\infty} \frac{1}{x^\alpha \sqrt{x^2 - 2}} dx$$

as $\alpha \in \mathbb{R}$;

b) calcolarlo for $\alpha = 1$.

Exercise . Sia $x_0 \in \mathbb{R}$ and define the sequence $\{a_n : n \in \mathbb{N}\}$ ponendo

$$a_0 = x_0 \text{ e, for every } n \geq 1, a_{n+1} = \sin a_n.$$

- a) prove that a_n is definitively monotonic for $n \rightarrow +\infty$;
- b) prove that $\lim_{n \rightarrow +\infty} a_n = 0$.

THEME 2

Exercise 1 [6 punti] Consider the function

$$f(x) := \log \frac{|x^2 - 3|}{x + 1}.$$

- i) Determine the domain D of f , its simmetries and study the sign; determine the limits of f at the extremes of D and the asymptotes;
- ii) study the derivability, compute the derivative and study the monotonicity of f and determine the points of extreme relative and absolute ;the study of the second derivative may be skipped;
- iii) draw a qualitative graph of f .

Exercise 2 [6 punti] Consider the sequence

$$a_n = \frac{(-1)^n e^{3n} \sinh \frac{1}{n}}{(n-1)!}, \quad n \in \mathbb{N}, n \geq 2.$$

- a) Compute $\lim_{n \rightarrow \infty} a_n$;
- b) study the absolute convergence and the convergence semplice of the series $\sum_{n=2}^{\infty} a_n$.

Exercise 3 [5 punti] Sia $f(z) = -z^2 + \bar{z}|z|$. Solve the equation

$$zf(z) = |z|^3 - 8i,$$

writing the solutions in algebraic form and disegnandole in the Gauss plane .

Exercise 4 [7 punti] Compute the limit

$$\lim_{x \rightarrow +\infty} \frac{\log(x+1) - \log(x+2) + \sinh \frac{1}{x}}{\cosh \sin \frac{1}{x} - e^{\frac{\alpha}{x^2}} - e^{-2x}}$$

as $\alpha \in \mathbb{R}$.

Exercise 5 [8 punti] a) Study the convergence of the generalized integral

$$\int_{\frac{1}{2}}^{+\infty} \frac{1}{x^\alpha \sqrt{4x^2 - 1}} dx$$

as $\alpha \in \mathbb{R}$;

b) calcolarlo for $\alpha = 1$.

THEME 3

Exercise 1 [6 punti] Consider the function

$$f(x) := \log \frac{|x^2 - 4|}{x - 1}.$$

- i) Determine the domain D of f , its simmetries and study the sign; determine the limits of f at the extremes of D and the asymptotes;
- ii) study the derivability, compute the derivative and study the monotonicity of f and determine the points of extreme relative and absolute ;the study of the second derivative may be skipped ;
- iii) draw a qualitative graph of f .

Exercise 2 [6 punti] Consider the sequence

$$a_n = \frac{(-1)^n e^{\frac{n}{2}} \arctan \frac{1}{n}}{(n-1)!}, \quad n \in \mathbb{N}, n \geq 2.$$

- a) Compute $\lim_{n \rightarrow \infty} a_n$;
- b) study the absolute convergence and the convergence semplice of the series $\sum_{n=2}^{\infty} a_n$.

Exercise 3 [5 punti] Sia $f(z) = z^2 + \bar{z}|z|$. Solve the equation

$$zf(z) = |z|^3 + 27i,$$

writing the solutions in algebraic form and disegnandole in the Gauss plane .

Exercise 4 [7 punti] Compute the limit

$$\lim_{x \rightarrow +\infty} \frac{\log(x-2) - \log(x-1) + \arctan \frac{1}{x}}{\cosh \sinh \frac{2}{x} - \cos \frac{\alpha}{x} - e^{-\frac{x}{2}}}$$

as $\alpha \in \mathbb{R}$.

Exercise 5 [8 punti] a) Study the convergence of the generalized integral

$$\int_2^{+\infty} \frac{1}{x^\alpha \sqrt{x^2 - 4}} dx$$

as $\alpha \in \mathbb{R}$;

b) calcolarlo for $\alpha = 1$.

THEME 4

Exercise 1 [6 punti] Consider the function

$$f(x) := \log \frac{|x^2 - 6|}{x + 1}.$$

- i) Determine the domain D of f , its simmetries and study the sign; determine the limits of f at the extremes of D and the asymptotes;
- ii) study the derivability, compute the derivative and study the monotonicity of f and determine the points of extreme relative and absolute ; the study of the second derivative may be skipped ;
- iii) draw a qualitative graph of f .

Exercise 2 [6 punti] Consider the sequence

$$a_n = \frac{(-1)^n e^{\frac{n}{3}} \tan \frac{1}{n}}{(n-1)!}, \quad n \in \mathbb{N}, n \geq 2.$$

- a) Compute $\lim_{n \rightarrow \infty} a_n$;
- b) study the absolute convergence and the convergence semplice of the series $\sum_{n=2}^{\infty} a_n$.

Exercise 3 [5 punti] Sia $f(z) = -z^2 + \bar{z}|z|$. Solve the equation

$$zf(z) = |z|^3 + 27i,$$

writing the solutions in algebraic form and disegnandole in the Gauss plane .

Exercise 4 [7 punti] Compute the limit

$$\lim_{x \rightarrow +\infty} \frac{\log(x+3) - \log(x+1) - \tan \frac{2}{x}}{\cosh \sinh \frac{3}{x} - \cosh \frac{\alpha}{x} - e^{-3x}}$$

as $\alpha \in \mathbb{R}$.

Exercise 5 [8 punti] a) Study the convergence of the generalized integral

$$\int_{\frac{1}{3}}^{+\infty} \frac{1}{x^\alpha \sqrt{9x^2 - 1}} dx$$

as $\alpha \in \mathbb{R}$;

b) calcolarlo for $\alpha = 1$.

Appello of the 16.02.2018

THEME 1

Exercise 1 [7 punti] Consider the function

$$f(x) = \begin{cases} e^{x-\frac{1}{|x-2|}} & \text{for } x \neq 2 \\ 0 & \text{for } x = 2. \end{cases}$$

- i) Determine the domain D of f , its simmetries and study the sign; determine the limits of f at the extremes of D and the asymptotes;
- ii) si dica if f is continuous in the whole \mathbb{R} .
- iii) compute the derivative and study the monotonicity of f and determine the points of extreme relative and absolute ; compute the main limits of f' ; in particolare si dica if f is differentiable in the whole \mathbb{R} ; the study of the second derivative can be skipped ;
- iv) draw a qualitative graph of f .

Exercise 2 [6 punti] Study as $x \in \mathbb{R}$ the convergence semplice and assoluta of the series

$$\sum_{n=1}^{\infty} \frac{(2x-1)^n}{(2n+3)^2}.$$

Exercise 3 [6 punti] Solve the equation

$$z^2\bar{z} + z\bar{z}^2 = 4 \operatorname{Im}(iz)$$

and draw the solutions on Gauss plane .

Exercise 4 [6 punti]

Compute the limit

$$\lim_{x \rightarrow 0} \frac{(4 \cos x - \alpha)^2 - 4x^4}{x^4 \sin^2 x}$$

as $\alpha \in \mathbb{R}$.

Exercise 5 [7 punti] a) Study the convergence of the generalized integral

$$\int_0^{\frac{\pi^2}{3}} x^\alpha \sin(\sqrt{3x}) dx$$

as $\alpha \in \mathbb{R}$;

b) calcolarlo for $\alpha = \frac{1}{2}$.

THEME 2

Exercise 1 [7 punti] Consider the function

$$f(x) = \begin{cases} e^{-x-\frac{1}{|x+2|}} & \text{for } x \neq -2 \\ 0 & \text{for } x = -2. \end{cases}$$

- i) Determine the domain D of f , its simmetries and study the sign; determine the limits of f at the extremes of D and the asymptotes;
- ii) si dica if f is continuous in the whole \mathbb{R} .
- iii) compute the derivative and study the monotonicity of f and determine the points of extreme relative and absolute ; compute the main limits of f' ; in particolare si dica if f is differentiable in the whole \mathbb{R} ; the study of the second derivative may be skipped ;
- iv) draw a qualitative graph of f .

Exercise 2 [6 punti] Study as $x \in \mathbb{R}$ the convergence semplice and assoluta of the series

$$\sum_{n=1}^{\infty} \frac{(x+2)^n}{(3n+2)^2}.$$

Exercise 3 [6 punti] Solve the equation

$$-\operatorname{Im}(z^2\bar{z} - z\bar{z}^2) = 8i(z - \bar{z})$$

and draw the solutions on Gauss plane .

Exercise 4 [6 punti]

Compute the limit

$$\lim_{x \rightarrow 0} \frac{4(\cosh x - \alpha)^2 - x^4}{x^4 \arctan^2 x}$$

as $\alpha \in \mathbb{R}$.

Exercise 5 [7 punti] a) Study the convergence of the generalized integral

$$\int_0^{\frac{\pi^3}{2}} x^{\alpha-1} \sin(\sqrt[3]{2x}) dx$$

as $\alpha \in \mathbb{R}$;

b) calcolarlo for $\alpha = 1$.

THEME 3

Exercise 1 [7 punti] Consider the function

$$f(x) = \begin{cases} e^{x-\frac{1}{|x-3|}} & \text{for } x \neq 3 \\ 0 & \text{for } x = 3 \end{cases}$$

- i) Determine the domain D of f , its simmetries and study the sign; determine the limits of f at the extremes of D and the asymptotes;
- ii) si dica if f is continuous in the whole \mathbb{R} .
- iii) compute the derivative and study the monotonicity of f and determine the points of extreme relative and absolute ; compute the main limits of f' ; in particolare si dica if f is differentiable in the whole \mathbb{R} ; the study of the second derivative may be skipped ;
- iv) draw a qualitative graph of f .

Exercise 2 [6 punti] Study as $x \in \mathbb{R}$ the convergence semplice and assoluta of the series

$$\sum_{n=1}^{\infty} \frac{(3x+1)^n}{(2n+5)^2}.$$

Exercise 3 [6 punti] Solve the equation

$$z\bar{z}^2 - z^2\bar{z} = 2i \operatorname{Im}(\bar{z} - z)$$

and draw the solutions on Gauss plane .

Exercise 4 [6 punti]

Compute the limit

$$\lim_{x \rightarrow 0} \frac{(e^{x^2} - 2\alpha)^2 - x^4}{x^4 \sinh^2 x}$$

as $\alpha \in \mathbb{R}$.

Exercise 5 [7 punti] a) Study the convergence of the generalized integral

$$\int_0^{\frac{\pi^2}{8}} x^{1-\alpha} \sin(\sqrt{2x}) dx$$

as $\alpha \in \mathbb{R}$;

b) calcolarlo for $\alpha = \frac{1}{2}$.

THEME 4

Exercise 1 [7 punti] Consider the function

$$f(x) = \begin{cases} e^{-x - \frac{1}{|x+3|}} & \text{for } x \neq -3 \\ 0 & \text{for } x = -3 \end{cases}$$

- i) Determine the domain D of f , its simmetries and study the sign; determine the limits of f at the extremes of D and the asymptotes;
- ii) si dica if f is continuous in the whole \mathbb{R} .
- iii) compute the derivative and study the monotonicity of f and determine the points of extreme relative and absolute ; compute the main limits of f' ; in particolare si dica if f is differentiable in the whole \mathbb{R} ; the

study of the second derivative may be skipped ;

iv) draw a qualitative graph of f .

Exercise 2 [6 punti] Study as $x \in \mathbb{R}$ the convergence semplice and assoluta of the series

$$\sum_{n=1}^{\infty} \frac{(x-3)^n}{(3n+5)^2}.$$

Exercise 3 [6 punti] Solve the equation

$$\operatorname{Im}(\bar{z}^2 z - z^2 \bar{z}) = 4 \operatorname{Re}(iz)$$

and draw the solutions on Gauss plane .

Exercise 4 [6 punti]

Compute the limit

$$\lim_{x \rightarrow 0} \frac{2(3\alpha - e^{x^2})^2 - 2x^4}{x^4 \tan^2 x}$$

as $\alpha \in \mathbb{R}$.

Exercise 5 [7 punti] a) Study the convergence of the generalized integral

$$\int_0^{\frac{\pi^3}{24}} x^\alpha \sin(\sqrt[3]{3x}) dx$$

as $\alpha \in \mathbb{R}$;

b) calcolarlo for $\alpha = 0$.

Appello of the 9.07.2018

THEME 1

Exercise 1 [6 punti] Consider the function

$$f(x) = \log |2 - 3e^{3x}|.$$

i) Si determini the domain D and study the sign of f ;

ii) si determinino the limits of f at the extremes of D and the asymptotes;

iii) find the derivative and study the monotonicity of f , determinando the points of extreme relative and absolute ; the study of the second derivative may be skipped ;

iv) draw a qualitative graph of f .

Exercise 2 [6 punti] Solve the inequality

$$|z|^2 \operatorname{Re}\left(\frac{1}{z}\right) \leq \operatorname{Im}(\bar{z}^2)$$

rappresentandone the solutions on Gauss plane .

Exercise 3 [6 punti] Compute the limit

$$\lim_{x \rightarrow +\infty} \frac{(\log(1+x) - \log x - \frac{\alpha}{x})^2}{(1 - \cos \frac{1}{x})^2 + e^{-x}}$$

as $\alpha \in \mathbb{R}$.

Exercise 4 [6 punti] Study as $\alpha \in \mathbb{R}$ the convergence of the series

$$\sum_{n=1}^{\infty} n \arctan \left(\frac{2^{\alpha n}}{n} \right).$$

Exercise 5 [8 punti] a) Compute a primitive di

$$f(x) = \frac{x^2}{(x^2+1)(x^2+2)}$$

(sugg.: cercare a decomposizione of the integrand of the tipo $\frac{A}{x^2+1} + \frac{B}{x^2+2}$).

b) Study the convergence of the generalized integral

$$\int_0^{+\infty} \log \frac{x^\alpha + 2}{x^\alpha + 1} dx.$$

as $\alpha > 0$.

c) Compute the integral for $\alpha = 2$.

THEME 2

Exercise 1 [6 punti] Consider the function

$$f(x) = \log |2e^{2x} - 3|.$$

- i) Si determini the domain D and study the sign of f ;
- ii) si determinino the limits of f at the extremes of D and the asymptotes;
- iii) find the derivative and study the monotonicity of f , determinandone the points of extreme relative and absolute ; the study of the second derivative may be skipped;
- iv) draw a qualitative graph of f .

Exercise 2 [6 punti] Solve the inequality

$$\operatorname{Im} \left(\frac{1}{z} \right) \geq \frac{\operatorname{Im}(z^2 - \bar{z}^2)}{|z|^2}$$

rappresentandone the solutions on Gauss plane .

Exercise 3 [6 punti] Compute the limit

$$\lim_{x \rightarrow +\infty} \frac{(\cosh \frac{1}{x} - 1)^2 - e^{-x}}{\left(\log(2+x) - \log x + \frac{2\alpha}{x} \right)^2}$$

as $\alpha \in \mathbb{R}$.

Exercise 4 [6 punti] Study as $\alpha \in \mathbb{R}$ the convergence of the series

$$\sum_{n=1}^{\infty} n^2 \arctan \left(\frac{4^{\alpha n}}{n^2} \right).$$

Exercise 5 [8 punti] a) Compute a primitive di

$$f(x) = \frac{x^2}{(x^2+4)(x^2+1)}$$

(sugg.: cercare a decomposizione of the integrand of the tipo $\frac{A}{x^2+1} + \frac{B}{x^2+4}$).

b) Study the convergence of the generalized integral

$$\int_0^{+\infty} \log \frac{x^\alpha + 1}{x^\alpha + 4} dx.$$

as $\alpha > 0$.

c) Compute the integral for $\alpha = 2$.

Appello of the 17.09.2018

THEME 1

Exercise 1 [7 punti] Consider the function

$$f(x) := \begin{cases} e^{-\frac{2}{|x|}} (2|x| - 3) & \text{for } x \neq 0 \\ 0 & \text{for } x = 0. \end{cases}$$

- i) Determine the domain D , le simmetries and study the sign of f ;
- ii) determine the limits of f at the extremes of D and the asymptotes;
- iii) compute the derivative and study the monotonicity of f and determine the points of extreme relative and absolute ; the study of the second derivative may be skipped ;
- iv) study the continuity and () the derivability of f (in particolare in $x = 0$);
- v) draw a qualitative graph of f .

Exercise 2 [6 punti] Sia

$$P_\lambda(z) = \lambda - 4t \operatorname{the} z + 2iz^2 + z^3.$$

Find $\lambda_0 \in \mathbb{C}$ in modo che $z = -2i$ sia a zero of P_{λ_0} . Solve the equation

$$P_{\lambda_0}(z) = 0$$

and express the solutions in algebraic form.

Exercise 3 [6 punti] Discutere for all values of the real parameter α the convergence of the series

$$\sum_{n=2}^{\infty} \frac{\log(n + \sin n)}{n^{\frac{\alpha}{2}} + 2}$$

Exercise 4 [6 punti] Compute as $\alpha \in \mathbb{R}^+$ the limit

$$\lim_{x \rightarrow 0^+} \frac{x - \sinh x - x^\alpha}{\cos x - 1 + x^{\frac{7}{3}} \log x}.$$

Exercise 5 [7 punti] Given the integral

$$\int_0^{\frac{1}{\sqrt{2}}} x^{\frac{\alpha}{2}} \arcsin 2x^2 dx,$$

- a) study the convergence as $\alpha \in \mathbb{R}$;
- b) calcolarlo for $\alpha = 2$.

THEME 2

Exercise 1 [7 punti] Consider the function

$$f(x) := \begin{cases} e^{-\frac{1}{|x|}} (2 - 3|x|) & \text{for } x \neq 0 \\ 0 & \text{for } x = 0. \end{cases}$$

- i) Find the domain D , le simmetries and study the sign of f ;
- ii) determine the limits of f at the extremes of D and the asymptotes;
- iii) compute the derivative and study the monotonicity of f and determine the points of extreme relative and absolute ; the study of the second derivative may be skipped ;
- iv) study the continuity and () the derivability of f (in particolare in $x = 0$);
- v) draw a qualitative graph of f .

Exercise 2 [6 punti] Sia

$$P_\lambda(z) = \lambda + 2iz + 3iz^2 + z^3.$$

Determine $\lambda_0 \in \mathbb{C}$ in modo che $z = -3i$ sia a zero of P_{λ_0} . Solve the equation

$$P_{\lambda_0}(z) = 0$$

and express the solutions in algebraic form.

Exercise 3 [6 punti] Discutere for all values of the real parameter α the convergence of the series

$$\sum_{n=1}^{\infty} \frac{\log(n + \cos n)}{n^{2\alpha} + 1}$$

Exercise 4 [6 punti] Compute as $\alpha \in \mathbb{R}^+$ the limit

$$\lim_{x \rightarrow 0^+} \frac{\sin x - x - x^\alpha}{\cosh x - 1 + x^{\frac{5}{2}} \log x}.$$

Exercise 5 [7 punti] Given the integral

$$\int_0^{\sqrt{2}} x^{2\alpha} \arcsin \frac{x^2}{2} dx,$$

- a) study the convergence as $\alpha \in \mathbb{R}$;
- b) calcolarlo for $\alpha = \frac{1}{2}$.

Appello of the 21.01.2019

THEME 1

Exercise 1 [6 punti] Consider the function

$$f(x) = e^{\frac{|x^2 - 16|}{x+3}}, \quad x \in D =]-\infty, -3[.$$

- i) determine the limits of f at the extremes of D and the asymptotes;
- ii) study the derivability, calcolare the derivata, study the monotonicity, determine the points of extreme relative and absolute and draw the graph.

Exercise 2 [4 punti] Compute the limit

$$\lim_{x \rightarrow 0^+} \frac{e^{2x} - 1 - \sin(2x)}{\sinh^2 x + x^{\frac{9}{2}}}.$$

Exercise 3 [4 punti] Solve the equation

$$iz^2 + (1 + 2i)z + 1 = 0$$

in $z \in \mathbb{C}$, writing the solutions in algebraic form.

Exercise 4 [5+3+3 punti] Siano $\alpha \in \mathbb{R}$ fissato and

$$f(t) := \frac{\log(1 + \frac{t}{2})}{t^{2\alpha}}.$$

- i) Compute $\int_1^2 f(t) dt$ con $\alpha = 1$.
- ii) Sia $F(x) := \int_2^x f(t) dt$ con $\alpha = \frac{1}{2}$. Scrivere la Taylor formula di seconda ordine per F centrata in $x = 2$.
- iii) Determinare per quali $\alpha \in \mathbb{R}$ esiste finito $\int_0^1 f(t) dt$.

Exercise 5 [7 punti] Studiare la convergenza semplice e assoluta della serie

$$\sum_{n=1}^{+\infty} \frac{(\log \alpha)^n}{1 + \sqrt{2n}}$$

per $\alpha \in]0, +\infty[$.

Exercise Determinare tutti i valori di $a \in \mathbb{R}$ tali che la funzione $f(x) = e^x - ax^3$ sia convessa in tutto il campo \mathbb{R} .

THEME 2

Exercise 1 [6 punti] Considerare la funzione

$$f(x) = e^{\frac{|x^2-4|}{x+1}}, \quad x \in D =]-\infty, -1[.$$

- i) determinare i limiti di f nei punti estremi di D e le asintoti;
- ii) studiare la derivabilità, calcolare la derivata, studiare la monotonicità, determinare i punti di estrema relativa e assoluta e disegnare il grafico.

Exercise 2 [4 punti] Calcolare il limite

$$\lim_{x \rightarrow 0^+} \frac{\sinh(3x) - \log(1 + 3x)}{\sin^2 x + x^{\frac{11}{2}}}.$$

Exercise 3 [4 punti] Risolvere l'equazione

$$iz^2 + (-1 - 2i)z + 1 = 0$$

in $z \in \mathbb{C}$, scrivendo le soluzioni in forma algebrica.

Exercise 4 [5+3+3 punti] Siano $\alpha \in \mathbb{R}$ fissato e

$$f(t) := \frac{\log(1 + \frac{t}{4})}{t^{\frac{\alpha}{2}}}.$$

- i) Calcolare $\int_1^4 f(t) dt$ con $\alpha = 4$.
- ii) Sia $F(x) := \int_4^x f(t) dt$ con $\alpha = 2$. Scrivere la Taylor formula di seconda ordine per F centrata in $x = 4$.

iii) Determine for which $\alpha \in \mathbb{R}$ esiste finito $\int_0^1 f(t) dt$.

Exercise 5 [7 punti] Study the convergence semplice and assoluta of the series

$$\sum_{n=1}^{+\infty} \frac{(\tan \alpha)^n}{\sqrt{2n} - 1}$$

as $\alpha \in]-\pi/2, +\pi/2[$.

Exercise Determine all the values of $a \in \mathbb{R}$ such that the function $f(x) = e^x - ax^3$ sia convex in the whole \mathbb{R} .

THEME 3

Exercise 1 [6 punti] Consider the function

$$f(x) = e^{\frac{|x^2-3|}{x-1}}, \quad x \in D =]-\infty, 1[.$$

- i) determine the limits of f at the extremes of D and the asymptotes;
- ii) study the derivability, calcolarne the derivata, study the monotonicity, determine the points of extreme relative and absolute and draw the graph.

Exercise 2 [4 punti] Compute the limit

$$\lim_{x \rightarrow 0^+} \frac{e^{3x} - 1 - \sinh(3x)}{\log^2(1+x) + x^{2\pi}}.$$

Exercise 3 [4 punti] Solve the equation

$$iz^2 + (1-2i)z - 1 = 0$$

in $z \in \mathbb{C}$, writing the solutions in algebraic form.

Exercise 4 [5+3+3 punti] Siano $\alpha \in \mathbb{R}$ fissato and

$$f(t) := \frac{\log(1+2t)}{t^{\alpha-1}}.$$

- i) Compute $\int_1^{\frac{3}{2}} f(t) dt$ con $\alpha = 3$.
- ii) Sia $F(x) := \int_3^x f(t) dt$ con $\alpha = 2$. Scrivere the Taylor formula of the second order for F centrata in $x = 3$.
- iii) Determine for which $\alpha \in \mathbb{R}$ esiste finito $\int_0^1 f(t) dt$.

Exercise 5 [7 punti] Study the convergence semplice and assoluta of the series

$$\sum_{n=2}^{+\infty} \frac{(1 + \log \alpha)^n}{\sqrt{n} - 1}$$

as $\alpha \in]0, +\infty[$.

Exercise Determine all the values of $a \in \mathbb{R}$ such that the function $f(x) = e^x - ax^3$ sia convex in the whole \mathbb{R} .

THEME 4

Exercise 1 [6 punti] Consider the function

$$f(x) = e^{\frac{|x^2 - 5|}{x-2}}, \quad x \in D =]-\infty, 2[.$$

- i) determine the limits of f at the extremes of D and the asymptotes;
- ii) study the derivability, calcolarne the derivata, study the monotonicity, determine the points of extreme relative and absolute and draw the graph.

Exercise 2 [4 punti] Compute the limit

$$\lim_{x \rightarrow 0^+} \frac{\sinh(2x) - \log(1 + 2x)}{\arctan(x^2) + x^{2e}}.$$

Exercise 3 [4 punti] Solve the equation

$$iz^2 + (-1 + 2i)z - 1 = 0$$

in $z \in \mathbb{C}$, writing the solutions in algebraic form.

Exercise 4 [5+3+3 punti] Siano $\alpha \in \mathbb{R}$ fissato and

$$f(t) := \frac{\log(1 + \frac{t}{3})}{t^{\alpha+1}}.$$

- i) Compute $\int_1^3 f(t) dt$ con $\alpha = 1$.
- ii) Sia $F(x) := \int_3^x f(t) dt$ con $\alpha = 0$. Scrivere the Taylor formula of the second order for F centrata in $x = 3$.
- iii) Determine for which $\alpha \in \mathbb{R}$ esiste finito $\int_0^1 f(t) dt$.

Exercise 5 [7 punti] Study the convergence semplice and assoluta of the series

$$\sum_{n=1}^{+\infty} \frac{(\tan 2\alpha)^n}{1 + \sqrt{n}}$$

as $\alpha \in]-\pi/4, +\pi/4[$.

Exercise Determine all the values of $a \in \mathbb{R}$ such that the function $f(x) = e^x - ax^3$ sia convex in the whole \mathbb{R} .

Appello of the 11.02.2019

THEME 1

Exercise 1 [6 punti] Sia

$$f(x) = |(x+3) \log(x+3)|, \quad x \in D =]-3, +\infty[.$$

- (i) Determine i limits of f at the extremes of D and the asymptotes; study the prolongabilità for continuity in $x = -3$;
- (ii) study the derivability, calcolarne the derivata, study the monotonicity, determine the points of extreme relative and absolute and draw the graph.

Exercise 2 [4 punti] Study the convergence of the series

$$\sum_{n=1}^{\infty} \frac{(1+n^2) \sin n}{n^4}$$

Exercise 3 [4 punti] Solve the inequality

$$\frac{1}{2} \leq \frac{(\operatorname{Re}(\bar{z} + i) - 1)^2}{4} + \frac{(\operatorname{Im}(\bar{z} + i) - 1)^2}{4} \leq 1$$

and draw the solutions on Gauss plane .

Exercise 4 [5 punti] Compute

$$\int_0^{+\infty} e^{-\sqrt{2x}} dx.$$

Exercise 5 [3+3 punti] Sia

$$f_\alpha(x) = \frac{e^{-\sqrt{2x}} - 1}{x^{\alpha-1}}.$$

(a) study the convergence of the integral

$$\int_0^{+\infty} f_\alpha(x) dx$$

as $\alpha \in \mathbb{R}$.

(b) Per $\alpha = 2$, sia $F(x) = \int_1^{\cos x} f_\alpha(t) dt$: si calcoli $F'(\pi/3)$.

Exercise 6 [7 punti] Compute the limit

$$\lim_{x \rightarrow 0^+} \frac{\cosh(\alpha x) - \cosh(e^{2x} - 1)}{x^3}$$

for all values of the parameter $\alpha > 0$.

Exercise Compute

$$\lim_{x \rightarrow +\infty} \int_x^{x+e^{-x}} e^t \arctan t dt.$$

THEME 2

Exercise 1 [6 punti] Sia

$$f(x) = |(x+2)\log(x+2)|, \quad x \in D =]-2, +\infty[.$$

- (i) Determine i limits of f at the extremes of D and the asymptotes; study the prolongabilità for continuity in $x = -2$;
(ii) study the derivability, calcolarne the derivata, study the monotonicity, determine the points of extreme relative and absolute and draw the graph.

Exercise 2 [4 punti] Study the convergence of the series

$$\sum_{n=2}^{\infty} \frac{n^3 \sin n}{1 - n^5}$$

Exercise 3 [4 punti] Solve the inequality

$$\frac{1}{3} \leq \frac{(\operatorname{Re}(\bar{z} + 2i) - 1)^2}{9} + \frac{(\operatorname{Im}(\bar{z} + 2i) - 1)^2}{9} \leq 1$$

and draw the solutions on Gauss plane .

Exercise 4 [5 punti] Compute

$$\int_0^{+\infty} e^{-\sqrt{3x}} dx.$$

Exercise 5 [3+3 punti] Sia

$$f_\alpha(x) = \frac{e^{-\sqrt{3x}} - 1}{x^{2\alpha+1}}.$$

- (a) study the convergence of the integral

$$\int_0^{+\infty} f_\alpha(x) dx$$

as $\alpha \in \mathbb{R}$.

- (b) Per $\alpha = 0$, sia $F(x) = \int_1^{\sin x} f_\alpha(t) dt$: si calcoli $F'(\pi/6)$.

Exercise 6 [7 punti] Compute the limit

$$\lim_{x \rightarrow 0^+} \frac{\cos(\alpha x) - \cos \log(1 + 5x)}{x^3}$$

for all values of the parameter $\alpha > 0$.

Exercise Compute

$$\lim_{x \rightarrow +\infty} \int_x^{x+e^{-x}} e^t \arctan t dt.$$

THEME 3

Exercise 1 [6 punti] Sia

$$f(x) = |(x+1)\log(x+1)|, \quad x \in D =]-1, +\infty[.$$

(i) Determine i limits of f at the extremes of D and the asymptotes; study the prolungabilità for continuity in $x = -1$;

(ii) study the derivability, calcolarne the derivata, study the monotonicity, determine the points of extreme relative and absolute and draw the graph.

Exercise 2 [4 punti] Study the convergence of the series

$$\sum_{n=2}^{\infty} \frac{n^2 \sin(n^2)}{1-n^5}$$

Exercise 3 [4 punti] Solve the inequality

$$\frac{1}{2} \leq \frac{(\operatorname{Re}(\bar{z}-i)-1)^2}{9} + \frac{(\operatorname{Im}(\bar{z}-i)-1)^2}{9} \leq 1$$

and draw the solutions on Gauss plane .

Exercise 4 [5 punti] Compute

$$\int_0^{+\infty} e^{-\sqrt{x/2}} dx.$$

Exercise 5 [3+3 punti] Sia

$$f_{\alpha}(x) = \frac{e^{-\sqrt{x/2}} - 1}{x^{\alpha-3}}.$$

(a) study the convergence of the integral

$$\int_0^{+\infty} f_{\alpha}(x) dx$$

as $\alpha \in \mathbb{R}$.

(b) Per $\alpha = 4$, sia $F(x) = \int_1^{\sinh x} f_{\alpha}(t) dt$: si calcoli $F'(\log 3)$.

Exercise 6 [7 punti] Compute the limit

$$\lim_{x \rightarrow 0^+} \frac{\cos(\alpha x) - \cos \log(1+2x)}{x^3}$$

for all values of the parameter $\alpha > 0$.

Exercise Compute

$$\lim_{x \rightarrow +\infty} \int_x^{x+e^{-x}} e^t \arctan t dt.$$

THEME 4

Exercise 1 [6 punti] Sia

$$f(x) = |(x+4) \log(x+4)|, \quad x \in D =]-4, +\infty[.$$

(i) Determine i limits of f at the extremes of D and the asymptotes; study the prolongabilità for continuity in $x = -4$;

(ii) study the derivability, calcolarne the derivata, study the monotonicity, determine the points of extreme relative and absolute and draw the graph.

Exercise 2 [4 punti] Study the convergence of the series

$$\sum_{n=1}^{\infty} \frac{(2-n^2) \sin(n^2)}{n^5}$$

Exercise 3 [4 punti] Solve the inequality

$$\frac{1}{3} \leq \frac{(\operatorname{Re}(\bar{z} - 2i) - 1)^2}{4} + \frac{(\operatorname{Im}(\bar{z} - 2i) - 1)^2}{4} \leq 1$$

and draw the solutions on Gauss plane .

Exercise 4 [5 punti] Compute

$$\int_0^{+\infty} e^{-\sqrt{x/3}} dx.$$

Exercise 5 [3+3 punti] Sia

$$f_{\alpha}(x) = \frac{e^{-\sqrt{x/3}} - 1}{x^{2\alpha-1}}.$$

(a) study the convergence of the integral

$$\int_0^{+\infty} f_{\alpha}(x) dx$$

as $\alpha \in \mathbb{R}$.

(b) Per $\alpha = 1$, sia $F(x) = \int_1^{\arctan x} f_{\alpha}(t) dt$: si calcoli $F'(\sqrt{3})$.

Exercise 6 [7 punti] Compute the limit

$$\lim_{x \rightarrow 0^+} \frac{\cosh(\alpha x) - \cosh(1 - e^{3x})}{x^3}$$

for all values of the parameter $\alpha > 0$.

Exercise Compute

$$\lim_{x \rightarrow +\infty} \int_x^{x+e^{-x}} e^t \arctan t dt.$$

Appello of the 8.07.2019

THEME 1

Exercise 1 [6 punti] Sia

$$f(x) = e^{\frac{2}{|2+\log x|}}.$$

- a) Determine the domain D of f ; determine the limits of f at the extremes of D and study the prolongabilità for continuity di f in $x = 0$;
- b) study the derivability, compute the derivative and study the monotonicity of f ; determine the points of extreme relative and absolute ;
- c) draw a qualitative graph of f .

Exercise 2 [4 punti] Study the convergence of the series

$$\sum_{n=1}^{\infty} \frac{\sin \frac{1}{n}}{1 - 2\sqrt{n}}.$$

Exercise 3 [4 punti] Solve the equation

$$\frac{z}{\bar{z}} = -\frac{(\operatorname{Im} z)^2}{|iz^2|}$$

and draw the solutions on Gauss plane .

Exercise 4 [5+3+4 punti] a) Compute a primitive of the function

$$e^x \log(1 + 2e^x).$$

Per $\alpha \in \mathbb{R}$, define $f_\alpha(x) = e^{\alpha x} \log(1 + 2e^x)$:

- b) study the convergence of the generalized integral

$$\int_0^{+\infty} f_\alpha(x) dx$$

as $\alpha \in \mathbb{R}$;

- c) find the Taylor expansion diorder 2 centered in $x_0 = 1$ of the function

$$F(x) = \int_1^x f_0(t) dt.$$

Exercise 5 [6 punti] Compute the limit

$$\lim_{x \rightarrow +\infty} x^\alpha \left(\sqrt[8]{x^2 - 2} - \sqrt[4]{x + 1} \right)$$

for all values of the parameter $\alpha > 0$.

THEME 2

Exercise 1 [6 punti] Sia

$$f(x) = e^{\frac{1}{|3+\log x|}}.$$

- a) Determine the domain D of f ; determine the limits of f at the extremes of D and study the prolongabilità for continuity di f in $x = 0$;
b) study the derivability, compute the derivative and study the monotonicity of f ; determine the points of extreme relative and absolute ;
c) draw a qualitative graph of f .

Exercise 2 [4 punti] Study the convergence of the series

$$\sum_{n=1}^{\infty} (1 - \sqrt{n}) \sinh \frac{1}{n^2}.$$

Exercise 3 [4 punti] Solve the equation

$$\frac{z}{\bar{z}} = \frac{(\operatorname{Re} z)^2}{|iz^2|}$$

and draw the solutions on Gauss plane .

Exercise 4 [5+3+4 punti] a) Compute a primitive of the function

$$e^x \log(1 + 3e^x).$$

Per $\alpha \in \mathbb{R}$, define $f_\alpha(x) = e^{\alpha x} \log(1 + 3e^x)$:

- b) study the convergence of the generalized integral

$$\int_0^{+\infty} f_\alpha(x) dx$$

as $\alpha \in \mathbb{R}$;

- c) find the Taylor expansion diorder 2 centered in $x_0 = 2$ of the function

$$F(x) = \int_2^x f_0(t) dt.$$

Exercise 5 [6 punti] Compute the limit

$$\lim_{x \rightarrow +\infty} x^\alpha \left(\sqrt[3]{x+2} - \sqrt[6]{x^2-1} \right)$$

for all values of the parameter $\alpha > 0$.

Appello of the 17.09.2019

THEME 1

Exercise 1 [7 punti] Sia

$$f(x) = \log |e^{3x} - 2|.$$

- a) Determine the domain D and study the sign of f ; determine the limits of f at the extremes of D and determine the asymptotes;
- b) study the derivability, compute the derivative and study the monotonicity of f ; determine the points of extreme relative and absolute ;
- c) draw a qualitative graph of f .

Exercise 2 [5 punti] Compute the limit

$$\lim_{x \rightarrow 0^+} \frac{e^{x-2x^2} - 1 - x}{\sinh x^2 + x^{7/3} \log x}.$$

Exercise 3 [4 punti] Solve the inequality

$$\operatorname{Re} z \leq \operatorname{Re} \left(\frac{3}{z} \right)$$

and draw the solutions on Gauss plane .

Exercise 4 [6+3 punti] a) Compute the indefinite integral

$$\int \left(\tan \frac{x}{2} \right)^3 dx \quad (\text{sugg.: eseguire la sostituzione } \tan \frac{x}{2} = u).$$

b) study the convergence of the generalized integral

$$\int_0^{\frac{\pi}{6}} \frac{\tan x}{x^{\alpha+2}} dx$$

as $\alpha \in \mathbb{R}$.

Exercise 5 [4+3 punti] (i) Si dimostri that the sequence

$$a_n = \log(n+1) - \log \sqrt{n^2 + \alpha n + 4}$$

is infinitesimal for $n \rightarrow \infty$ (for every α) and for $\alpha = 2$ compute the order ;

(ii) study the convergence of the series

$$\sum_{n=2}^{\infty} a_n$$

as $\alpha \in \mathbb{R}$.

THEME 2

Exercise 1 [7 punti] Sia

$$f(x) = \log |e^{2x} - 3|.$$

- a) Determine the domain D and study the sign of f ; determine the limits of f at the extremes of D and determine the asymptotes;
 b) study the derivability, compute the derivative and study the monotonicity of f ; determine the points of extreme relative and absolute ;
 c) draw a qualitative graph of f .

Exercise 2 [5 punti] Compute the limit

$$\lim_{x \rightarrow 0^+} \frac{e^{x-3x^2} - 1 - x}{\sin x^2 + x^{5/2} \log x}.$$

Exercise 3 [4 punti] Solve the inequality

$$\operatorname{Re} z \leq \operatorname{Re} \left(\frac{4}{z} \right)$$

and draw the solutions on Gauss plane .

Exercise 4 [6+3 punti] a) Compute the indefinite integral

$$\int (\tan 2x)^3 dx \quad (\text{sugg.: eseguire la sostituzione } \tan 2x = u).$$

b) study the convergence of the generalized integral

$$\int_0^{\frac{\pi}{6}} \frac{\tan x}{x^{2\alpha-1}} dx$$

as $\alpha \in \mathbb{R}$.

Exercise 5 [4+3 punti] (i) Si dimostri that the sequence

$$a_n = \log(n+1) - \log \sqrt{n^2 + \alpha n + 3}$$

is infinitesimal for $n \rightarrow \infty$ (for every α) and for $\alpha = 2$ compute the order ;

(ii) study the convergence of the series

$$\sum_{n=2}^{\infty} a_n$$

as $\alpha \in \mathbb{R}$.

Appello of the 20.01.2020

THEME 1

Exercise 1 [7 punti] Consider the function

$$f(x) = \sin(2 \arctan(|x|^3))$$

- i) determine the domain D , the sign, simmetries, i limits at the extremes of D and the asymptotes;
- ii) study the derivability, calcolarne the derivata, study the monotonicity, determine the points of extreme relative and absolute ; the study of the second derivative may be skipped .
- iii) draw the qualitative graph .

Exercise 2 [6 punti] Compute the limit

$$\lim_{x \rightarrow 0^+} (1 + \sin x)^{x^a}$$

as $a \in \mathbb{R}$, usando the form “ $\exp\{\log \dots\}$ ”.

Exercise 3 [4 punti] Trovare the zeros in \mathbb{C} di

$$(z^3 + 5)(z^2 + z + 1) = 0.$$

Exercise 4 [4+3 punti] Siano $\alpha \in \mathbb{R}$ fissato and

$$f_\alpha(t) := \frac{e^{2t} + 2e^t}{(e^t - 1)^\alpha}.$$

- i) Compute a primitive of f_α con $\alpha = 1$.
- ii) Determine for which $\alpha \in \mathbb{R}$ esiste finito $\int_0^1 f_\alpha(t) dt$.

Exercise 5 [6 punti] Study the convergence semplice and assoluta of the series

$$\sum_{n=1}^{+\infty} \frac{(3 \sin x)^n n}{n^2 + \sqrt{n}}$$

as $x \in [-\frac{\pi}{2}, \frac{\pi}{2}]$.

Exercise Sia $\{a_n\}$ a sequence tale che $a_n > 0$ and $\frac{a_{n+1}}{a_n} \geq \frac{n}{n+1}$ for every $n \in \mathbb{N}$. Si dimostri che $\sum_{n=1}^{\infty} a_n$ diverges.

Tempo a disposizione: 2 ore and 45 minuti.

THEME 2

Exercise 1 [7 punti] Consider the function

$$f(x) = 1 - \sin(2 \arctan(|x|^3))$$

- i) determine the domain D , the sign, simmetries, i limits at the extremes of D and the asymptotes;
- ii) study the derivability, calcolarne the derivata, study the monotonicity, determine the points of extreme relative and absolute ; the study of the second derivative may be skipped .

iii) draw the qualitative graph .

Exercise 2 [6 punti] Compute the limit

$$\lim_{x \rightarrow 0^+} (1 - \sinh x)^{x^a}$$

as $a \in \mathbb{R}$, usando the form “ $\exp\{\log \dots\}$ ”.

Exercise 3 [4 punti] Trovare the zeros in \mathbb{C} di

$$(z^2 - z + 1)(z^3 + 4) = 0.$$

Exercise 4 [4+3 punti] Siano $\alpha \in \mathbb{R}$ fissato and

$$f_\alpha(t) := \frac{e^{2t} - 3e^t}{(e^t - 1)^\alpha}.$$

- i) Compute a primitive of f_α con $\alpha = 1$.
- ii) Determine for which $\alpha \in \mathbb{R}$ esiste finito $\int_0^1 f_\alpha(t) dt$.

Exercise 5 [6 punti] Study the convergence semplice and assoluta of the series

$$\sum_{n=1}^{+\infty} \frac{(4 \cos x)^n n}{n^2 + 1}$$

as $x \in [0, \pi]$.

Exercise Sia $\{a_n\}$ a sequence tale che $a_n > 0$ and $\frac{a_{n+1}}{a_n} \geq \frac{n}{n+1}$ for every $n \in \mathbb{N}$. Si dimostri che $\sum_{n=1}^{\infty} a_n$ diverges.

Tempo a disposizione: 2 ore and 45 minuti.

THEME 3

Exercise 1 [7 punti] Consider the function

$$f(x) = \sin(2 \arctan(|x|^5))$$

- i) determine the domain D , the sign, simmetries , i limits at the extremes of D and the asymptotes;
- ii) study the derivability, calcolarne the derivata, study the monotonicity, determine the points of extreme relative and absolute ; the study of the second derivative may be skipped .
- iii) draw the qualitative graph .

Exercise 2 [6 punti] Compute the limit

$$\lim_{x \rightarrow 0^+} (1 - \sin x)^{x^a}$$

as $a \in \mathbb{R}$, usando the form “ $\exp\{\log \dots\}$ ”.

Exercise 3 [4 punti] Trovare the zeros in \mathbb{C} di

$$(z^3 + 3)(z^2 + z + 2) = 0.$$

Exercise 4 [4+3 punti] Siano $\alpha \in \mathbb{R}$ fissato and

$$f_\alpha(t) := \frac{e^{2t} - 2e^t}{(e^t - 1)^\alpha}.$$

- i) Compute a primitive of f_α con $\alpha = 1$.
- ii) Determine for which $\alpha \in \mathbb{R}$ esiste finito $\int_0^1 f_\alpha(t) dt$.

Exercise 5 [6 punti] Study the convergence semplice and assoluta of the series

$$\sum_{n=1}^{+\infty} \frac{(4 \sin x)^n n}{n^2 + 2\sqrt{n}}$$

as $x \in [-\frac{\pi}{2}, \frac{\pi}{2}]$.

Exercise Sia $\{a_n\}$ a sequence tale che $a_n > 0$ and $\frac{a_{n+1}}{a_n} \geq \frac{n}{n+1}$ for every $n \in \mathbb{N}$. Si dimostri che $\sum_{n=1}^{\infty} a_n$ diverges.

Tempo a disposizione: 2 ore and 45 minuti.

THEME 4

Exercise 1 [7 punti] Consider the function

$$f(x) = 1 - \sin(2 \arctan(|x|^5))$$

- i) determine the domain D , the sign, simmetries, i limits at the extremes of D and the asymptotes;
- ii) study the derivability, calcolarne the derivata, study the monotonicity, determine the points of extreme relative and absolute ; the study of the second derivative may be skipped.
- iii) draw the qualitative graph .

Exercise 2 [6 punti] Compute the limit

$$\lim_{x \rightarrow 0^+} (1 + \sinh x)^{x^a}$$

as $a \in \mathbb{R}$, usando the form “ $\exp\{\log \dots\}$ ”.

Exercise 3 [4 punti] Trovare the zeros in \mathbb{C} di

$$(z^2 - z + 2)(z^3 + 2) = 0.$$

Exercise 4 [4+3 punti] Siano $\alpha \in \mathbb{R}$ fissato and

$$f_\alpha(t) := \frac{e^{2t} + 3e^t}{(e^t - 1)^\alpha}.$$

- i) Compute a primitive of f_α con $\alpha = 1$.
- ii) Determine for which $\alpha \in \mathbb{R}$ esiste finito $\int_0^1 f_\alpha(t) dt$.

Exercise 5 [6 punti] Study the convergence semplice and assoluta of the series

$$\sum_{n=1}^{+\infty} \frac{(3 \cos x)^n n}{n^2 + 2}$$

as $x \in [0, \pi]$.

Exercise Sia $\{a_n\}$ a sequence tale che $a_n > 0$ and $\frac{a_{n+1}}{a_n} \geq \frac{n}{n+1}$ for every $n \in \mathbb{N}$. Si dimostri che $\sum_{n=1}^{\infty} a_n$ diverges.

Tempo a disposizione: 2 ore and 45 minuti.

Appello of the 10.02.2020

THEME 1

Exercise 1 [7 punti] Consider the function

$$f(x) = \exp \left\{ \left| \frac{x}{x+1} \right| \right\}.$$

- i) Find the domain D , i limits at the extremes of D and the asymptotes;
- ii) study the derivability, calcolarne the derivata, study the monotonicity, determine the points of extreme relative and absolute ;
- iii) draw the qualitative graph .

Exercise 2 [5 punti] Study the convergence of the series

$$\sum_{k=1}^{\infty} 3^k \frac{k!}{k^k}.$$

Exercise 3 [5 punti] Solve in \mathbb{C} nella form preferita (algebrica, esponenziale, trigonometrica):

$$z^3 = \frac{1}{-\frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2}i}.$$

Exercise 4 [4+3 punti] Siano $\alpha \in \mathbb{R}$ and

$$f_{\alpha}(t) := \frac{e^{-2/t}}{3t^{\alpha}}.$$

- i) Compute a primitive of f_{α} con $\alpha = 3$.
- ii) Determine for which $\alpha \in \mathbb{R}$ esiste finito $\int_0^{+\infty} f_{\alpha}(t) dt$.

Exercise 5 [6 punti] Compute the seguente limit

$$\lim_{x \rightarrow 0^+} \frac{\sin(x - x^3) - \log(1 + \sinh x) + \alpha x^2}{x^3}$$

as $\alpha \in \mathbb{R}$.

Exercise Sia $\alpha \in [0, +\infty[$ and si definisca

$$F_{\alpha}(x) := \int_0^x t^{\alpha} e^{-t^2} dt, \quad x \geq 0.$$

Establish for which values of α one has che F_{α} is concave sull'interval $[1, +\infty[$. There are values $\alpha > 0$ so that F_{α} sia concave su $[0, +\infty[$?

Tempo a disposizione: 2 ore and 45 minuti.

THEME 2

Exercise 1 [7 punti] Consider the function

$$f(x) = \exp \left\{ \left| \frac{x+1}{x} \right| \right\}.$$

- i) Find the domain D , i limits at the extremes of D and the asymptotes;
- ii) study the derivability, calcolarne the derivata, study the monotonicity, determine the points of extreme relative and absolute ;

iii) draw the qualitative graph .

Exercise 2 [5 punti] Study the convergence of the series

$$\sum_{k=1}^{\infty} 4^k \frac{k!}{k^k}.$$

Exercise 3 [5 punti] Solve in \mathbb{C} nella form preferita (algebrica, esponenziale, trigonometrica):

$$z^3 = \frac{1}{\frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2}i}.$$

Exercise 4 [4+3 punti] Siano $\alpha \in \mathbb{R}$ and

$$f_{\alpha}(t) := \frac{2e^{-3/t}}{t^{\alpha}}.$$

- i) Compute a primitive of f_{α} con $\alpha = 3$.
- ii) Determine for which $\alpha \in \mathbb{R}$ esiste finito $\int_0^{+\infty} f_{\alpha}(t) dt$.

Exercise 5 [6 punti] Compute the seguente limit

$$\lim_{x \rightarrow 0^+} \frac{\sinh(x - x^3) - \log(1 + \sin x) + \alpha x^2}{x^3}$$

as $\alpha \in \mathbb{R}$.

Exercise Sia $\alpha \in [0, +\infty[$ and si definisca

$$F_{\alpha}(x) := \int_0^x t^{\alpha} e^{-t^2} dt, \quad x \geq 0.$$

Establish for which values of α one has che F_{α} is concave sull'interval $[1, +\infty[$. There are values $\alpha > 0$ so that F_{α} sia concave su $[0, +\infty[$? Tempo a disposizione: 2 ore and 45 minuti.

THEME 3

Exercise 1 [7 punti] Consider the function

$$f(x) = \exp \left\{ \left| \frac{x}{x-1} \right| \right\}.$$

- i) Find the domain D , i limits at the extremes of D and the asymptotes;
- ii) study the derivability, calcolarne the derivata, study the monotonicity, determine the points of extreme relative and absolute ;

iii) draw the qualitative graph .

Exercise 2 [5 punti] Study the convergence of the series

$$\sum_{k=1}^{\infty} 5^k \frac{k!}{k^k}.$$

Exercise 3 [5 punti] Solve in \mathbb{C} nella form preferita (algebrica, esponenziale, trigonometrica):

$$z^3 = \frac{1}{-\frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}i}.$$

Exercise 4 [4+3 punti] Siano $\alpha \in \mathbb{R}$ and

$$f_{\alpha}(t) := \frac{3e^{-2/t}}{t^{\alpha}}.$$

- i) Compute a primitive of f_{α} con $\alpha = 3$.
- ii) Determine for which $\alpha \in \mathbb{R}$ esiste finito $\int_0^{+\infty} f_{\alpha}(t) dt$.

Exercise 5 [6 punti] Compute the seguente limit

$$\lim_{x \rightarrow 0^+} \frac{\sin(x + x^3) - \log(1 + \sinh x) + \alpha x^2}{x^3}$$

as $\alpha \in \mathbb{R}$.

Exercise Sia $\alpha \in [0, +\infty[$ and si definisca

$$F_{\alpha}(x) := \int_0^x t^{\alpha} e^{-t^2} dt, \quad x \geq 0.$$

Establish for which values of α one has che F_{α} is concave sull'interval $[1, +\infty[$. There are values $\alpha > 0$ so that F_{α} sia concave su $[0, +\infty[$?

Tempo a disposizione: 2 ore and 45 minuti.

THEME 4

Exercise 1 [7 punti] Consider the function

$$f(x) = \exp \left\{ \left| \frac{x-1}{x} \right| \right\}.$$

- i) Find the domain D , i limits at the extremes of D and the asymptotes;

- ii) study the derivability, calcolarne the derivata, study the monotonicity, determine the points of extreme relative and absolute ;
- iii) draw the qualitative graph .

Exercise 2 [5 punti] Study the convergence of the series

$$\sum_{k=1}^{\infty} 6^k \frac{k!}{k^k}.$$

Exercise 3 [5 punti] Solve in \mathbb{C} nella form preferita (algebrica, esponenziale, trigonometrica):

$$z^3 = \frac{1}{\frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}i}.$$

Exercise 4 [4+3 punti] Siano $\alpha \in \mathbb{R}$ and

$$f_{\alpha}(t) := \frac{e^{-3/t}}{2t^{\alpha}}.$$

- i) Compute a primitive of f_{α} con $\alpha = 3$.
- ii) Determine for which $\alpha \in \mathbb{R}$ esiste finito $\int_0^{+\infty} f_{\alpha}(t) dt$.

Exercise 5 [6 punti] Compute the seguente limit

$$\lim_{x \rightarrow 0^+} \frac{\sinh(x + x^3) - \log(1 + \sin x) + \alpha x^2}{x^3}$$

as $\alpha \in \mathbb{R}$.

Exercise Sia $\alpha \in [0, +\infty[$ and si definisca

$$F_{\alpha}(x) := \int_0^x t^{\alpha} e^{-t^2} dt, \quad x \geq 0.$$

Establish for which values of α one has che F_{α} is concave sull'interval $[1, +\infty[$. There are values $\alpha > 0$ so that F_{α} sia concave su $[0, +\infty[$?

Tempo a disposizione: 2 ore and 45 minuti.

Appello of the 06.07.2020 - Modalità telematica (causa COVID)

THEME 1

Exercise 1 [6 punti] Consider the function

$$f(x) = |(x+3)\log(x+3)|, \quad x \in D =]-3, +\infty[.$$

(i) Compute

$$\lim_{x \rightarrow -3^+} f(x), \quad \lim_{x \rightarrow +\infty} f(x).$$

(ii) Compute the first derivative of the function f , study the monotonicity intervals and draw the graph of f .

Exercise 2 [6 punti] Find the solutions of the equation

$$z^3 = 8i$$

in algebraic and exponential form (or trigonometric), and draw them on the Gauss plane .

Exercise 3 [6 punti] Study the convergence of the series

$$\sum_{n=1}^{\infty} \frac{(1+n^2) \log n}{n^4}.$$

Exercise 4 [6 punti] Compute

$$\int_0^{+\infty} e^{-\sqrt{2x}} dx.$$

Exercise 5 [6 punti] Compute the limit

$$\lim_{x \rightarrow +\infty} x^{\frac{4}{3}} \left(\sqrt[3]{x+2} - \sqrt[6]{x^2-1} \right)^2.$$

Tempo a disposizione: 1 ore and 30 minuti.

Appello of the 14.09.2020 - Modalità telematica (causa COVID)

THEME 1

Exercise 1 [6 punti] Consider the function

$$f(x) = \arctan \left(\frac{x+1}{x-1} \right), \quad x \in (1, \infty).$$

(i) Individuarne the asymptotes.

(ii) If ne determini the monotonicity .

Exercise 2 [6 punti] Consider the complex number $z = \sqrt{3} - i$.

(i) Scrivere lo in exponential form .

(ii) Compute the real part of z^6 .

Exercise 3 [6 punti] Establish the convergence semplice and assoluta of the series

$$\sum_{n=1}^{\infty} (-1)^n \frac{n}{n^2 + 1}.$$

Exercise 4 [6 punti] Compute the limit

$$\lim_{x \rightarrow 0^+} \frac{\log(1 + \sinh x) - \sin x}{x^2}.$$

Exercise 5 [6 punti] Consider the generalized integral

$$\int_1^{\infty} \log\left(\frac{x^\alpha}{x^\alpha + 1}\right) dx.$$

- (i) Compute the integral for $\alpha = 2$.
- (ii) Establish for which $\alpha \in [0, \infty)$ it converges.

Tempo a disposizione: 1 ore and 30 minuti.

6 Esercizi scelti dai temi d'esame di anni passati

Functions .

1) (20.02.2013) Given the function

$$f(x) = x \left| 3 + \frac{1}{\log(2x)} \right|,$$

- (a) determine the domain, calcolarne the limits at the extremes and determine asymptotes;
 - (b) study the prolongabilità at the extremes of the domain and the derivability;
 - (c) compute f' and determine the monotonicity intervals and the points of extreme (maximum and minimum) relative and absolute of f ;
 - (d) compute the main limits of f' ;
 - (e) draw a qualitative graph of f (the study of the concavity and of the convexity is not required).
- 2) (3.02.2014) Consider the function

$$f(x) = \arctan \left(\frac{2x}{\log|x| - 1} \right).$$

- 1) Determine the domain and discuss the simmetria and the sign of f .
- 2) Compute the main limits of f , determine the asymptotes and discuss brevemente the continuity.
- 3) Compute f' and determine the monotonicity intervals and the points of extreme of f .
- 4) Compute the main limits of f' and study the derivability of f in $x = 0$.
- 5) Draw a graph of f the study of the second derivative may be skipped.

3) (26.01.2015) Consider the function

$$f(x) = |x+1| e^{\frac{-1}{|x+3|}}.$$

- (a) Determine the domain D of f ; determine the limits of f at the extremes of D and the asymptotes; study the continuity and the prolungamenti for continuity;
- (b) study the derivability, compute the derivative and study the monotonicity of f ; determine the points of extreme relative and absolute and compute the main limits of f' ;
- (c) draw a qualitative graph of f .

Series

1) (16.09.2013) Discutere, for all values of the parameter $\alpha \in \mathbb{R}$, the convergence of the series

$$\sum_{n=1}^{\infty} \log \left(n \left(e^{\frac{1}{n}} - 1 \right) - \frac{\alpha}{n} \right).$$

2) (26.01.2015) Determine all the $x \in \mathbb{R}$ such that the series

$$\sum_{n=2}^{\infty} \frac{\log n}{n-1} (x-2)^n$$

converga, resp. converga assolutamente.

- 3) (25.01.2016) Determine all the $x \in \mathbb{R}$ such that the series

$$\sum_{n=5}^{+\infty} \frac{(\log(x-3))^n}{n-1}$$

converga, resp. converga assolutamente.

Limits

- 1) (20.02.2013) Compute the limit

$$\lim_{x \rightarrow 0^+} \frac{x^{7/2} \log^2 x - 1 + \sin x^2 + \cos(1 - e^{\sqrt{2}x})}{\sinh x - x^\alpha}$$

for all values of the parameter $\alpha > 0$.

- 2) (3.02.2014) Compute the limit

$$\lim_{x \rightarrow 0^+} \frac{\sinh x^\alpha - \cos(\sqrt{x}) \log(1 + \sin x)}{\log \cos 2x + x^3 \log x}$$

for all values of the parameter $\alpha > 0$.

- 3) (20.02.2015) (a) Compute the order diinfinitesimal 1 di

$$e^{x-x^2} - \cos(\alpha x) - \sin x$$

for $x \rightarrow 0$ as $\alpha \in \mathbb{R}$;

(b) compute the limit

$$\lim_{x \rightarrow 0} \frac{e^{x-x^2} - \cos(\alpha x) - \sin x}{\sinh x - \log(1 + \sin x)}$$

as $\alpha \in \mathbb{R}$.

Esercizi sui numeri complessi

- 1) (7.02.2012) Solve the equation

$$i \operatorname{Re} z + z^2 = |z|^2 - 1$$

and draw the solutions on Gauss plane .

- 2) (23.02.2012) Write in algebraic form the zeros of the polynomial

$$(z^2 + iz + 2)(z^3 - 8i).$$

- 3) (18.09.2012) Express in algebraic form the solutions of the equation

$$z^6 - iz^3 + 2 = 0$$

and rappresentarle on Gauss plane .

4) (5.02.2013) Compute tutte the solutions $z \in \mathbb{C}$ of the equation

$$\left(\frac{2z+1}{2z-1}\right)^3 = 1,$$

scriverele in algebraic form and rappresentarle in the Gauss plane .

5) (15.07.2013) Compute tutte the solutions $z \in \mathbb{C}$ of the equation

$$z^5 = -16\bar{z}$$

writing them prima in trigonometric form /esponenziale and in algebraic form; draw them infine on Gauss plane .

6) (15-07.2014) Express in trigonometric form the solutions of the equation

$$\frac{z^4}{z^4 + 1} = 1 - \frac{i}{\sqrt{3}}, \quad z \in \mathbb{C}$$

and draw them in the Gauss plane .

7) (20.02.2015) Si risolva the inequality

$$\operatorname{Re}((z+i)^2) \leq \operatorname{Im}(i(\bar{z}-2i)^2) \quad (1)$$

and se ne disegni the insieme of the solutions in the Gauss plane .

8) (16.07.2015) Si risolva the equation

$$\left(\frac{1}{18} - \frac{i\sqrt{3}}{18}\right)\bar{z}^2 = 1,$$

disegnandone the solutions in the Gauss plane .

9) (15.02.2016) Determine tutte the solutions of the 'equation

$$\bar{z}^2 = 2\operatorname{th}ez, \quad z \in \mathbb{C},$$

writing them in algebraic form and rappresentandole on Gauss plane .

10) (11.07.2016) Solve in the Gauss plane the equation

$$2\bar{z}^3 = 3i,$$

rappresentandone the solutions in algebraic form.

Integrali

1) (7.02.2012) Compute the integral

$$\int_0^8 e^{\sqrt[3]{x}} dx.$$

2) (23.02.2012) Given the function

$$f(x) = \frac{2e^x + 1}{e^{2x} + 2e^x + 2},$$

- (a) compute a primitiva;
 (b) si provi that the generalized integral $\int_0^{+\infty} f(x) dx$ and converging and lo si calcoli.
 3) (17.07.2012) (a) Compute the order diinfinito for $x \rightarrow 3$ of the function

$$g(x) = \frac{x}{9 - x^2};$$

b) dire for which $\alpha \geq 0$ converges the integral

$$I = \int_0^3 \frac{x}{(9 - x^2)^\alpha} dx;$$

c) calcolarlo for $\alpha = \frac{1}{2}$.

4) (5.02.2013) Compute the integral

$$\int_{\log 8}^{+\infty} \frac{\sqrt{e^x + 1}}{e^x - 3} dx.$$

5) (20.02.2013) Compute the integral

$$\int_{\frac{2}{\pi}}^{+\infty} \frac{1}{x^4} \sin \frac{1}{x} dx$$

6) (15.07.2013) a) Study the convergence of the generalized integral

$$\int_0^{+\infty} \frac{e^{2\alpha x} - 1}{e^{2x} + 1} dx$$

as $\alpha \in \mathbb{R}$.

b) Compute the integral for $\alpha = 1/2$.

7) (3.02.2014) Study the convergence of the integral

$$\int_0^{+\infty} \frac{\arctan x}{(x+2)^{\frac{\alpha-1}{2}} (4+x)^{2\alpha}} dx$$

as $\alpha \in \mathbb{R}$ and calcolarlo for $\alpha = 1$.

8) (15.07.2014) Trovare for which $\alpha \in \mathbb{R}$ converges the integral

$$\int_0^{+\infty} \frac{1}{x^\alpha (3 + 2\sqrt{x} + x)} dx$$

and calcolarlo for $\alpha = 1/2$.

9) (12.09.2014) Determine the $\alpha \in \mathbb{R}$ for the quali the integral

$$\int_0^4 \frac{\sqrt{x}}{(4-x)^\alpha} dx$$

converges and calcolarlo for $\alpha = 1/2$.

10) (25.01.2016) Compute the integral

$$\int_0^{1/2} (\arcsin 2x)^2 dx$$

11) (11.07.2016) Establish for which $\alpha \in \mathbb{R}$ the seguente integral is converging

$$\int_0^{\pi/8} \frac{\sin 2x}{|\log(\cos 2x)|^\alpha \cos 2x} dx$$

and calcolarne the value for $\alpha = 1/2$.

7 Ulteriori esercizi (a cura di C. Sartori)

FUNZIONI

Exercise. Determine, as $\lambda > 1$ the numero disolutions of the equation

$$\lambda^x = x^\lambda.$$

Soluzione. The equation (that ha the soluzione λ) is equivalente a

$$x \log \lambda = \lambda \log x.$$

Posto $f(x) = x \log \lambda$, $g(x) = \lambda \log x$, one has $f'(x) = \log \lambda$, $g'(x) = \frac{\lambda}{x}$ and hence le two functions they are tangenti if

$$\begin{cases} x \log \lambda = \lambda \log x \\ \log \lambda = \frac{\lambda}{x}. \end{cases}$$

Si ricava $\lambda = \lambda \log x$ that is, $x = e$ and hence $\log \lambda = \frac{\lambda}{e}$ from which $\lambda = e$. The function $\log x$ is tangent alla retta $y = \frac{\log \lambda}{\lambda} x$ if $\lambda = e$. the coefficiente angolare of the retta ha a maximum for $\lambda = e$ and hence confrontando the graph of $\log x$ con quello of the retta $y = \frac{\log \lambda}{\lambda} x$ si ottengono always two solutions $\forall \lambda > 1$. Per $\lambda = e$ one has a sola soluzione.

Exercise Given the function $f : \mathbb{R} \rightarrow \mathbb{R}$ defined da

$$f(x) = 3x^4 + 4(2a - a^2)x^3 - 12a^3x^2 + a^6,$$

dove $a > 0$ is a parameter fissato. Determine

a) i points of maximum and minimum of f and the values of f in tali punti;

$x = -2a, a^2$ points of minimum ; $x = 0$ point of maximum; $f(-2a) = -16a^4 - 16a^5 + a^6$, $f(a^2) = -a^8 - 4a^7 + a^6$ $f(0) = a^6$.

b) i values of a so that the equation $f = 0$ ha 2 zeros positivi;

Basta imporre $f(a^2) < 0$ that implica $a > -2 + \sqrt{5}$;

c) i values of a so that the equation $f = 0$ ha non piu' dia zero negative ;

Basta imporre $f(-2a) \geq 0$ that implica $a \geq 8 + \sqrt{80}$.

d) i values of $a \geq 0$ so that f is convex.

$$a = 0$$

Exercise. Determine, as $\lambda \in \mathbb{R}$ the numero disolutions of the equation

$$\frac{1}{10(1+x^2)} + |1 - \sqrt{|x|}| = \lambda.$$

Soluzione. Studio the function $f(x) = \frac{1}{10(1+x^2)} + |1 - \sqrt{|x|}|$ is pari and hence basta studiarla for $x \geq 0$. One has $f(0) = \frac{11}{10}$, $\lim_{x \rightarrow +\infty} = +\infty$ e

$$f'(x) = \begin{cases} \frac{-2x}{10(1+x^2)^2} - \frac{1}{2\sqrt{x}}, & 0 < x < 1 \\ \frac{-2x}{10(1+x^2)^2} + \frac{1}{2\sqrt{x}}, & x > 1. \end{cases}$$

f is decreasing in $(0, 1)$ and increasing in $(1, +\infty)$, hence $x = 0$ is point direlative maximum and $x = 1$ is point of absolute minimum. The graph is porta alle solutions

$$\begin{cases} \lambda < \frac{1}{20} & \text{ness a soluzione} \\ \lambda = \frac{1}{20} & 2 \text{ solutions} \\ \frac{1}{20} < \lambda < \frac{11}{10} & 4 \text{ solutions} \\ \lambda = \frac{11}{10} & 3 \text{ solutions} \\ \lambda > \frac{11}{10} & 2 \text{ solutions} . \end{cases}$$

Exercise Given the function

$$f(x) = 4x^3 - 4ax^2 + a^2x - 1,$$

determine for which values of $a > 0$

- a) $f(x)$ ha esattamente three zeros;
- b) tali zeros they are all positivi.

Soluzione.

a)

$$f'(x) = 12x^2 - 8ax + a^2 = 0 \iff x = \frac{a}{2}, x = \frac{a}{6}.$$

$x = \frac{a}{2}$ is point of maximum and $x = \frac{a}{6}$ is point of minimum. In order to find three zeros one must assume $f(\frac{a}{2}) < 0 < f(\frac{a}{6})$, that is verified if and only if $a > \sqrt[3]{2}$.

- b) Since $f(0) = -1 < 0$ for $0 < x < \frac{a}{6}$ there is a zero, as well as there is a zero in $(\frac{a}{6}, \frac{a}{2})$ and, finally a zero $x > \frac{a}{2}$, $\lim_{x \rightarrow +\infty} f(x) = +\infty$.

Exercise. Given the function

$$f_a(x) = x^a - ax^2, \quad a > 0,$$

compute $\sup\{f_a(x), x \geq 0\}$ and $\inf\{f_a(x), x \geq 0\}$, specifying if they are maximum o minimum .

Soluzione.

$$\begin{aligned} a > 2 &\Rightarrow \lim_{x \rightarrow +\infty} f_a(x) = +\infty = \sup\{f_a(x), x \geq 0\}; \\ a \leq 2 &\Rightarrow \lim_{x \rightarrow +\infty} f_a(x) = -\infty = \inf\{f_a(x), x \geq 0\}. \end{aligned}$$

One has $f'_a(x) = ax^{a-1} - 2ax = 0 \iff x = 0, 2^{\frac{1}{a-2}}$ if $a \neq 2$. $2^{\frac{1}{a-2}}$ is of minimum if $a > 2$, of maximum if $a < 2$. Therefore

$$a > 2 \Rightarrow \min\{f_a(x), x \geq 0\} = 2^{\frac{a}{a-2}} - a2^{\frac{2}{a-2}};$$

$$a < 2 \Rightarrow \max\{f_a(x), x \geq 0\} = 2^{\frac{a}{a-2}} - a2^{\frac{2}{a-2}};$$

$$a = 2 \Rightarrow \max\{f_2(x), x \geq 0\} = 0.$$

Exercise. Study, as $\lambda \in \mathbb{R}$, the number of solutions of the equation

$$-e^x + e^4|x - 1| = \lambda.$$

Sol. Study the function

$$f(x) = -e^x + e^4|x - 1|.$$

$$\text{Dom } f = \mathbb{R}, \lim_{x \rightarrow +\infty} f(x) = -\infty, \lim_{x \rightarrow -\infty} f(x) = +\infty.$$

$$f'(x) = \begin{cases} -e^x + e^4, & \text{for } x > 1 \\ -e^x - e^4, & \text{for } x < 1. \end{cases}$$

Find a point of maximum in $(4, 2e^4)$ and a point of minimum (angle) in $(1, -e)$. Therefore

$$\begin{cases} \lambda > 2e^4, \lambda < -e & 1 \text{ sol.}, \\ -e < \lambda < 2e^4 & 3 \text{ sol.}, \\ \lambda = -e, 2e^4 & 2 \text{ sol..} \end{cases}$$

Exercise. Sia

$$f(x) = \ln(x+4) + \frac{x+8}{x+4}.$$

- Compute the intervals of concavity and convexity of f on its domain.
- Find the maximum interval A containing -3 where f is invertible.
- Let g the inverse function of f restricted to A . Compute $g'(f(-3))$.

SOL. $\text{Dom } f = \{x > -4\}$. $f'(x) = x/(4+x)^2$, $f''(x) = (4-x)/(4+x)^3$. One has $f''(x) > 0$ for $-4 < x < 4$ and the function is convex, for $x > 4$ concave. The maximal neighbourhood of -3 in which f is monotonic (decreasing) and hence invertible is $-4 < x < 0$. One has $f(-3) = 5$, and

$$g'(f(-3)) = \frac{1}{f'(-3)} = -\frac{1}{3}.$$

FUNZIONI INTEGRALI

Exercise. Study the convexity and concavity of the function

$$F(x) = \int_2^x g(\sin t) dt, \quad x \in \mathbb{R},$$

dove g is a function differentiable in \mathbb{R} and tale che $g'(x) < 0$.

Soluzione One has

$$F'(x) = g(\sin x), \text{ e } F''(x) = g'(\sin x) \cos x,$$

from which

$$F''(x) > 0 \iff \cos x < 0 \iff \frac{\pi}{2} + 2K\pi < x < \frac{3\pi}{2} + 2K\pi$$

for K intero; in the union of tali intervalli F is convex, and in the complementare is concave.

Exercise. Study the function

$$F(x) = \int_0^x \frac{(t+1)(3-t)}{\arctan(1+t^2)} dt,$$

specifying, in particolare, the intervals dicrescenza and decrescenza.

Compute $\lim_{x \rightarrow +\infty} F(x)$ and $\lim_{x \rightarrow -\infty} F(x)$ and tracciare a qualitativa graph .

Soluzione. One has $F'(x) = \frac{(x+1)(3-x)}{\arctan(1+x^2)}$ and $F'(x) > 0 \iff -1 < x < 3$. $\lim_{x \rightarrow +\infty} F'(x) = -\infty$ and $\lim_{x \rightarrow -\infty} F'(x) = -\infty$, from which si ricava $F'(x) < -1$ for $|x| > M$ and hence $\lim_{x \rightarrow +\infty} F(x) = -\infty$ and $\lim_{x \rightarrow -\infty} F(x) = +\infty$.

LIMITI

Exercise Compute the seguenti limits (the terzo as $\alpha \in \mathbb{R}$),

$$\begin{aligned} \lim_{x \rightarrow 3} \frac{x^6 - 3^6}{x^8 - 3^8} &= 1/12, \quad \lim_{n \rightarrow +\infty} \left(1 + \frac{1}{n^n}\right)^{n!} = 1, \\ \lim_{x \rightarrow 0^+} \frac{\sqrt{x^\alpha + x} + \sqrt{x}}{x} &= \lim_{x \rightarrow 0^+} \frac{\sqrt{x^\alpha + x} + \sqrt{x}}{x} \\ &= \begin{cases} +\infty & \text{if } \alpha < 2/3 \\ 1 & \text{if } \alpha = 2/3 \\ 0 & \text{if } \alpha > 2/3 \end{cases} \end{aligned}$$

8 Soluzioni of the Temi 1 of the prove scritte of the quattro anni precedenti

Appello of the 23.01.2017

THEME 1

Exercise 1 Compute the integral

$$\int_{\log(3)}^2 \frac{e^x}{e^{2x} - 4} dx$$

Solution. One has

$$\begin{aligned} \int_{\log(3)}^2 \frac{e^x}{e^{2x} - 4} dx &= (\text{setting } e^x = t, \text{ so that } dx = dt/t) \int_3^{e^2} \frac{1}{t^2 - 4} dt \\ &= -\frac{1}{4} \int_3^{e^2} \frac{1}{t+2} - \frac{1}{t-2} dt = \frac{1}{4} \ln \frac{|t-2|}{|t+2|} \Big|_3^{e^2} \\ &= \frac{1}{4} \left[\log 5 \frac{e^4 - 2}{e^4 + 2} \right] = \frac{1}{2} \left(\operatorname{settanh} \frac{3}{2} - \operatorname{settanh} \frac{e^2}{2} \right). \end{aligned}$$

Exercise 2 Solve the inequality

$$|2\bar{z}^2 - 2z^2| < 3$$

and draw the solutions on Gauss plane .

Solution. One has

$$2\bar{z}^2 - 2z^2 = 4(\bar{z} - z)(\bar{z} + z) = (\text{setting } z = x + iy) 8ixy,$$

from which

$$|2\bar{z}^2 - 2z^2| = 8|xy|.$$

The soluzione is hence

$$\{z = x + iy \in \mathbb{C} : |xy| < \frac{3}{8}\},$$

rappresentata in the picture 1.



Figura 1: Solutions of exercise 2 (Theme 1).

Exercise 3 Study the convergence of the series

$$\sum_{n=1}^{+\infty} n^2 (\cos(1/n) - 1 + \sin(1/2n^\alpha))$$

for all values of the parameter $\alpha > 0$.

Solution. Dagli sviluppi di Mac Laurin di $\cos x$ and of $\sin x$ one has, for $n \rightarrow +\infty$,

$$\cos(1/n) - 1 + \sin(1/2n^\alpha) = -\frac{1}{2n^2} + \frac{1}{4! \cdot n^4} + o\left(\frac{1}{n^4}\right) + \frac{1}{2n^\alpha} - \frac{1}{3!8n^{3\alpha}} + \frac{1}{5!32n^{5\alpha}} + o\left(\frac{1}{n^{5\alpha}}\right),$$

so that the general term of the series, for $n \rightarrow +\infty$, is asymptotic to

$$\begin{cases} (\text{if } \alpha < 2) & \frac{n^2}{2n^\alpha} \\ (\text{if } \alpha = 2) & 1/24n^2 \\ (\text{if } \alpha > 2) & -1/2 \end{cases}$$

and hence ha sign definitively constant for $n \rightarrow +\infty$. If $\alpha \neq 2$ the general term of the series is not infinitesimal 1 and hence the series diverges ($\rightarrow -\infty$). Per $\alpha = 2$ the series converges.

Exercise 4 Consider the function

$$f(x) := \arcsin \frac{|x| - 4}{2 + x^2}.$$

- i) Determine the domain D of f , its simmetries and study the sign; determine the limits of f at the extremes of D ;
- ii) study the derivability, compute the derivative and study the monotonicity of f ; determine the points of extreme relative and absolute and compute the main limits of f' ;
- iii) draw a qualitative graph of f .

Solution. (i) The function is pari. $D = \{x \in \mathbb{R} : -1 \leq \frac{|x|-4}{2+x^2} \leq 1\}$. The inequality $\frac{|x|-4}{2+x^2} \leq 1$ equivale a $|x| - 6 - x^2 \leq 0$, that is always verificata, while $\frac{|x|-4}{2+x^2} \geq -1$ equivale a $x^2 + |x| - 2 \geq 0$, that is verificata for $x \leq -1$ and $x \geq 1$. Therefore $D = [1, +\infty[\cup]-\infty, -1]$. D'ora in assumeremo always $x \geq 0$. The function is continuous in D , $f(1) = \arcsin(-1) = -\pi/2$ and $\lim_{x \rightarrow +\infty} f(x) = \arcsin 0 = 0$, horizontal asymptote. The sign of f is dato dal sign of the argument of the arcoseno, so that $f(x) \geq 0$ if and only if $x - 4 \geq 0$ and hence $x \geq 4$.

(ii) In D si posthey are applicare le regole diderivazione if the argument of the arcoseno is diverso da ± 1 , that is, for $x > 1$. Per tali x one has

$$f'(x) = \frac{x^2 + 2 - 2x(x-4)}{(2+x^2)^2} \frac{1}{\sqrt{1 - \left(\frac{x-4}{2+x^2}\right)^2}} = \frac{-x^2 + 8x + 2}{(1+2x^2)\sqrt{2x^2+x-3}},$$

from which one deduces that $f'(x) \leq 0$ if and only if $-x^2 + 8x + 2 \leq 0$, for $x > 1$, that is, for $1 < x < 4+3\sqrt{2}$, that therefore is the point di absolute maximum, while $x = 1$ is the point of absolute minimum. One has

$$\lim_{x \rightarrow 1^+} f'(x) = +\infty,$$

grafici/2017-I-T1f.pdf

Figura 2: the graph of f (Theme 1).

so that the graph of f , rappresentato nella figura 2, ha tangent verticale in $(1, \pi/2)$.

Exercise 5 Study the convergence of the generalized integral

$$\int_0^{+\infty} \frac{|\arctan(x-1)| \arctan x}{|1-x^2|^\alpha (\sinh \sqrt{x})^\beta} dx$$

as $\alpha, \beta \in \mathbb{R}$.

Solution. The integranda $f(x)$ is continuous in $]0, 1[\cup]1, +\infty[$, so that bisogna study the convergence of the integral separatamente for $x \rightarrow 0^+$, for $x \rightarrow 1$ and for $x \rightarrow +\infty$.

Per $x \rightarrow 0$,

$$f(x) \sim \frac{x \arctan 1}{x^{\beta/2}} = \arctan 1 \frac{1}{x^{\frac{\beta}{2}-1}},$$

and hence the integral converges in 0 if and only if $\beta < 4$.

Per $x \rightarrow 1$,

$$f(x) \sim \frac{\arctan 1 |x-1|}{|x-1|^\alpha |x+1|^\alpha (\sinh \sqrt{2})^\beta} = \frac{\arctan 1}{2^\alpha (\sinh \sqrt{2})^\beta} \frac{1}{|x-1|^{\alpha-1}},$$

and hence the integral converges in 1 if and only if $\alpha < 2$.

Per $x \rightarrow +\infty$, if $\beta > 0$

$$f(x) \leq \frac{\pi^2}{4 (\sinh \sqrt{x})^\beta} \leq \frac{\pi^2}{2^{(2-\beta)} e^{(\beta \sqrt{x})}}.$$

Quest'ultima expression is $o(1/x^2)$ for $x \rightarrow +\infty$ and hence converges.

If $\beta = 0$,

$$f(x) \sim \pi^2/4x^{2\alpha},$$

hence converges if $\alpha > 1/2$. If $\beta < 0$,

$$f(x) \sim \pi^2 e^{-\beta/2}/2^{2-\beta} > 1/x$$

for $x \rightarrow +\infty$ and hence the integral diverges. In sintesi, the integral converges if $\alpha < 2$ and $0 < \beta < 4$ o if $\beta = 0$ and $1/2 < \alpha < 2$.

Exercise . Sia I a interval chiuso and limitato and sia $f : I \rightarrow \mathbb{R}$ a function continuous and tale che $f(x) \in I$ for every $x \in I$. Dimostrare that esiste almeno a $x \in I$ tale che $f(x) = x$.

Solution. Consideriamo the function $g(x) = f(x) - x$, that vogliamo dimostrare that si annulla in almeno a point of $I := [a, b]$. If $g(a), g(b) \neq 0$ allora necessariamente $g(a) > 0$ and $g(b) < 0$, so that for the teorema degli zeros esiste $\bar{x} \in]a, b[$ tale che $g(\bar{x}) = 0$.

Appello of the 13.02.2017

THEME 1

Exercise 1 Consider the function

$$f(x) := \log |x^2 - 2x - 3|.$$

- i) Determine the domain D and study the sign of f ; determine the limits of f at the extremes of D and the asymptotes;
- ii) study the derivability, compute the derivata, study the monotonicity and determine the points of extreme relative and absolute of f ;
- (iii) compute f'' and study the concavity and the convexity of f ;
- (iv) draw a qualitative graph of f .

Soluzione. i) Clearly $D = \{x \in \mathbb{R} : x^2 - 2x - 3 \neq 0\} = \mathbb{R} \setminus \{-1, 3\}$. Per the sign abbiamo

$$f(x) \geq 0, \iff |x^2 - 2x - 3| \geq 1, \iff x^2 - 2x - 3 \leq -1, \vee x^2 - 2x - 3 \geq +1.$$

Abbiamo che $x^2 - 2x - 2 \leq 0$ if and only if $x_0 := 1 - \sqrt{3} \leq x \leq 1 + \sqrt{3} =: x_1$ and $x^2 - 2x - 4 \geq 0$ if and only if $x \leq 1 - \sqrt{5} =: x_2$ oppure $x \geq 1 + \sqrt{5} =: x_3$. Therefore $f(x) \leq 0$ if and only if x appartiene ad uno of the two intervals $[x_2, x_0]$ and $[x_1, x_3]$. As for i limits, one has:

and chiaro che $x^2 + 3x - 4 \rightarrow +\infty$ for $x \rightarrow \pm\infty$, cosicché $\lim_{x \rightarrow \pm\infty} f(x) = +\infty$. Yet non ci they are asymptotes poiché, for $x \rightarrow \pm\infty$,

$$\frac{f(x)}{x} = \frac{\log(x^2 - 2x - 3)}{x} \sim \frac{\log x^2}{x} = \frac{2 \log |x|}{x} \rightarrow 0,$$

essendo $\log|x| = o(x)$ for $x \rightarrow \pm\infty$. Per $x \rightarrow -1, 3$ one has always che $|x^2 + 3x - 4| \rightarrow 0+$ hence in ogni case $f(x) \rightarrow -\infty$ so that one has the asymptotes verticali $x = -1, 3$.

ii) The function is superposition dicontinuous functions ove definite, hence is continuous on tutto the proprio domain. Furthermore, is superposition didifferentiable functions, eccetto quando $x^2 + 3x - 4 = 0$, that for ò they are punti that non appartengono al domain of f : si conclude che f is differentiable in the proprio domain. Since $(\log|y|)' = \frac{1}{y}$ one has immediately che

$$f'(x) = \frac{2x - 2}{x^2 - 2x - 3}.$$

Let us study the sign of f' . the sign of the denominator is positive for $x < -1$ oppure $x > 3$. the numerator is positive for $x > 1$. We deduce the tabella seguente:

	$-\infty$	-1	-1	1	3	3	$+\infty$
$\operatorname{sgn}(2x - 2)$	—	—	+	+	+	+	+
$\operatorname{sgn}(x^2 - 2x - 3)$	+	—	—	—	+	+	+
$\operatorname{sgn} f'$	—	+	—	—	+	+	+
f	↘	↗	↘	↗	↗	↗	↗

I punti $x = -1, 3$ non appartengono al domain, while $x = 1$ is a localmaximum stretto. There are no nán massimi nán minimi globali essendo f il limitata sia inferiormente that superiormente.

iii) Clearly f' is differentiable ove defined in quanto rational function: we have che

$$f''(x) = \frac{-2x^2 + 4x - 10}{(x^2 - 2x - 3)^2}.$$

Therefore $f'' \geq 0$ if and only if $2x^2 - 4x - 10 \leq 0$, that is, mai. One concludes that $f'' < 0$ ovunque (where defined) so that the function is concave in ciascuno degli intervals that compongono the suo domain.

iv) the graph of f is rappresentato figura 3.

Exercise 2 Study the convergence of the series

$$\sum_{n=1}^{+\infty} \frac{1}{2^n} \frac{n^n}{n!}.$$

Soluzione. The series is clearly a termini with constant sign . one can applicare the criterio asymptotic of the rapporto. Detto a_n the general term, one has

$$\frac{a_{n+1}}{a_n} = \frac{1}{2} \frac{(n+1)^{n+1}}{(n+1)!} \frac{n!}{n^n} = \frac{1}{2} \frac{(n+1)^n}{n^n} = \frac{1}{2} \left(1 + \frac{1}{n}\right)^n \rightarrow \frac{1}{2}e > 1.$$

Hence the series diverges.

Exercise 3 Given

$$f(z) = \frac{2 + iz}{iz + 1},$$

determine the domain and determine all the $z \in \mathbb{C}$ tali che $f(z) = z$. Express tutte the solutions in algebraic form.

grafici/2017-II-T1f.pdf

Figura 3: the graph of f (Theme 1).

Soluzione. Perché the fraction sia defined occorre che $iz + 1 \neq 0$, that is, che $z \neq -\frac{1}{i} = \frac{i}{-i \cdot i} = i$. Ora, for $z \neq i$,

$$f(z) = z \iff 2 + iz = z(iz + 1) \iff iz^2 + (1 - i)z - 2 = 0.$$

Thisis un'equation disecondo grado a coefficienti complessi, and the formula risolutiva tradizionale funziona allo stesso modo (pur diintendere the root come radice complessa). One has hence

$$\begin{aligned} z_{1,2} &= \frac{i - 1 + \sqrt{1 - 1 - 2i + 8i}}{2i} = \frac{i - 1 + \sqrt{6i}}{2i} = \frac{i - 1 \pm \sqrt{6}e^{i\frac{\pi}{4}}}{2i} \\ &= \frac{1}{2} + \frac{i}{2} \pm \frac{\frac{\sqrt{12}}{2}(1+i)}{2i} = \frac{1 \pm \sqrt{3}}{2} + \frac{1 \mp \sqrt{3}}{2}i \end{aligned}$$

Exercice 4 Compute the limit

$$\lim_{x \rightarrow 0^+} \frac{\arctan x - \sin x + x^{\frac{10}{3}} \log x}{x^\alpha (1 - \cos^2 x)}$$

as $\alpha > 0$.

Soluzione. Osservato che, in virtù of the notable limit $\lim_{x \rightarrow 0^+} x^\gamma \log x = 0$ for every $\gamma \in \mathbb{R}$, one has subito that si tratta of a indeterminate form of the tipo $\frac{0}{0}$. Determiniamo the termini principali col metodo degli sviluppi asintotici. Abbiamo che, for $x \rightarrow 0^+$,

$$N(x) := x - \frac{x^3}{3} + o(x^3) - \left(x - \frac{x^3}{6} + o(x^3) \right) + x^{\frac{10}{3}} \log x = -\frac{x^3}{6} + o(x^3) + x^{\frac{10}{3}} \log x.$$

Let us observe that $x^{\frac{10}{3}} \log x = o(x^3)$ for $x \rightarrow 0^+$: infatti

$$\frac{x^{\frac{10}{3}} \log x}{x^3} = x^{\frac{10}{3}-3} \log x \rightarrow 0, \text{ essendo } \frac{10}{3} - 3 > 0,$$

always in virtù of the notable limit sopra richiamato. Therefore $N(x) = \frac{x^3}{6} + o(x^3)$ for $x \rightarrow 0^+$. Quanto al denominator , one can osservare preliminary che

$$(1 - \cos^2 x) = (1 - \cos x)(1 + \cos x) \sim \frac{x^2}{2} \cdot 2 = x^2,$$

so that $D(x) := x^\alpha(1 - \cos^2 x) \sim x^\alpha \cdot x^2 = x^{\alpha+2}$ for $x \rightarrow 0^+$. In conclusione, for $x \rightarrow 0^+$ one has

$$\frac{N(x)}{D(x)} \sim \frac{\frac{x^3}{6}}{x^{\alpha+2}} \rightarrow \begin{cases} 0, & 1 - \alpha > 0, \iff \alpha < 1, \\ -\frac{1}{6}, & 1 - \alpha = 0, \iff \alpha = 1, \\ -\infty, & 1 - \alpha < 0, \iff \alpha > 1. \end{cases}$$

Exercise 5 Study the convergence of the generalized integral

$$\int_2^{+\infty} \frac{1}{x^\alpha \sqrt{x-2}} dx$$

as $\alpha \in \mathbb{R}$ and calcolarlo for $\alpha = 1$.

Soluzione. Sia $f_\alpha(x) := \frac{1}{x^\alpha \sqrt{x-2}}$ the function integranda. Notiamo that it is continuous in $]2, +\infty[$ and hence the integral is generalizzato sia in $x = 2$ that for $x \rightarrow +\infty$. Avendo clearly f_α also sign constant , andiamo a study the comportamento asymptotic at the extremes of the integration interval . Per $x \rightarrow +\infty$ we have che

$$f_\alpha(x) \sim \frac{1}{x^\alpha \sqrt{x}} = \frac{1}{x^{\alpha+1/2}},$$

so that $\int_2^{+\infty} f_\alpha(x) dx < +\infty$ if and only if $\int_2^{+\infty} \frac{1}{x^{\alpha+1/2}} dx < +\infty$ that is, if and only if $\alpha + 1/2 > 1$, ovvero $\alpha > 1/2$. Per $x \rightarrow 2+$ one has che

$$f_\alpha(x) \sim \frac{1}{2^\alpha \sqrt{x-2}},$$

that is integrabile in $x = 2+$. In conclusione, f_α is integrabile in senso generalizzato in $[2, +\infty[$ if and only if $\alpha > 1/2$.

Calcoliamo the integral in the case $\alpha = 1$. Siccome is generalizzato we have che

$$\int_2^{+\infty} \frac{1}{x \sqrt{x-2}} dx = \lim_{a \rightarrow 2+, b \rightarrow +\infty} \int_a^b \frac{1}{x \sqrt{x-2}} dx.$$

Sostituendo $x - 2 = y^2$ ($y > 0$), one has

$$\int \frac{1}{x \sqrt{x-2}} dx = \int \frac{2y}{(y^2 + 2)y} dy = \frac{1}{2} \int \frac{2}{\left(\frac{y}{\sqrt{2}}\right)^2 + 1} dy.$$

Sostituendo ancora $y/\sqrt{2} = t$, one has

$$\int \frac{1}{\left(\frac{y}{\sqrt{2}}\right)^2 + 1} dy = \int \frac{1}{t^2 + 1} dt = \arctan t + c = \arctan \frac{y}{\sqrt{2}} + c.$$

Therefore,

$$\int_2^{+\infty} \frac{1}{x\sqrt{x-2}} dx = \lim_{a \rightarrow 2+, b \rightarrow +\infty} \left(\arctan \sqrt{\frac{b-2}{\sqrt{2}}} - \arctan \sqrt{\frac{a-2}{\sqrt{2}}} \right) = \frac{\pi}{\sqrt{2}}.$$

Appello of the 10.07.2017

THEME 1

Exercise 1 Consider the function

$$f(x) := \log |e^{2x} - 4|.$$

- i) Determine the domain D and study the sign of f ; determine the limits of f at the extremes of D and the asymptotes;
- ii) study the derivability, compute the derivata, study the monotonicity and determine the points of extreme relative and absolute of f ;
- iii) compute f'' and study the concavity and the convexity of f ;
- iv) draw a qualitative graph of f .

Solution. i) the domain is

$$D = \{x : e^{2x} \neq 4\} = \mathbb{R} \setminus \{\log 2\}.$$

SI ha $f(x) \geq 0$ if and only if $|e^{2x} - 4| \geq 1$, that is, if and only if $e^{2x} \geq 5$ oppure $e^{2x} \leq 3$, hence

$$f\left(\frac{\log 5}{2}\right) = f\left(\frac{\log 3}{2}\right) = 0 \quad \text{and } f(x) > 0 \text{ if and only if } x > \frac{\log 5}{2} \text{ oppure } x < \frac{\log 3}{2}.$$

One has moreover

$$\lim_{x \rightarrow -\infty} f(x) = \log 4, \quad \lim_{x \rightarrow \log 2} f(x) = -\infty, \quad \lim_{x \rightarrow +\infty} f(x) = +\infty.$$

As for the oblique asymptote for $x \rightarrow +\infty$, one has

$$\lim_{x \rightarrow +\infty} \frac{f(x)}{x} = \lim_{x \rightarrow +\infty} \frac{\log(e^{2x} - 4)}{x} = 2, \quad \lim_{x \rightarrow +\infty} (f(x) - 2x) = \lim_{x \rightarrow +\infty} \log(e^{2x} - 4) - 2x = \lim_{x \rightarrow +\infty} \log \frac{e^{2x} - 4}{e^{2x}} = 0.$$

Therefore $y = 2x$ is oblique asymptote for $x \rightarrow +\infty$, $y = 2\log 2$ is horizontal asymptote for $x \rightarrow -\infty$ and in $x = \log 2$ one has a vertical asymptote.

- ii) f is differentiable in the whole D , where one has

$$f'(x) = \frac{2e^{2x}}{e^{2x} - 4}.$$

f is therefore strictly decreasing for $x < \log 2$ and strictly increasing for $x > \log 2$. Non risultano hence points of extreme.

iii) Un calcolo diretto gives

$$f''(x) = \frac{-16e^{2x}}{(e^{2x} - 4)^2},$$

so that f is concave in $]-\infty, \log 2[$ and in $\] \log 2, +\infty [$.

iv) the graph is in the picture ??.

Exercise 2 Draw in the Gauss plane the insieme

$$S := \left\{ z \in \mathbb{C} : \operatorname{Re} \frac{z-1}{z-i} \geq 0, |z+1-i| \leq 1 \right\}.$$

Solution. Si tratta in first luogo didetermine the real part of $\frac{z-1}{z+i}$. One has, setting $z = x + iy$,

$$\operatorname{Re} \frac{x-1+iy}{x+i(y-1)} = \operatorname{Re} \frac{(x-1+iy)(x-i(y-1))}{x^2+(y-1)^2} = \frac{x(x-1)+y(y-1)}{x^2+(y-1)^2}.$$

One has therefore

$$\begin{aligned} S &= \{(x, y) \in \mathbb{R}^2 : x^2 - x + y^2 - y \geq 0, (x+1)^2 + (y-1)^2 \leq 1\} \\ &= \left\{ (x, y) \in \mathbb{R}^2 : \left(x - \frac{1}{2}\right)^2 + \left(y - \frac{1}{2}\right)^2 \geq \frac{1}{2}, (x+1)^2 + (y-1)^2 \leq 1 \right\}, \end{aligned}$$

that is, the parte esterna al cerchio centered in $(\frac{1}{2}, \frac{1}{2})$ and raggio $\frac{1}{\sqrt{2}}$ and interna al cerchio centered in $(-1, 1)$ and raggio 1, rappresentata in the picture ??.

Figure 5: Solutions of exercise 2 (Theme 1).

Exercise 3 Compute the integral

$$\int e^{2x} \arctan(3e^x) dx.$$

Solution. Eseguendo the sostituzione $x = \log t$ one has

$$\begin{aligned} \int e^{2x} \arctan(3e^x) dx &= \int t \arctan(3t) dt = \frac{t^2}{2} \arctan 3t - \frac{3}{2} \int \frac{t^2}{1+9t^2} dt \\ &= \frac{t^2}{2} \arctan 3t - \frac{3}{2} \left[\frac{1}{9} \int \frac{1+9t^2}{1+9t^2} dt - \frac{1}{9} \int \frac{1}{1+(3t)^2} dt \right] \\ &= \frac{t^2}{2} \arctan 3t - \frac{t}{6} + \frac{\arctan 3t}{18} + c \\ &= \frac{e^{2x}}{2} \arctan 3e^x - \frac{e^x}{6} + \frac{\arctan 3e^x}{18} + c, \quad c \in \mathbb{R}. \end{aligned}$$

Exercise 4 Compute the limit

$$\lim_{x \rightarrow 0^+} \frac{\arctan \sin x - \sinh x}{x^\alpha (1 - \cos^2 x)}$$

for all values of the parameter $\alpha > 0$.

Solution. Da $\arctan y = y - \frac{y^3}{3} + o(y^3)$ per $y \rightarrow 0$ si deduce, per $x \rightarrow 0$,

$$\arctan \sin x = \sin x - \frac{\sin^3 x}{3} + o(x^3) = x - \frac{x^3}{6} - \frac{x^3}{3} + o(x^3) = x - \frac{x^3}{2} + o(x^3),$$

so that, per $x \rightarrow 0$,

$$\arctan \sin x - \sinh x = x - \frac{x^3}{2} - \left(x + \frac{x^3}{6} \right) + o(x^3) = -\frac{2x^3}{3} + o(x^3).$$

Therefore one has

$$\lim_{x \rightarrow 0^+} \frac{\arctan \sin x - \sinh x}{x^\alpha (1 - \cos^2 x)} = \lim_{x \rightarrow 0^+} \frac{-2x^3/3 + o(x^3)}{x^{\alpha+2} + o(x^{2+\alpha})} = \begin{cases} 0 & \text{for } \alpha < 1 \\ -\frac{2}{3} & \text{for } \alpha = 1 \\ -\infty & \text{for } \alpha > 1. \end{cases}$$

Exercise 5 Study the convergence semplice and assoluta di

$$\sum_{n=2}^{+\infty} \frac{(1-e^a)^n}{n + \sqrt{n}}$$

as $a \in \mathbb{R}$.

Solution. Per la convergenza assoluta si può usare il Test della Radice, che dà

$$\lim_{n \rightarrow \infty} \sqrt[n]{\frac{|1-e^a|^n}{n + \sqrt{n}}} = |1-e^a|.$$

La serie converge assolutamente (e quindi semplicemente) se $|1-e^a| < 1$ e diverge assolutamente se $|1-e^a| > 1$, in quanto il termine generale non è infinitesimale. Per $|1-e^a| = 1$ il Test della Radice non dà informazioni. Risolvendo le disequazioni si deduce che la serie converge assolutamente per $a < \log 2$ e non converge per $a > \log 2$. Per $a = \log 2$ la serie diventa

$$\sum_{n=2}^{+\infty} \frac{(-1)^n}{n + \sqrt{n}}.$$

Per l'asintoticità con la serie armonica $\sum 1/n$ questa serie non converge assolutamente. Inoltre, converge per il criterio di Leibniz, essendo il termine generale a segno alternato e – in valore assoluto – infinitesimale e decrescente.

THEME 1

Exercise 1 Consider the function

$$f(x) := \frac{3x}{\log|2x|}.$$

- i) Determine the domain D and study the simmetries and the sign of f ; determine the limits of f at the extremes of D , the prolongabilità of f and the asymptotes;
- ii) study the derivability, compute the derivative and its main limits, study the monotonicity and determine the points of extreme relative and absolute of f ;
- iii) compute f'' and study the concavity and the convexity of f ;
- iv) draw a qualitative graph of f .

Solution. i) the domain is $D = \{x : x \neq 0, \log|2x| \neq 0\} = \{x : x \neq 0, x \neq \pm\frac{1}{2}\}$. The function is visibilmente dispari, so that the study in $[0, +\infty[$. Per $x > 0$, $f(x) > 0$ if and only if $x > \frac{1}{2}$. One has

$$\lim_{x \rightarrow 0^+} f(x) = 0 \quad (\text{so that } f \text{ is prolongabile con continuity in } x = 0)$$

$$\lim_{x \rightarrow \frac{1}{2}^-} f(x) = -\infty$$

$$\lim_{x \rightarrow \frac{1}{2}^+} f(x) = +\infty$$

$$\lim_{x \rightarrow +\infty} f(x) = +\infty$$

$$\lim_{x \rightarrow +\infty} \frac{f(x)}{x} = 0 \quad (\text{so that non c' è oblique asymptote for } x \rightarrow +\infty).$$

- ii) Per $x > 0, x \neq \frac{1}{2}$ one has

$$f'(x) = \frac{3 \log 2x - 3}{\log^2 2x}.$$

Essendo f prolungabile con continuity in $x = 0$, vediamo if the prolungamento of f is differentiable in 0. A tale scopo calcoliamo

$$\lim_{x \rightarrow 0^+} f'(x) = 0,$$

so that the prolungata of f is differentiable also in $x = 0$, con derivative nulla. The sign of f' dipende solo dal sign of $\log 2x - 1$, that is positive if and only if $x > e/2$. Therefore $e/2$ is a point of strict local minimum.

There are no extremes absolute .

iii) Per $x > 0, x \neq \frac{1}{2}$ one has

$$f''(x) = 3 \frac{\frac{\log^2 2x}{x} - 2(\log 2x - 1)\frac{\log 2x}{x}}{\log^4 2x} = 3 \frac{2 - \log 2x}{x \log^3 2x},$$

that one has > 0 if and only if $\frac{1}{2} < x < \frac{e^2}{2}$, that is, f is convex in the interval $\left[\frac{1}{2}, \frac{e^2}{2}\right]$ and concave negli intervalli $]0, \frac{1}{2}[$ and $[\frac{1}{2}, +\infty[$.

iv) the graph of f is riportato nella figura ??.

Figure 6: the graph of f (Theme 1).

Exercise 2 Given the polynomial

$$z^4 + z^3 + 8i$$

determine first a integer root and then le other roots , writing them in algebraic form.

Solution. Per tentativi, a radice intera is $z = -1$: infatti

$$(-1)^4 + (-1)^3 - 8i + 8i = 0.$$

Eseguendo the divisione dipolinomi, oppure, più semplicemente, raccogliendo z^3 nei primi two addendi and $8i$ negli ultimi two , one has

$$z^4 + z^3 + 8i = (z + 1)(z^3 + 8i),$$

so that le restanti three radici they are le radici cubiche of $-8i = 8e^{\frac{3}{2}\pi i}$, that is, they are

$$2e^{i\frac{\pi}{2}} = 2i, 2e^{(\frac{1}{2}+\frac{2}{3})\pi i} = 2e^{\frac{7}{6}\pi i} = -\sqrt{3} - i, 2e^{(\frac{1}{2}+\frac{4}{3})\pi i} = 2e^{\frac{11}{6}\pi i} = \sqrt{3} - i.$$

Exercise 3 Study the convergence of the series

$$\sum_{n=1}^{\infty} \left(1 + \frac{3x}{n}\right)^{n^2}$$

as $x \in \mathbb{R}$.

Solution. The series is a termini definitively positivi for every $x \in \mathbb{R}$. the Root Test gives

$$\lim_{n \rightarrow +\infty} \sqrt[n]{\left(1 + \frac{3x}{n}\right)^{n^2}} = \lim_{n \rightarrow +\infty} \left(1 + \frac{3x}{n}\right)^n = e^{3x}.$$

The series therefore converges for every $x < 0$ and diverges for every $x > 0$.
 Per $x = 0$ the Root Test nongives informazioni, ma for tale x the series ha
 for general term 1 and hence diverges.

Exercise 4 Compute, for all values of the real parameter α , the limit

$$\lim_{x \rightarrow 0^+} \frac{\cosh(\alpha x) - e^{x^2} + x \log(\cos x)}{x - \sin x + e^{-1/x^2}}.$$

Solution. the numerator, for $x \rightarrow 0$, si sviluppa come

$$\begin{aligned}\cosh \alpha x &= 1 + \frac{1}{2} \alpha^2 x^2 + \frac{1}{24} \alpha^4 x^4 + o(x^4) = 1 + \frac{1}{2} \alpha^2 x^2 + o(x^3) \\ -e^{x^2} &= -1 - x^2 - \frac{1}{2} x^4 + o(x^4) = -1 - x^2 + o(x^3) \\ x \log \cos x &= x \log \left(1 - \frac{1}{2} x^2 + \frac{1}{24} x^4 + o(x^4)\right) = x \left(\frac{-x^2}{2} + o(x^2)\right) = \frac{-x^3}{2} + o(x^3),\end{aligned}$$

so that

$$\cosh(\alpha x) - e^{x^2} + x \log(\cos x) = x^2 \left(\frac{\alpha^2}{2} - 1\right) - \frac{x^3}{2} + o(x^3) = \begin{cases} x^2 \left(\frac{\alpha^2}{2} - 1\right) + o(x^2) & \text{if } \alpha \neq \pm\sqrt{2} \\ -\frac{x^3}{2} + o(x^3) & \text{if } \alpha = \pm\sqrt{2}. \end{cases}$$

the denominator , for $x \rightarrow 0$, si sviluppa come

$$x - \sin x + e^{-1/x^2} = \frac{x^3}{6} + o(x^3),$$

in quanto $e^{-1/x^2} = o(x^\beta)$ for every β reale. The limit hence vale

$$\begin{aligned}\lim_{x \rightarrow 0^+} \frac{\cosh(\alpha x) - e^{x^2} + x \log(\cos x)}{x - \sin x + e^{-1/x^2}} &= \lim_{x \rightarrow 0^+} \frac{x^2 \left(\frac{\alpha^2}{2} - 1\right) - \frac{x^3}{2} + o(x^3)}{\frac{x^3}{6} + o(x^3)} \\ &= \lim_{x \rightarrow 0^+} \begin{cases} \frac{x^2 \left(\frac{\alpha^2}{2} - 1\right) + o(x^2)}{\frac{x^3}{6} + o(x^3)} & \text{if } \alpha \neq \pm\sqrt{2} \\ \frac{-\frac{x^3}{2} + o(x^3)}{\frac{x^3}{6} + o(x^3)} & \text{if } \alpha = \pm\sqrt{2}. \end{cases} \\ &= \begin{cases} +\infty & \text{if } |\alpha| < \sqrt{2} \\ -\infty & \text{if } |\alpha| > \sqrt{2} \\ -3 & \text{if } \alpha = \pm\sqrt{2}. \end{cases}\end{aligned}$$

Exercise 5 Study the convergence of the generalized integral

$$\int_0^{+\infty} xe^{ax} (2 + \cos x) dx$$

as $a \in \mathbb{R}$. Compute

$$\int_0^{+\infty} xe^{-x} \cos x dx$$

(sugg.: compute preliminarily a primitive of $e^{-x} \cos x$).

Solution. Una primitive of the integrand può essere calcolata for every a , so that the discussione of the convergence può essere fatta sia direttamente dalla definizione, sia mediante criteri of convergence . Usando the comparison criterion one has, for $a \geq 0$,

$$xe^{ax}(2 + \cos x) \geq x \text{ for every } x \geq 0$$

and hence the integral diverges. Per $a < 0$ the asymptotic comparison dà, ad esempio,

$$xe^{ax}(2 + \cos x) = o(e^{ax/2}),$$

perché

$$\lim_{x \rightarrow +\infty} \frac{xe^{ax}(2 + \cos x)}{e^{ax/2}} \leq \lim_{x \rightarrow +\infty} \frac{3xe^{ax}}{e^{ax/2}} = \lim_{x \rightarrow +\infty} 3xe^{\frac{ax}{2}} = 0.$$

Siccome $\int_0^{+\infty} e^{ax/2} dx < +\infty$, the integral converges.

Per the primitiva, calcoliamo preliminarly

$$\begin{aligned} \int e^{-x} \cos x dx &= -e^{-x} \cos x - \int e^{-x} \sin x dx \\ &= -e^{-x} \cos x + e^{-x} \sin x - \int e^{-x} \cos x dx, \end{aligned}$$

so that

$$\int e^{-x} \cos x dx = \frac{e^{-x}}{2}(\sin x - \cos x) + c.$$

Ora integriamo by parts prendendo x come factor finito and $e^{-x} \cos x$ come differential factor. Risulta

$$\int xe^{-x} \cos x dx = x \frac{e^{-x}}{2}(\sin x - \cos x) - \int \frac{e^{-x}}{2}(\sin x - \cos x) dx.$$

Calcoliamo separatamente

$$\begin{aligned} \int e^{-x} \sin x dx &= -e^{-x} \sin x + \int e^{-x} \cos x dx \\ &= -e^{-x} \sin x - e^{-x} \cos x - \int e^{-x} \sin x dx, \end{aligned}$$

so that

$$\int e^{-x} \sin x dx = -\frac{e^{-x}}{2}(\sin x + \cos x) + c.$$

In definitiva,

$$\begin{aligned} \int_0^{+\infty} xe^{-x} \cos x dx &= \lim_{b \rightarrow +\infty} \left[-x \frac{e^{-x}}{2} (\sin x - \cos x) \Big|_0^b + \frac{1}{4} e^{-x} (\sin x + \cos x) \Big|_0^b \right. \\ &\quad \left. + \frac{1}{4} e^{-x} (\sin x - \cos x) \Big|_0^b \right] \\ &= 0. \end{aligned}$$

(NB. Non è strano che il risultato sia nullo: il integrando non ha segno costante.)

Appello of the 29.01.2018

THEME 1

Exercise 1 Consider the function

$$f(x) := \log \frac{|x^2 - 5|}{x+1}.$$

- i) Determine the domain D of f , its simmetries and study the sign; determine the limits of f at the extremes of D and the asymptotes;
- ii) study the derivability, compute the derivative and study the monotonicity of f and determine the points of extreme relative and absolute;
- iii) draw a qualitative graph of f .

Solution. i) Da $\frac{|x^2 - 5|}{x+1} > 0$ segue che $D = \{x > -1, x \neq \sqrt{5}\}$. There are no simmetries evidenti. $f(x) \leq 0$ if and only if

$$x > -1$$

e

$$|x^2 - 5| \leq x + 1 \Leftrightarrow -x - 1 \leq x^2 - 5 \leq x + 1 \Leftrightarrow \begin{cases} x^2 + x - 4 \geq 0 \\ x^2 - x - 6 \leq 0, \end{cases}$$

that is, if and only if $\frac{-1 + \sqrt{17}}{2} \leq x \leq 3$.

One has moreover

$$\begin{aligned} \lim_{x \rightarrow -1^+} f(x) &= +\infty \\ \lim_{x \rightarrow \sqrt{5}} f(x) &= -\infty \\ \lim_{x \rightarrow +\infty} f(x) &= +\infty. \end{aligned}$$

Siccome

$$\lim_{x \rightarrow +\infty} \frac{f(x)}{x} = 0,$$

non ci they are asymptotes obliqui, monly one two asymptotes verticali (oltre ad a asintoto orizzontale “così alto that non si vede” (cit.))

ii) Le regole diderivazione posthey are essere applicate in the whole D , perché the point in which the argument of the modulo si annulla non appartiene al domain. Siccome $f(x) = \log|x^2 - 5| - \log(x + 1)$ and ricordando che $\frac{d}{dx} \log|g(x)| = \frac{g'(x)}{g(x)}$ dove $g(x) \neq 0$, one has for every $x \in D$

$$f'(x) = \frac{2x}{x^2 - 5} - \frac{1}{x + 1} = \frac{x^2 + 2x + 5}{(x^2 - 5)(x + 1)}.$$

Siccome the polynomial al numerator is always positive, $f'(x) > 0$, and hence f is increasing, if and only if $x > \sqrt{5}$. There are no points of extreme.

iii) the graph of f is in the picture ??.



Figure 7: the graph of f (Theme 1).

Exercise 2 Consider the sequence

$$a_n = \frac{(-1)^n e^{2n} \sin \frac{1}{n}}{(n-1)!}, \quad n \in \mathbb{N}, n \geq 2.$$

- a) Compute $\lim_{n \rightarrow \infty} a_n$;
- b) study the absolute convergence and the convergence semplice of the series

$\sum_{n=2}^{\infty} a_n$.

Solution. a) Siccome $a_n \sim \frac{(e^2)^n}{n!}$ for $n \rightarrow \infty$, $\lim_{n \rightarrow \infty} a_n = 0$ (ricordando a limit fondamentale).

b) the criterion of the asymptotic comparison and the criterio of the rapporto danno

$$\lim_{n \rightarrow \infty} \frac{e^{2(n+1)}}{(n+1)!} \frac{n!}{e^{2n}} = \lim_{n \rightarrow \infty} \frac{e^2}{n+1} = 0,$$

so that the series absolutely converges and hence converges.

the fatto che $a_n \rightarrow 0$ si poteva also dedurre direttamente dalla convergence of the series .

NOTA: applicando the criterio di Leibniz one may dedurre direttamente the convergence of the series . Risulta che $|a_n|$ is decreasing if and only if $e^2 \leq n$, the that is vero for every $n > 2$ (the dimostrazione richiede a po' dilavoro). Resta comunque da verificare the absolute convergence . Siccome in questo case is vera, the uso of the criterio di Leibniz is of the tutto inutile.

Exercise 3 Sia $f(z) = z^2 + \bar{z}|z|$. Solve the equation

$$zf(z) = |z|^3 - 8i,$$

writing the solutions in algebraic form and disegnandole in the Gauss plane

Solution. The equation is

$$z^3 + z\bar{z}|z| = |z|^3 - 8i.$$

Siccome $z\bar{z}|z| = |z|^2|z| = |z|^3$, the equation diventa

$$z^3 = -8i.$$

Le three radici cubiche of $-8i = 8e^{i3\pi/2}$ they are date da

$$2e^{i\frac{\pi}{2}} = 2i, \quad 2e^{i\frac{7\pi}{6}} = -\sqrt{3} - i, \quad 2e^{i\frac{11\pi}{6}} = \sqrt{3} - i,$$

rappresentate in the picture ??.


Exercise 4 Compute the limit

$$\lim_{x \rightarrow +\infty} \frac{\log(x+3) - \log(x+1) - \sin \frac{2}{x}}{\cos \sin \frac{1}{2x} - e^{\frac{\alpha}{x^2}} - e^{-x}}$$

as $\alpha \in \mathbb{R}$.

Solution. the numerator:

$$\begin{aligned} \log(x+3) - \log(x+1) - \sin \frac{2}{x} &= \log x + \log \left(1 + \frac{3}{x}\right) - \log x - \log \left(1 + \frac{1}{x}\right) - \sin \frac{2}{x} \\ &= \log \left(1 + \frac{3}{x}\right) - \log \left(1 + \frac{1}{x}\right) - \sin \frac{2}{x} \\ &= \frac{3}{x} - \frac{9}{2x^2} - \frac{1}{x} + \frac{1}{2x^2} - \frac{2}{x} + o\left(\frac{2}{x^2}\right) \\ &= -\frac{4}{x^2} + o\left(\frac{1}{x^2}\right) \end{aligned}$$



grafici/2018-I-T1c-eps-converted-to.pdf

Figure 8: Solutions of exercise 3 (Theme 1).

for $x \rightarrow +\infty$. the denominator (ricordando che $e^{-x} = o(1/x^\alpha)$ for $x \rightarrow +\infty$ for every α):

$$\begin{aligned} \cos \sin \frac{1}{x} - e^{\frac{\alpha}{x^2}} - e^{-x} &= 1 - \frac{1}{2} \sin^2 \frac{1}{2x} + \frac{1}{24} \sin^4 \frac{1}{2x} - \left(1 + \frac{\alpha}{x^2} + \frac{\alpha^2}{2x^4}\right) + o\left(\frac{1}{x^4}\right) \\ &= -\frac{1}{2} \left(\frac{1}{2x} - \frac{1}{6(2x)^3}\right)^2 + \frac{1}{24(2x)^4} - \frac{\alpha}{x^2} - \frac{\alpha^2}{2x^4} + o\left(\frac{1}{x^4}\right) \\ &= \left(-\frac{1}{8} - \alpha\right) \frac{1}{x^2} + \left(\frac{1}{96} + \frac{1}{24 \cdot 2^4} - \frac{\alpha^2}{2}\right) \frac{1}{x^4} + o\left(\frac{1}{x^4}\right) \\ &= \begin{cases} -\left(\frac{1}{8} + \alpha\right) \frac{1}{x^2} + o\left(\frac{1}{x^2}\right) & \text{if } \alpha \neq -\frac{1}{2} \\ \frac{1}{192} \frac{1}{x^4} + o\left(\frac{1}{x^4}\right) & \text{if } \alpha = -\frac{1}{2} \end{cases} \end{aligned}$$

for $x \rightarrow +\infty$. Di conseguenza,

$$\lim_{x \rightarrow +\infty} \frac{\log(x+3) - \log(x+1) - \sin \frac{2}{x}}{\cosh \sin \frac{1}{2x} - e^{\frac{\alpha}{x^2}} - e^{-x}} = \lim_{x \rightarrow +\infty} \begin{cases} \frac{\frac{-4}{x^2} + o\left(\frac{1}{x^2}\right)}{-\left(\frac{1}{8} + \alpha\right) \frac{1}{x^2} + o\left(\frac{1}{x^2}\right)} = \frac{32}{1+8\alpha} & \text{if } \alpha \neq -\frac{1}{8} \\ \frac{\frac{-4}{x^2} + o\left(\frac{1}{x^2}\right)}{\frac{1}{192} \frac{1}{x^4} + o\left(\frac{1}{x^4}\right)} = -\infty & \text{if } \alpha = \frac{1}{8}. \end{cases}$$

NOTA: the numerator poteva also essere scritto come

$$\begin{aligned}
\log(x+3) - \log(x+1) - \sin \frac{2}{x} &= \log \frac{x+3}{x+1} - \sin \frac{2}{x} = \log \left(1 + \frac{2}{x+1}\right) - \sin \frac{2}{x} \\
&= \frac{2}{x+1} - \frac{1}{2} \left(\frac{2}{x+1}\right)^2 - \frac{2}{x} + o\left(\frac{2}{x^2}\right) \\
&= -2 \frac{2x+1}{x(x+1)^2} + o\left(\frac{2}{x^2}\right) \\
&= -2 \frac{1+2x}{x(x+1)^2} + o\left(\frac{2}{x^2}\right) \sim -\frac{4}{x^2}
\end{aligned}$$

for $x \rightarrow +\infty$. The maggior parte degli studenti that ha svolto the calcolo in questo modo ha tralasciato the termine di order 2 nello sviluppo of the logarithm.

Exercise 5 a) Study the convergence of the generalized integral

$$\int_{\sqrt{2}}^{+\infty} \frac{1}{x^\alpha \sqrt{x^2 - 2}} dx$$

as $\alpha \in \mathbb{R}$;

b) calcolarlo for $\alpha = 1$.

Solution. a) The integranda $f(x)$ is continuous in $\sqrt{2}, +\infty$, so that si deve controllare the convergence sia for $x \rightarrow \sqrt{2}^+$ that for $x \rightarrow +\infty$. Per $x \rightarrow \sqrt{2}^+$,

$$f(x) \sim \frac{1}{\sqrt{x - \sqrt{2}}},$$

so that the integral converges for every α . Per $x \rightarrow +\infty$,

$$f(x) \sim \frac{1}{x^{\alpha+1}},$$

so that the integral converges if and only if $\alpha > 0$.

b) Con the sostituzione $x = \sqrt{2} \cosh t$, one has (for $t > 0$)

$$\int_{\sqrt{2}}^{+\infty} \frac{1}{x \sqrt{x^2 - 2}} dx = \int_0^{+\infty} \frac{\sqrt{2} \sinh t}{2 \cosh t \sinh t} dt = \sqrt{2} \int_0^{+\infty} \frac{e^t}{1 + e^{2t}} dt = \sqrt{2} \arctan e^t \Big|_0^{+\infty} = \sqrt{2} \left(\frac{\pi}{2} - \frac{\pi}{4} \right) = \frac{\sqrt{2}\pi}{4}.$$

In alternativa, con the sostituzione $y = \sqrt{x^2 - 2}$, seguita dalla sostituzione $z = y/\sqrt{2}$, one gets ,

$$\int_2^{+\infty} \frac{1}{x \sqrt{x^2 - 4}} dx = \int_0^{+\infty} \frac{1}{y^2 + 4} dy = \frac{1}{\sqrt{2}} \int_0^{+\infty} \frac{1}{z^2 + 1} dz = \frac{1}{\sqrt{2}} \arctan z \Big|_0^{+\infty} = \frac{1}{\sqrt{2}} \frac{\pi}{2}.$$

Un terzo modo di compute the integral is the seguente:

$$\int_{\sqrt{2}}^{+\infty} \frac{1}{x \sqrt{x^2 - 2}} dx = \int_{\sqrt{2}}^{+\infty} \frac{dx}{x^2 \sqrt{1 - 2/x^2}} = \frac{\sqrt{2}}{2} \int_1^{+\infty} \frac{dt}{t^2 \sqrt{1 - 1/t^2}} = \frac{1}{\sqrt{2}} \arcsin \frac{1}{t} \Big|_1^{+\infty} = \frac{1}{\sqrt{2}} \frac{\pi}{2}.$$

Exercise . Sia $x_0 \in \mathbb{R}$ and si definisca the sequence $\{a_n : n \in \mathbb{N}\}$ ponendo

$$a_0 = x_0 \text{ e, for every } n \geq 1, a_{n+1} = \sin a_n.$$

- a) prove that a_n is definitively monotonic for $n \rightarrow +\infty$;
- b) prove that $\lim_{n \rightarrow +\infty} a_n = 0$.

Solution. a) Per $n \geq 1$ one has $|a_n| = |\sin(a_{n-1})| \leq 1$. If $a_1 \in [0, 1]$, allora da $\sin x \leq x \forall x \geq 0$ si ricava $a_{n+1} = \sin a_n \leq a_n$ and hence the sequence is definitively decreasing. If instead $a_1 \in [-1, 0]$ one gets the sequence is definitively increasing.

b) In ogni case the sequence ha a limit $\ell \in [-1, 1]$. If for assurdo fosse $\ell \neq 0$ si avrebbe, essendo the function seno continuous,

$$\lim_{n \rightarrow +\infty} \frac{|a_{n+1}|}{|a_n|} = \frac{|\sin \ell|}{|\ell|} < 1,$$

the that implicherebbe the convergence of the series $\sum_{n=0}^{\infty} |a_n|$, the that a sua volta implicherebbe che a_n converges a 0, cosicché $0 = \ell \neq 0$. Hence $\ell = 0$. In alternativa, always for the continuity di sin,

$$\ell = \lim a_{n+1} = \lim \sin a_n = \sin \ell$$

that ha $\ell = 0$ come unica soluzione.

Appello of the 16.02.2018

THEME 1

Exercise 1 Consider the function

$$f(x) = \begin{cases} e^{x - \frac{1}{|x-2|}} & \text{for } x \neq 2 \\ 0 & \text{for } x = 2. \end{cases}$$

- i) Determine the domain D of f , its simmetries and study the sign; In order to determine i limits of f at the extremes of D and the asymptotes;
- ii) si dica if f is continuous in the whole \mathbb{R} .
- iii) compute the derivative and study the monotonicity of f and determine the points of extreme relative and absolute ; compute the main limits of f' ; in particolare si dica if f is differentiable in the whole \mathbb{R} ; the study of the second derivative may be skipped
- iv) draw a qualitative graph of f .

Solution. i)DOMINIO: $|x - 2| \neq 0 \iff x \neq 2$, hence $D = \mathbb{R} \setminus \{2\} \cup \{2\} = \mathbb{R}$
LIMITI:

$$\lim_{x \rightarrow 2} f(x) = e^2 \cdot e^{-\infty} = e^2 \cdot 0 = 0 \quad \lim_{x \rightarrow +\infty} f(x) = e^{+\infty} = +\infty \quad \lim_{x \rightarrow -\infty} f(x) = e^{-\infty} = 0$$

ASINTOTI

$$\lim_{x \rightarrow +\infty} f(x)/x = \lim_{x \rightarrow +\infty} \frac{e^x}{x} = +\infty$$

hence non ci they are asymptotes obliqui.

ii) CONTINUITY: The function is continuous in $\mathbb{R} \setminus \{2\}$ because superposition discontinuous. It is continuous also for $x = 2$ since $\lim_{x \rightarrow 2} = 0 = f(2)$. Hence f is continuous.

iii) if $x > 2$ one has

$$f'(x) = \left(e^{x-\frac{1}{x-2}}\right) \left(1 + \frac{1}{(x-2)^2}\right);$$

if $x < 2$ one has

$$f'(x) = \left(e^{x+\frac{1}{x-2}}\right) \left(1 - \frac{1}{(x-2)^2}\right).$$

Hence $f'(x) \geq 0$ if

$$\begin{cases} x > 2 \\ 1 + \frac{1}{(x-2)^2} \geq 0 \end{cases} \cup \begin{cases} x < 2 \\ 1 - \frac{1}{(x-2)^2} \geq 0 \end{cases}$$

that is, if

$$\begin{aligned} x \in]2, +\infty[& \left(\bigcup \left(]-\infty, 2[\cap \{x : (x-2)^2 \geq 1\} \right) \right) \\ & =]2, +\infty[\left(\bigcup \left(]-\infty, 2[\cap \{x : (x-2) \leq -1 \text{ oppure } (x-2) \geq 1\} \right) \right) \end{aligned}$$

that is, if

$$x \in]2, +\infty[\cup]-\infty, 1].$$

Furthermore,, since

$$\lim_{x \rightarrow 2^-} f'(x) = \lim_{x \rightarrow 2^-} \left(e^{x+\frac{1}{x-2}}\right) \left(1 - \frac{1}{(x-2)^2}\right) = -e^2 \lim_{x \rightarrow 2^-} \frac{e^{\frac{1}{x-2}}}{(x-2)^2} = 0,$$

$$\lim_{x \rightarrow 2^+} f'(x) = \lim_{x \rightarrow 2^+} \left(e^{x-\frac{1}{x-2}}\right) \left(1 + \frac{1}{(x-2)^2}\right) = -e^2 \lim_{x \rightarrow 2^+} \frac{e^{-\frac{1}{x-2}}}{(x-2)^2} = 0$$

one has che f is differentiable in $x = 2$ and $f'(2) = 0$. Concludendo, f is differentiable on tutto the domain $D = \mathbb{R}$, anzi is di classe C^1 .

Dalthe study of the monotonicity f ha a relative maximum in $x = 1$ and a absolute minimum in $x = 0$.

iv) the graph is in the picture ??.



grafici/2018-II-T1f-eps-converted-to.pdf

Figure 9: the graph of f (Theme 1).

Exercise 2 Study as $x \in \mathbb{R}$ the convergence semplice and assoluta of the series

$$\sum_{n=1}^{\infty} \frac{(2x-1)^n}{(2n+3)^2}.$$

Solution. Let us study the absolute convergence con the criterio of the rapporto

$$\lim_{n \rightarrow \infty} \frac{|2x-1|^{n+1}}{(2n+5)^2} \frac{(2n+3)^2}{|2x-1|^n} = |2x-1| \lim_{n \rightarrow \infty} \frac{(2n+3)^2}{(2n+5)^2} = |2x-1|$$

o, alternativamente, con Root Test

$$\lim_{n \rightarrow \infty} \sqrt[n]{\frac{|2x-1|^n}{(2n+3)^2}} = \lim_{n \rightarrow \infty} |2x-1| \sqrt[n]{\frac{1}{(2n+3)^2}} = |2x-1|.$$

Therefore the series absolutely converges – and hence converges– for $0 < x < 1$ and diverges assolutamente and does not converge (perché the general term is not infinitesimal) for $x < 0$ and for $x > 1$. Per $x = 0$ and $x = 1$ the Root Test and of the rapporto non danno informazioni. Per $x = 0, x = 1$

the series diventa

$$\sum_{n=1}^{\infty} \frac{(-1)^n}{(2x-1)^2}, \quad \sum_{n=1}^{\infty} \frac{1}{(2x-1)^2},$$

rispettivamente, and hence absolutely converges, and hence semplicemente, for asymptotic comparison con the series converging

$$\sum_{n=1}^{\infty} \frac{1}{n^2}.$$

Per $0 \leq x < 1/2$ the convergence semplice one may also dedurre dal criterio di Leibniz.

Exercise 3 Solve the equation

$$z^2\bar{z} + z\bar{z}^2 = 4 \operatorname{Im}(iz)$$

and diventasegnarne the solutions on Gauss plane .

Solution. Poniamo $z = \rho(\cos 0a + i \sin 0a)$. The equation diventa

$$2\rho^3 \cos 0a = 4\rho \cos 0a.$$

Hence, $\rho = 0$, that is, $z = 0$, oppure

$$\rho^2 \cos 0a = 2 \cos 0a,$$

vale a diventare $\rho^2 = 2$, o $z = \pm\rho i$, $\rho > 0$. Concludendo, l'insieme of the solutions on Gauss plane is the unione of the retta verticale for the origine and the circolo diraggio $\sqrt{2}$, rappresentati in the picture ??.




Figure 10: Solutions of exercise 3 (Theme 1).

Exercise 4

a) Compute the limit

$$\lim_{x \rightarrow 0} \frac{(4 \cos x - \alpha)^2 - 4x^4}{x^4 \sin^2 x}$$

as $\alpha \in \mathbb{R}$.

Solution. the denominator is asymptotic to x^6 for $x \rightarrow 0$. the numerator: one has, for $x \rightarrow 0$,

$$\begin{aligned} (4 \cos x - \alpha)^2 - 4x^4 &= (4 - \alpha - 2x^2 + \frac{x^4}{6} + o(x^4))^2 - 4x^4 \\ &= \begin{cases} 4 - \alpha + o(1) & \text{for } \alpha \neq 4 \\ 4x^4 - \frac{2x^6}{3} - 4x^4 + o(x^6) & \text{for } \alpha = 4 \end{cases} \\ &= \begin{cases} 4 - \alpha + o(1) & \text{for } \alpha \neq 4 \\ -\frac{2x^6}{3} + o(x^6) & \text{for } \alpha = 4. \end{cases} \end{aligned}$$

One has therefore

$$\lim_{x \rightarrow 0} \frac{4(\cos x - \alpha)^2 - x^4}{x^4 \sin^2 x} = \begin{cases} +\infty & \text{for } \alpha \neq 4 \\ -\frac{2}{3} & \text{for } \alpha = 4. \end{cases}$$

Exercise 5 a) Study the convergence of the generalized integral

$$\int_0^{\frac{\pi^2}{3}} x^\alpha \sin(\sqrt{3x}) dx$$

as $\alpha \in \mathbb{R}$;

b) calcolarlo for $\alpha = \frac{1}{2}$.

Solution. a) The integranda $g(x)$ is continuous in the integration interval, possibly except at the first extreme. Per $x \rightarrow 0^+$ one has

$$g(x) \sim \sqrt{3} x^{\alpha+\frac{1}{2}}.$$

The integral is converging if and only if the exponent is greater than -1 , that is, if and only if $\alpha > -\frac{3}{2}$.

b) One has, con the sostituzione $3x = t^2$, which gives $dx = \frac{2}{3}t dt$,

$$\begin{aligned} \int_0^{\frac{\pi^2}{3}} x^{\frac{1}{2}} \sin(\sqrt{3x}) dx &= \frac{2}{3\sqrt{3}} \int_0^{\pi} t^2 \sin t dt \\ &\quad (\text{by parts}) = \frac{2}{3\sqrt{3}} \left(-t^2 \cos t \Big|_0^{\pi} + 2 \int_0^{\pi} t \cos t dt \right) \\ &\quad (\text{by parts}) = \frac{2}{3\sqrt{3}} \pi^2 + \frac{2}{3\sqrt{3}} \left(2t \sin t \Big|_0^{\pi} - 2 \int_0^{\pi} \sin t dt \right) \\ &= \frac{2}{3\sqrt{3}} (\pi^2 - 4). \end{aligned}$$

Appello of the 9.07.2018

THEME 1

Exercise 1 Consider the function

$$f(x) = \log |2 - 3e^{3x}|.$$

- i) Si determini the domain D and study the sign of f ;
- ii) si determinino the limits of f at the extremes of D and the asymptotes;
- iii) find the derivative and study the monotonicity of f , determinandone the points of extreme relative and absolute ; the study of the second derivative may be skipped;
- iv) si diventasegni a qualitative graph of f .

Solution. i) the domain of f is dato dalla condizione $3e^{3x} \neq 2$, that is,

$$D = \{x \in \mathbb{R} : x \neq \frac{\log \frac{2}{3}}{3}\}.$$

the sign of f is positive if and only if $|2 - 3e^{3x}| > 1$. Elevando al quadrato one gets the inequality equivalente

$$9e^{6x} - 12e^{3x} + 3 > 0.$$

setting $e^{3x} = y$ and diventavidendo for 3, one gets the inequality $3y^2 - 4y + 1 > 0$, that ha for solutions $y < 1/3$, $y > 1$. Therefore $f(x) \geq 0$ if and only if

$$x \leq \frac{-\log 3}{3} \text{ oppure } x \geq 0.$$

In alternativa: if $2 - 3e^{3x} \geq 0$, one has:

$$|2 - 3e^{3x}| > 1 \iff 2 - 3e^{3x} > 1 \iff e^{3x} < \frac{1}{3} \iff x < \frac{1}{3} \log\left(\frac{1}{3}\right) = -\frac{1}{3} \log 3.$$

If instead $2 - 3e^{3x} < 0$:

$$|2 - 3e^{3x}| > 1 \iff 3e^{3x} - 2 > 1 \iff e^{3x} > 1 \iff x > \frac{1}{3} \log(1) = 0.$$

Therefore $f(x) \geq 0$ if and only if

$$x \leq \frac{-\log 3}{3} \text{ oppure } x \geq 0.$$

ii) One has

$$\lim_{x \rightarrow -\infty} f(x) = \lim_{x \rightarrow -\infty} \log(2 - 3e^{3x}) = \log 2,$$

perché $\lim_{x \rightarrow -\infty} e^{3x} = 0$, hence the retta $y = \log 2$ is a horizontal asymptote.
Furthermore,

$$\lim_{x \rightarrow +\infty} f(x) = \lim_{x \rightarrow +\infty} \log(3e^{3x} - 2) = +\infty,$$

e

$$\lim_{x \rightarrow +\infty} \frac{f(x)}{x} = \lim_{x \rightarrow +\infty} \frac{\log(3e^{3x} - 2)}{x} = \lim_{x \rightarrow +\infty} \frac{f(x)}{x} = \lim_{x \rightarrow +\infty} \frac{3x + \log(3 - 2e^{-3x})}{x} = 3,$$

$$\lim_{x \rightarrow +\infty} f(x) - 3x = \lim_{x \rightarrow +\infty} \log(3 - 2e^{-3x}) = \log 3.$$

Therefore the retta $y = 3x + \log 3$ is oblique asymptote for $x \rightarrow +\infty$.

Finally ,

$$\lim_{x \rightarrow \frac{\log \frac{2}{3}}{-3}} f(x) = \lim_{y \rightarrow 0^+} \log y = -\infty,$$

$x = \frac{\log \frac{2}{3}}{-3}$ is a vertical asymptote.

(iii) Le regole dederivazione posthey are essere applicate in the whole D , perché the point in which the argument of the modulo si annulla non appartiene al domain. Ricordando che $\frac{d}{dx} \log |g(x)| = \frac{g'(x)}{g(x)}$ dove $g(x) \neq 0$, one has for every $x \in D$

$$f'(x) = \frac{9e^{3x}}{3e^{3x} - 2}.$$

Siccome the numerator is always positive, $f'(x) > 0$, and hence f is increasing, if and only if $x > \frac{\log \frac{2}{3}}{-3}$. There are no points of extreme.

(iv) the graph is in the picture ??.

Figure 11: the graph of f (Theme 1).

Exercise 2 Solve the inequality

$$|z|^2 \operatorname{Re}\left(\frac{1}{z}\right) \leq \operatorname{Im}(\bar{z}^2)$$

rappresentandone the solutions on Gauss plane .

Solution. Notiamo prima that bisogna avere $z \neq 0$. Poniamo $z = x + iy$. Siccome, for $z \neq 0$,

$$\operatorname{Re}\left(\frac{1}{z}\right) = \operatorname{Re}\left(\frac{\bar{z}}{z\bar{z}}\right) = \operatorname{Re}\left(\frac{x - iy}{|z|^2}\right) = \frac{x}{|z|^2},$$

the inequality, for $z \neq 0$, is equivalente a

$$x \leq \operatorname{Im}((x - iy)^2) = \operatorname{Im}(x^2 - y^2 - 2ixy) = -2xy,$$

that a sua volta is equivalente a

$$x(1+2y) \leq 0, \quad x^2 + y^2 \neq 0,$$

that ha for solutions the insieme

$$\left(\{(x, y) \in \mathbb{R}^2 : x \geq 0, y \leq -\frac{1}{2}\} \cup \{(x, y) \in \mathbb{R}^2 : x \leq 0, y \geq -\frac{1}{2}\} \right) \setminus \{(0, 0)\}.$$

Solutions they are in the picture ???. **NB:** $z = 0$ is da togliere!

Figure 12: Solutions of exercise 2 (Theme 1).

Exercise 3 Compute the limit

$$\lim_{x \rightarrow +\infty} \frac{(\log(1+x) - \log x - \frac{\alpha}{x})^2}{(1 - \cos \frac{1}{x})^2 + e^{-x}}$$

as $\alpha \in \mathbb{R}$.

Solution. Per $x \rightarrow +\infty$ one has

$$\log(1+x) - \log x - \frac{\alpha}{x} = \log x + \log \left(1 + \frac{1}{x}\right) - \log x - \frac{\alpha}{x} = \frac{1}{x} - \frac{1}{2x^2} - \frac{\alpha}{x} + o\left(\frac{1}{x^2}\right) = \begin{cases} \frac{1-\alpha}{x} + o\left(\frac{1}{x}\right) & \text{for } \alpha \neq 1 \\ -\frac{1}{2x^2} + o\left(\frac{1}{x^2}\right) & \text{for } \alpha = 1. \end{cases}$$

One has therefore, for $x \rightarrow +\infty$,

$$\left(\log(1+x) - \log x - \frac{\alpha}{x}\right)^2 = \begin{cases} \frac{(1-\alpha)^2}{x^2} + o\left(\frac{1}{x^2}\right) & \text{for } \alpha \neq 1 \\ \frac{1}{4x^4} + o\left(\frac{1}{x^4}\right) & \text{for } \alpha = 1. \end{cases}$$

Per the denominator one has

$$\left(1 - \cos \frac{1}{x}\right)^2 + e^{-x} = \left(\frac{1}{2x^2} + o\left(\frac{1}{x^2}\right)\right)^2 + e^{-x} = \frac{1}{4x^4} + o\left(\frac{1}{x^4}\right),$$

poiché $e^{-x} = o\left(\frac{1}{x^n}\right)$ for $x \rightarrow +\infty$ qualunque sia $n > 0$. Therefore one has

$$\lim_{x \rightarrow +\infty} \frac{(\log(1+x) - \log x - \frac{\alpha}{x})^2}{(1 - \cos \frac{1}{x})^2 + e^{-x}} = \begin{cases} +\infty & \text{for } \alpha \neq 1 \\ 1 & \text{for } \alpha = 1. \end{cases}$$

Exercise 4 Study as $\alpha \in \mathbb{R}$ the convergence of the series

$$\sum_{n=1}^{\infty} n \arctan \left(\frac{2^{\alpha n}}{n} \right).$$

Solution. The series is a termini positivi. Osserviamo innanzitutto that for $\alpha > 0$ the general term is not infinitesimal, in quanto $\lim_{n \rightarrow \infty} 2^{\alpha n}/n = +\infty$, so that $\lim_{n \rightarrow \infty} \arctan\left(\frac{2^{\alpha n}}{n}\right) = \pi/2$, and hence

$$\lim_{n \rightarrow \infty} n \arctan\left(\frac{2^{\alpha n}}{n}\right) = +\infty.$$

Therefore for $\alpha > 0$ the series diverges. Per $\alpha \leq 0$ one can usare the criterion of the asymptotic comparison, that dice that the series ha lo stesso character of the series

$$\sum_{n=1}^{\infty} n \frac{2^{\alpha n}}{n} = \sum_{n=1}^{\infty} 2^{\alpha n}.$$

Quest'ultima is the series geometrica diragione 2^α , that converges if and only if $2^\alpha < 1$, hence if and only if $\alpha < 0$.

Exercise 5 a) Compute a primitive di

$$f(x) = \frac{x^2}{(x^2+1)(x^2+2)}$$

(sugg.: cercare a decomposizione of the integrand of the tipo $\frac{A}{x^2+1} + \frac{B}{x^2+2}$).

b) Study the convergence of the generalized integral

$$\int_0^{+\infty} \log \frac{x^\alpha + 2}{x^\alpha + 1} dx.$$

as $\alpha > 0$.

c) Compute the integral for $\alpha = 2$.

Solution. a) One has

$$\frac{x^2}{(x^2+1)(x^2+2)} = \frac{A}{x^2+1} + \frac{B}{x^2+2} = \frac{x^2(A+B) + 2A+B}{(x^2+1)(x^2+2)},$$

from which

$$A + B = 1, 2A + B = 0, \text{ that is, } A = -1, B = 2.$$

Therefore

$$\begin{aligned} \int f(x) dx &= \int \left(\frac{-1}{x^2+1} + \frac{2}{x^2+2} \right) dx = -\arctan x + \int \frac{1}{\left(\frac{x}{\sqrt{2}}\right)^2 + 1} dx \\ &= -\arctan x + \sqrt{2} \int \frac{1}{t^2+1} dt = -\arctan x + \sqrt{2} \arctan \frac{x}{\sqrt{2}} + k, k \in \mathbb{R}. \end{aligned}$$

b) The integrand is continuo in $[0, +\infty[$, so that bisogna controllare the convergence of the integral solo for $x \rightarrow +\infty$. Siccome the integrand is positive, usiamo the criterion of the asymptotic comparison . One has

$$\log \frac{x^\alpha + 2}{x^\alpha + 1} = \log \left(1 + \frac{2}{x^\alpha} \right) - \log \left(1 + \frac{1}{x^\alpha} \right) = \frac{1}{x^\alpha} + o\left(\frac{1}{x^\alpha}\right)$$

for $x \rightarrow +\infty$. Therefore the integral converges if and only if $\alpha > 1$.

c) Integrando by parts one has

$$\begin{aligned} \int_0^c \log \frac{x^2+2}{x^2+1} dx &= x \log \frac{x^2+2}{x^2+1} \Big|_0^c - \int_0^c x \frac{x^2+1}{x^2+2} \frac{2x(x^2+1)-2x(x^2+2)}{(x^2+1)^2} dx \\ &= c \log \frac{c^2+2}{c^2+1} - \int_0^c \frac{-2x^2}{(x^2+2)(x^2+1)} dx = \quad [\text{tenendo conto of the calcolo fatto in a}] \\ &= c \log \frac{c^2+2}{c^2+1} + 2 \left(-\arctan c + \sqrt{2} \arctan \frac{c}{\sqrt{2}} \right). \end{aligned}$$

Therefore

$$\int_0^{+\infty} \log \frac{x^2+2}{x^2+1} dx = \lim_{c \rightarrow +\infty} \left(c \log \frac{c^2+2}{c^2+1} + 2 \left(-\arctan c + \sqrt{2} \arctan \frac{c}{\sqrt{2}} \right) \right) = \pi(\sqrt{2}-1),$$

in quanto

$$\lim_{c \rightarrow +\infty} c \log \frac{c^2+2}{c^2+1} = \lim_{c \rightarrow +\infty} c \left(\frac{1}{c^2} + o\left(\frac{1}{c^2}\right) \right) = 0.$$

Appello of the 17.09.2018

THEME 1

Exercise 1 Consider the function

$$f(x) := \begin{cases} e^{-\frac{2}{|x|}} (2|x| - 3) & \text{for } x \neq 0 \\ 0 & \text{for } x = 0. \end{cases}$$

- i) Determine the domain D , le simmetries and study the sign of f ;
- ii) In order to determine i limits of f at the extremes of D and the asymptotes;
- iii) compute the derivative and study the monotonicity of f and determine the points of extreme relative and absolute ; the study of the second derivative may be skipped ;
- iv) study the continuity and () the derivability of f (in particolare in $x = 0$);
- v) draw a qualitative graph of f .

Solution. i) $D = \mathbb{R}$, ovviamente and the function is pari. One has

$$f(x) \geq 0 \text{ if and only if } |x| \geq \frac{3}{2} \text{ oppure } x = 0.$$

D'ora in study f for $x \geq 0$.

ii) One has

$$\lim_{x \rightarrow +\infty} f(x) = \lim_{x \rightarrow +\infty} e^{-\frac{2}{x}} (2x - 3) = +\infty.$$

Per the calcolo of the asintoto one has

$$\lim_{x \rightarrow +\infty} \frac{f(x)}{x} = \lim_{x \rightarrow +\infty} e^{-\frac{2}{x}} \frac{2x - 3}{x} = 2$$

e

$$\lim_{x \rightarrow +\infty} f(x) - 2x = \lim_{x \rightarrow +\infty} \left(2x(e^{-\frac{2}{x}} - 1) - 3e^{-\frac{2}{x}} \right) = \lim_{x \rightarrow +\infty} \left(2x \left(-\frac{2}{x} + o\left(\frac{1}{x}\right) \right) - 3e^{-\frac{2}{x}} \right) = -7,$$

so that the retta $y = 2x - 7$ is oblique asymptote for $x \rightarrow +\infty$.

iii) Per $x > 0$ si posthey are applicare le regole diderivazione, dato that one has $f(x) = e^{-\frac{2}{x}}(2x - 3)$. Therefore

$$f'(x) = 2e^{-\frac{2}{x}} + \frac{2e^{-\frac{2}{x}}}{x^2}(2x - 3) = \frac{2e^{-\frac{2}{x}}}{x^2}(x^2 + 2x - 3).$$

One has therefore che $f'(x) \geq 0$ if and only if $x^2 + 2x - 3 \geq 0$, that is, (for $x > 0$) if and only if $x \geq 1$. Therefore $x = 1$ is the point of absolute minimum, and is a minimum stretto, while $x = 0$ is a point of relative maximum stretto, in quanto $f(x) < 0 = f(0)$ for $0 < |x| < \frac{3}{2}$ (mostrato in (i)).

iv) The function is continuous in $]0, +\infty[$ in quanto superposition difunctions elementari. Per study the continuity in 0 bisogna compute

$$\lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} e^{-\frac{2}{x}}(2x - 3) = -3 \lim_{x \rightarrow 0^+} e^{-\frac{2}{x}} = 0 = f(0).$$

Therefore f is continuous also in $x = 0$. Per study the derivability in $x = 0$ one may compute the limit

$$\lim_{x \rightarrow 0^+} f'(x) = \lim_{x \rightarrow 0^+} \frac{2e^{-\frac{2}{x}}}{x^2}(x^2 + 2x - 3) = -3 \lim_{x \rightarrow 0^+} \frac{2e^{-\frac{2}{x}}}{x^2} = 0$$

for the noto confronto tra esponenziali and potenze. Therefore f is differentiable also in $x = 0$ (and the derivative is continuous also in $x = 0$).

v) the graph of f is in the picture

Figure 13: the graph of f (Theme 1).

Exercice 2 Sia

$$P_\lambda(z) = \lambda - 4t\bar{z} + 2iz^2 + z^3.$$

Find $\lambda_0 \in \mathbb{C}$ in modo che $z = -2i$ sia a zero of P_{λ_0} . Solve the equation

$$P_{\lambda_0}(z) = 0$$

and express the solutions in algebraic form.

Solution. $P_\lambda(-2i) = \lambda - 8 - 8i + 8i$, from which $P_\lambda(-2i) = 0$ if and only if $\lambda = 8$. the polynomial dicui trovare the zeros is hence $P_{\lambda_0}(z) = 8 - 4\lambda z + 2iz^2 + z^3$. Siccome $z = -2i$ is a zero of P , P is divisible for $z + 2i$ and one has, in particolare,

$$P_{\lambda_0}(z) = (z + 2i)(z^2 - 4i).$$

Le altre solutions of the equation $P_{\lambda_0}(z) = 0$ they are therefore le two radici quadrate of $4i = 4e^{i\frac{\pi}{2}}$, that is, they are

$$\pm 2e^{i\frac{\pi}{4}} = \pm \sqrt{2}(1+i).$$

Exercise 3 Discutere for all values of the real parameter α the convergence of the series

$$\sum_{n=2}^{\infty} \frac{\log(n + \sin n)}{n^{\frac{\alpha}{2}} + 2}$$

Solution. The series is a termini positivi and one may hence usare the criterion of the asymptotic comparison . One has

$$\log(n + \sin n) \sim \log n \text{ for } n \rightarrow \infty,$$

perché

$$\lim_{n \rightarrow \infty} \frac{\log(n + \sin n)}{\log n} = \lim_{n \rightarrow \infty} \frac{\log n + \log\left(1 + \frac{\sin n}{n}\right)}{\log n} = \lim_{n \rightarrow \infty} \frac{\log n + \frac{\sin n}{n} + o\left(\frac{1}{n}\right)}{\log n} = 1.$$

Furthermore,

$$\frac{1}{n^{\frac{\alpha}{2}} + 2} \sim \frac{1}{n^{\frac{\alpha}{2}}} \text{ for } n \rightarrow \infty.$$

The series converges therefore if and only if converges the series

$$\sum_{n=1}^{\infty} \frac{\log n}{n^{\frac{\alpha}{2}}}.$$

Quest'ultima converges if and only if $\frac{\alpha}{2} > 1$, that is, if and only if $\alpha > 2$. Infatti, if $\frac{\alpha}{2} \leq 1$, the general term of the series is $\geq \frac{1}{n}$ and hence the series diverges. If instead $\frac{\alpha}{2} > 1$ and scelgo $1 < \beta < \frac{\alpha}{2}$, allora, for $n \rightarrow \infty$,

$$\frac{\log n}{n^{\frac{\alpha}{2}}} = o\left(\frac{1}{n^\beta}\right),$$

dal limit fondamentale

$$\lim_{n \rightarrow \infty} \frac{\log n}{n^\gamma} = 0 \text{ for every } \gamma > 0$$

and the series $\sum_{n=1}^{\infty} \frac{1}{n^\beta}$ converges.

Exercise 4 Compute as $\alpha \in \mathbb{R}^+$ the limit

$$\lim_{x \rightarrow 0^+} \frac{x - \sinh x - x^\alpha}{\cos x - 1 + x^{\frac{7}{3}} \log x}.$$

Solution. One has, for $x \rightarrow 0^+$,

$$x - \sinh x - x^\alpha = -\frac{x^3}{6} + o(x^3) - x^\alpha \sim \begin{cases} -x^\alpha & \text{if } \alpha < 3 \\ -\frac{7}{6}x^3 & \text{if } \alpha = 3 \\ -\frac{x^3}{6} & \text{if } \alpha > 3 \end{cases}$$

$$\cos x - 1 + x^{\frac{7}{3}} \log x = -\frac{x^2}{2} + o(x^2) + x^{\frac{7}{3}} \log x = -\frac{x^2}{2} + o(x^2) \sim -\frac{x^2}{2}$$

in quanto

$$\lim_{x \rightarrow 0^+} \frac{x^{\frac{7}{3}} \log x}{x^2} = \lim_{x \rightarrow 0^+} \sqrt[3]{x} \log x = 0.$$

Therefore ,

$$\lim_{x \rightarrow 0^+} \frac{x - \sinh x - x^\alpha}{\cos x - 1 + x^{\frac{7}{3}} \log x} = \begin{cases} +\infty & \text{if } \alpha < 2 \\ 2 & \text{if } \alpha = 2 \\ 0 & \text{if } \alpha > 2. \end{cases}$$

Exercise 5 Given the integral

$$\int_0^{\frac{1}{\sqrt{2}}} x^{\frac{\alpha}{2}} \arcsin 2x^2 dx,$$

a) study the convergence as $\alpha \in \mathbb{R}$;

b) calcolarlo for $\alpha = 2$.

Solution. a) The integrand $g(x) = x^{\frac{\alpha}{2}} \arcsin 2x^2$ is positive, so that one may use the criterion of the asymptotic comparison . One has, for $x \rightarrow 0^+$,

$$g(x) \sim 2x^{\frac{\alpha}{2}+2},$$

so that the integral converges if and only if $\frac{\alpha}{2} + 2 > -1$, that is, if and only if $\alpha > -6$.

b) One has

$$\begin{aligned} \int_0^{\frac{1}{\sqrt{2}}} x \arcsin 2x^2 dx &= (\text{by parts}) \quad \frac{x^2}{2} \arcsin 2x^2 \Big|_0^{\frac{1}{\sqrt{2}}} - \int_0^{\frac{1}{\sqrt{2}}} \frac{x^2}{2} \frac{4x}{\sqrt{1-4x^4}} dx \\ &= \frac{\pi}{8} - \int_0^{\frac{1}{\sqrt{2}}} \frac{2x^3}{\sqrt{1-4x^4}} dx = \frac{\pi}{8} + \frac{\sqrt{1-4x^4}}{4} \Big|_0^{\frac{1}{\sqrt{2}}} = \frac{\pi}{8} - \frac{1}{4}. \end{aligned}$$

Appello of the 21.01.2019

THEME 1

Exercise 1 Consider the function

$$f(x) = e^{\frac{|x^2 - 16|}{x+3}}, \quad x \in D =]-\infty, -3[.$$

- i) Determine the limits of f at the extremes of D and the asymptotes; study the prolongabilità for continuity in $x = -3$;
- ii) study the derivability, calcolarne the derivata, study the monotonicity and determine the points of extreme relative and absolute .

Solution.

- i) Let us observe that

$$\lim_{x \rightarrow -\infty} \frac{|x^2 - 16|}{x+3} = -\infty, \quad \lim_{x \rightarrow -3^-} \frac{|x^2 - 16|}{x+3} = -\infty$$

hence con a change of variable

$$\lim_{x \rightarrow -\infty} f(x) = \lim_{y \rightarrow -\infty} e^y = 0, \quad \lim_{x \rightarrow -3^-} f(x) = \lim_{y \rightarrow -\infty} e^y = 0.$$

In particolare f ha a horizontal asymptote ($y = 0$) for $x \rightarrow -\infty$. Furthermore, f può essere prolungata come function continuous da sinistra in -3 setting $f(-3) = 0$.

- ii) Calcoliamo, for $x \neq -4$,

$$\begin{aligned} f'(x) &= e^{\frac{|x^2 - 16|}{x+3}} \frac{d}{dx} \frac{|x^2 - 16|}{x+3} = e^{\frac{|x^2 - 16|}{x+3}} \frac{\operatorname{sgn}(x^2 - 16) 2x(x+3) - |x^2 - 16|}{(x+3)^2} \\ &= e^{\frac{|x^2 - 16|}{x+3}} \operatorname{sgn}(x^2 - 16) \frac{2x(x+3) - (x^2 - 16)}{(x+3)^2} \\ &= e^{\frac{|x^2 - 16|}{x+3}} \operatorname{sgn}(x^2 - 16) \frac{x^2 + 6x + 16}{(x+3)^2}, \end{aligned}$$

dove “sgn” indica the function sign.

Osservando che $e^{\frac{|x^2 - 16|}{x+3}} > 0$ and $(x+3)^2 > 0$ for every $x \in D$, vogliamo valutare the sign of

$$(x^2 + 6x + 16)\operatorname{sgn}(x^2 - 16)$$

Calcolando the discriminante of $x^2 + 6x + 16$, $\Delta = 36 - 64 < 0$ one gets $x^2 + 6x + 16 > 0$ for every x . Furthermore,

$$\operatorname{sgn}(x^2 - 16) > 0 \Leftrightarrow x^2 - 16 > 0 \Leftrightarrow |x| > 4 \Leftrightarrow x < -4 \text{ o } x > 4.$$

Since ci interessano solo the values of $x \in D$, ovvero $x < -3$, we get $\operatorname{sgn}(x^2 - 16) > 0$ for $x < -4$ and $\operatorname{sgn}(x^2 - 16) < 0$ for $-4 < x < -3$. Ne one has

$f'(x) > 0$ (and hence f increasing) for $x < -4$, $f'(x) < 0$ (and hence f decreasing) for $x \in]-4, -3[$,

from which segue che -4 è un punto assoluto massimo e per il teorema di Fermat, non posti altri punti di estremo.

Finally $x = -4$ è il punto unico in cui f non è differentiabile (è un punto angolare) perché

$$\lim_{x \rightarrow -4^+} f'(x) = -8 = -\lim_{x \rightarrow -4^-} f'(x).$$

the graph of f is in the picture ??.




Figure 14: the graph of f (Theme 1).

Exercise 2 Compute the limit

$$\lim_{x \rightarrow 0^+} \frac{e^{2x} - 1 - \sin(2x)}{\sinh^2 x + x^{\frac{9}{2}}}.$$

Solution. By making use of the Taylor development $e^y = 1 + y + \frac{y^2}{2} + o(y^2)$, $\sin y = y + o(y^2)$ con $y = 2x$ we get

$$e^{2x} = 1 + 2x + 2x^2 + o(x^2), \quad \sin 2x = 2x + o(x^2), \quad \text{for } x \rightarrow 0$$

and therefore the numerator can be written as

$$e^{2x} - 1 - \sin 2x = 2x^2 + o(x^2) \quad \text{for } x \rightarrow 0$$

Writing $\sinh x = x + o(x)$ we have $\sinh^2 x = (x + o(x))^2 = x^2 + o(x^2)$ for $x \rightarrow 0$. Furthermore, since $\frac{9}{2} > 2$, it is $x^{\frac{9}{2}} = o(x^2)$ for $x \rightarrow 0$. It follows

$$\sinh^2 x + x^{\frac{9}{2}} = x^2 + o(x^2).$$

Da cui

$$\lim_{x \rightarrow 0} \frac{e^{2x} - 1 - \sin 2x}{\sinh^2 x + x^{\frac{9}{2}}} = \lim_{x \rightarrow 0} \frac{2x^2 + o(x^2)}{x^2 + o(x^2)} = 2.$$

Exercise 3 Solve the equation

$$iz^2 + (1 + 2i)z + 1 = 0$$

in $z \in \mathbb{C}$, writing the solutions in algebraic form.

Solution. Vale

$$z = \frac{-1 - 2i + \sqrt{(1 + 2i)^2 - 4i}}{2i} = \frac{-1 - 2i + \sqrt{-3}}{2i},$$

dove $\sqrt{-3}$ denota le due radici complesse di -3 , that they are $\pm i\sqrt{3}$ (while $\sqrt{3}$ denota la radice quadrata positiva di 3). Questo one may verificare scrivendo le radici nella form ρe^{i0a} , richiedendo che

$$3 = 3e^{i0} = (\rho e^{i0a})^2 = \rho^2 e^{2i0a},$$

from which $\rho = \sqrt{3}$ and $0a = k\pi$ for $k \in \mathbb{Z}$. Abbiamo hence that le two radici they are

$$z_{\pm} = \frac{-1 - 2i \pm i\sqrt{3}}{2i} = -1 \pm \frac{\sqrt{3}}{2} + \frac{i}{2}.$$

Exercise 4

Siano $\alpha \in \mathbb{R}$ fissato and

$$f(t) := \frac{\log(1 + \frac{t}{2})}{t^{2\alpha}}.$$

- i) Compute $\int_1^2 f(t) dt$ con $\alpha = 1$.
- ii) Sia $F(x) := \int_2^x f(t) dt$ con $\alpha = \frac{1}{2}$. Scrivere the formula of Taylor of the second order for F centrata in $x = 2$.
- iii) Determine for which $\alpha \in \mathbb{R}$ esiste finito $\int_0^1 f(t) dt$.

Solution. i) Integriamo by parts and we get

$$\int_1^2 \frac{\log(1 + \frac{t}{2})}{t^2} dt = -\frac{\log(1 + \frac{t}{2})}{t} \Big|_1^2 + \int_1^2 \frac{1}{t(2+t)} dt = -\frac{\log 2}{2} + \log \frac{3}{2} + \int_1^2 \frac{1}{t(2+t)} dt$$

To compute the second integral usiamo the metodo of the fratti semplici:
poniamo

$$\frac{1}{t(2+t)} = \frac{A}{t} + \frac{B}{2+t} = \frac{2A + At + Bt}{t(2+t)},$$

from which $A = \frac{1}{2}$ and $B = -\frac{1}{2}$. In conclusione

$$\int_1^2 \frac{1}{t(2+t)} dt = \frac{1}{2} \int_1^2 \frac{1}{t} dt - \frac{1}{2} \int_1^2 \frac{1}{2+t} dt = \frac{1}{2} (\log t - \log(2+t)) \Big|_1^2 = \frac{1}{2} \log \frac{3}{2}.$$

In conclusione

$$\int_1^2 \frac{\log\left(1 + \frac{t}{2}\right)}{t^2} dt = -\frac{\log 2}{2} + \log \frac{3}{2} + \frac{1}{2} \log \frac{3}{2} = \log \frac{3\sqrt{3}}{4}.$$

ii) the polynomial di Taylor is

$$T_F^{2,2}(x) = F(2) + F'(2)(x-2) + \frac{F''(2)}{2}(x-2)^2,$$

therefore devo compute

$$F(0) = 0$$

$$F'(x) = \frac{d}{dx} \int_1^x f(t) dt = f(x) = \frac{\log\left(1 + \frac{x}{2}\right)}{x} \Rightarrow F'(2) = \frac{\log 2}{2},$$

e

$$F''(x) = f'(x) = \frac{\frac{1}{2+x}x - \log(1 + \frac{x}{2})}{x^2} \Rightarrow F''(2) = \frac{1}{8} - \frac{\log 2}{4}.$$

Ne segue

$$f(x) = \frac{\log 2}{2}(x-2) + \frac{1}{2} \left(\frac{1}{8} - \frac{\log 2}{4} \right) (x-2)^2 + o(x-2)^2$$

for $x \rightarrow 2$.

iii) Let us observe that for $\alpha \leq 0$ the function f is palesemente continuous and limitata su $[0, 1]$, so that the integral esiste finito. Per $\alpha > 0$ dobbiamo valutare the comportamento asymptotic of $f(t)$ for $t \rightarrow 0^+$, essendo comunque f continuous and limitata on ogni interval $[\delta, 1]$ for every $0 < \delta < 1$. Abbiamo

$$f(t) = \frac{\log\left(1 + \frac{t}{2}\right)}{t^{2\alpha}} = \frac{\frac{t}{2} + o(t)}{t^{2\alpha}} \sim \frac{1}{2t^{2\alpha-1}}, \quad \text{for } t \rightarrow 0^+.$$

Per the criterion of the asymptotic comparison it is hence

$$\int_0^1 f(t) dt \text{ converges} \Leftrightarrow \int_0^\delta \frac{1}{2t^{2\alpha-1}} dt \text{ converges for some } \delta > 0 \Leftrightarrow 2\alpha-1 < 1.$$

Hence the integral converges if and only if $\alpha < 1$.

Exercise 5 Study the convergence semplice and assoluta of the series

$$\sum_{n=0}^{+\infty} \frac{(\log \alpha)^n}{1 + \sqrt{2n}}$$

as $\alpha \in]0, +\infty[$.

Solution. Let us study the convergence of the series

$$\sum_{n=0}^{+\infty} \frac{y^n}{1 + \sqrt{2n}}.$$

Per $|y| < 1$ the series absolutely converges . Questo può essere easily provato usando the Root Test, essendo

$$\lim_{n \rightarrow \infty} \left(\frac{|y|^n}{1 + \sqrt{2n}} \right)^{\frac{1}{n}} = |y| < 1$$

oppure osservando che $n|y|^n \rightarrow 0$ for $|y| < 1$, hence $|y|^n \leq \frac{1}{n}$ definitively for $n \rightarrow \infty$ and visto that the series

$$\sum_{n=0}^{+\infty} \frac{1}{n(1 + \sqrt{2n})}$$

converges ($\frac{1}{n(1 + \sqrt{2n})} \sim \frac{1}{n^{\frac{3}{2}}}$) possiamo concludere usando the teorema of the confronto.

Per $|y| > 1$ the general term of the series diverges, hence the series non può convergere.

Per $y = 1$ the series diverges for asymptotic comparison con the series $\sum_{n=0}^{+\infty} \frac{1}{\sqrt{2n}}$.

Finally , for $y = -1$ the series converges for the criterio diLeibniz, essendo the modulo of the general term of the series decreasing a 0. Yet , for the case precedente, the series does not converge assolutamente.

Sostituendo $\log \alpha = y$ we get that the series originale absolutely converges if and only if $-1 < \log \alpha < 1$, ovvero if and only if $\frac{1}{e} < \alpha < e$, simply converges if and only if $-1 \leq \log \alpha < 1$, ovvero if and only if $\frac{1}{e} \leq \alpha < e$ and diverges in all the altri casi, ovvero $0 < \alpha < \frac{1}{e}$ and $\alpha \geq e$.

Exercise Determine all the values of $a \in \mathbb{R}$ such that the function $f(x) = e^x - ax^3$ sia convex in the whole \mathbb{R} .

Solution. Da $f''(x) = e^x - 6ax$, one has che f is convex if and only if

(A) $f''(x) = e^x - 6ax \geq 0$ for every $x \in \mathbb{R}$

Ora, if $a < 0$,

$$\lim_{x \rightarrow -\infty} f''(x) = \lim_{x \rightarrow -\infty} e^x - 66ax \geq 0 = -\infty$$

and hence (A) is not verified.

If $a = 0$ instead (A) is verified.

If $a > 0$ study the function $g(x) := f''(x) = e^x - 6ax$. One has $g'(x) = e^x - 6a \geq 0 \iff x \geq \log(6a)$. Hence g ha a absolute minimum in $x = \log(6a)$. Therefore (A) is verificata if and only if $g(\log(6a)) =$

$6a - 6a \log(6a) \geq 0$, that is, if and only if $1 - 1 \log(6a) \geq 0$, hence if and only if $a \leq \frac{e}{6}$.

In conclusione f is convex if and only if $a \leq \frac{e}{6}$.

xxxxxxxxxxxxxxxxxxxxxx

Appello of the 11.02.2019

THEME 1

Exercise 1. Sia

$$f(x) = |(x+3) \log(x+3)|, \quad x \in D =]-3, +\infty[.$$

- (i) Determine i limits of f at the extremes of D and the asymptotes; study the prolungabilità for continuity in $x = -3$;
- (ii) study the derivability, calcolarne the derivata, study the monotonicity, determine the points diextreme relative and absolute and draw the graph.

Solution.

- (i) Con the change of variable $y = x + 3$ we get

$$\lim_{x \rightarrow -3^+} f(x) = \lim_{y \rightarrow 0^+} |y \log y| = 0.$$

Questo in particolare implica che f one may prolungare for continuity in $x = -3$ setting $f(-3) = 0$.

Clearly vale

$$\lim_{x \rightarrow \infty} |x+3| = \infty, \quad \lim_{x \rightarrow \infty} |\log(x+3)| = \infty \quad \Rightarrow \quad \lim_{x \rightarrow \infty} f(x) = \infty.$$

D'altronde

$$\lim_{x \rightarrow \infty} \frac{f(x)}{x} = \lim_{x \rightarrow \infty} \frac{|x+3|}{x} |\log(x+3)| = 1 \cdot \lim_{x \rightarrow \infty} |\log(x+3)| = \infty,$$

hence the function non ha a oblique asymptote for $x \rightarrow \infty$.

(ii) Let us observe that in the domain D the function $(x+3) \log(x+3)$ si annullonly one for $x+3 = 1$, ovvero $x = -2$. Hence in $D \setminus \{-2\}$ the function f is differentiable in quanto prodotto and superposition didifferentiable functions, and si calcola

$$f'(x) = \operatorname{sgn}((x+3) \log(x+3))((x+3) \log(x+3))' = \operatorname{sgn}((x+3) \log(x+3))(\log(x+3)+1),$$

ovvero

$$f'(x) = -(\log(x+3) + 1) \text{ for } -3 < x < -2$$

$$f'(x) = \log(x+3) + 1 \text{ for } x > -2.$$

Si vede easily that $f'(x) > 0$ for every $x > -2$, hence f is strictly monotonic increasing for $x > -2$.

Per $-3 < x < -2$ vale

$$f'(x) > 0 \Leftrightarrow \log(x+3) < -1 \Leftrightarrow x+3 < \frac{1}{e} \Leftrightarrow x < -3 + \frac{1}{e}.$$

Con analoghi calcoli one has hence che

$$f'(x) > 0 \text{ for } -3 < x < -3 + \frac{1}{e}, \quad f'(x) = 0 \text{ for } x = -3 + \frac{1}{e}, \quad f'(x) < 0 \text{ for } -3 + \frac{1}{e} < x < -2.$$

Ne segue che f is strictly monotonic increasing for $-3 < x < -3 + \frac{1}{e}$ and strictly monotonic decreasing for $-3 + \frac{1}{e} < x < -2$.

Therefore $-3 + \frac{1}{e}$ is a point of local maximum , while -2 is a point of absolute minimum (infatti $f(x) \geq 0$ for every $x \in D$ and $f(-2) = 0$).

Si può easily osservare che

$$\lim_{x \rightarrow -2^-} f'(x) = -1, \quad \lim_{x \rightarrow -2^+} f'(x) = 1$$

and questo (for a teorema eventualmente visto a lezione) implica che f is not differentiable for $x = -2$.

the graph of f is in the picture

Exercise 2. Study the convergence of the series

$$\sum_{n=1}^{\infty} \frac{(1+n^2) \sin n}{n^4}$$

Solution. Let us observe that $|\sin n| \leq 1$ for every n , and hence

$$\sum_{n=1}^{\infty} \left| \frac{(1+n^2) \sin n}{n^4} \right| = \sum_{n=1}^{\infty} |\sin n| \left| \frac{1+n^2}{n^4} \right| \leq \sum_{n=1}^{\infty} \frac{1+n^2}{n^4}.$$

Since abbiamo

$$\frac{1+n^2}{n^4} \sim \frac{n^2}{n^4} = \frac{1}{n^2} \quad \text{for } n \rightarrow \infty,$$

o (equivalentemente) scrivendo

$$\frac{1+n^2}{n^4} = \frac{n^2(1+o(1/n^2))}{n^4} = \frac{1+o(1)}{n^2},$$

for the criterio of convergence asymptotic deduce that the series a termini positivi

$$\sum_{n=1}^{\infty} \frac{1+n^2}{n^4}$$

converges, and hence for the principio of the confronto the series originale absolutely converges .

Exercise 3 [4 punti] Solve the inequality

$$\frac{1}{2} \leq \frac{(\operatorname{Re}(\bar{z} + i) - 1)^2}{4} + \frac{(\operatorname{Im}(\bar{z} + i) - 1)^2}{4} \leq 1$$

and draw the solutions on Gauss plane .

Solution. Scriviamo in algebraic form $z = x + iy$, $\bar{z} = x - iy$. Hence

$$\operatorname{Re}(\bar{z} + i) = \operatorname{Re}(x - iy + i) = x, \quad \operatorname{Im}(\bar{z} + i) = \operatorname{Im}(x - iy + i) = 1 - y.$$

The inequality può essere therefore riscritta come

$$\frac{1}{2} \leq \frac{(x - 1)^2}{4} + \frac{(-y)^2}{4} \leq 1,$$

ovvero

$$2 \leq (x - 1)^2 + y^2 \leq 4.$$

Ricordando che $(x - x_0)^2 + (y - y_0)^2 = r^2$ is the equation of a circonferenza diraggio r centrata in (x_0, y_0) , we get that the inequality determina the corona circolare compresa tra le circonferenze diraggi $\sqrt{2}$ and 2 and centrate in $(1, 0)$.

the disign of the solutions is in the picture ??.




Figure 15: The soluzione of the exercise 3 (Theme 1).

Exercise 4. Compute

$$\int_0^{+\infty} e^{-\sqrt{2x}} dx.$$

Solution. By making use the change of variable $\sqrt{2x} = y$, from which $x = \frac{y^2}{2}$ and $dx = ydy$, we get

$$\int_0^{+\infty} e^{-\sqrt{2x}} dx = \int_0^{+\infty} e^{-y} y dy.$$

Integrand by parts one has

$$\int_0^{+\infty} e^{-y} y dy = [-e^{-y} y]_0^{+\infty} + \int_0^{+\infty} e^{-y} dy = 0 + [-e^{-y}]_0^{+\infty} = 0 - (-1) = 1.$$

Exercise 5. Sia

$$f_\alpha(x) = \frac{e^{-\sqrt{2x}} - 1}{x^{\alpha-1}}.$$

(a) study the convergence of the integral

$$\int_0^{+\infty} f_\alpha(x) dx$$

as $\alpha \in \mathbb{R}$.

(b) Per $\alpha = 2$, sia $F(x) = \int_1^{\cos x} f_\alpha(t) dt$: si calcoli $F'(\pi/3)$.

Solution. (a) Let us observe that the function f_α is continuous for $0 < x < +\infty$. Consideriamo

$$\int_0^1 f_\alpha(x) dx. \quad (2)$$

Essendo $e^{-\sqrt{2x}} = 1 - \sqrt{2x} + o(\sqrt{x})$ for $x \rightarrow 0$, abbiamo

$$f_\alpha(x) = \frac{-\sqrt{2x} + o(\sqrt{x})}{x^{\alpha-1}} = \frac{-\sqrt{2} + o(1)}{x^{\alpha-\frac{3}{2}}} \sim \frac{-\sqrt{2}}{x^{\alpha-\frac{3}{2}}},$$

hence, for the criterio of convergence asymptotic, the integral in (converges if and only if

$$\int_0^1 \frac{-\sqrt{2}}{x^{\alpha-\frac{3}{2}}} dx$$

converges, ovvero (portando $-\sqrt{2}$ fuori dall'integral) if and only if $\alpha - \frac{3}{2} < 1$, hence if and only if $\alpha < \frac{5}{2}$.

Let us study ora

$$\int_1^{+\infty} f_\alpha(x) dx. \quad (3)$$

Since $e^{-\sqrt{2x}} \rightarrow 0$ for $x \rightarrow \infty$ abbiamo

$$f_\alpha(x) \sim \frac{-1}{x^{\alpha-1}}$$

and for the criterio asymptotic of convergence, the integral in (converges if and only if

$$\int_1^{+\infty} \frac{-1}{x^{\alpha-1}} dx.$$

converges, ovvero if and only if $\alpha - 1 > 1$, hence if and only if $\alpha > 2$.

Therefore the integral originale converges if and only if $2 < \alpha < \frac{5}{2}$.

(b) Scriviamo

$$G(y) = \int_1^y f_2(t) dt = \int_1^y \frac{e^{-\sqrt{2t}} - 1}{t} dt.$$

Per the teorema fondamentale of the calcolo vale

$$G'(y) = f_2(y) = \frac{e^{-\sqrt{2y}} - 1}{y}.$$

Abbiamo $F(x) = G(\cos x)$. Per the chain rule, hence

$$F'(\pi/3) = G'(\cos(\pi/3))(-\sin(\pi/3)) = -\frac{\sqrt{3}}{2}G'(1/2) = -\frac{\sqrt{3}}{2} \frac{e^{-1} - 1}{1/2} = -\sqrt{3}(1 - 1/e).$$

Exercise 6 Compute the limit

$$\lim_{x \rightarrow 0^+} \frac{\cosh(\alpha x) - \cosh(e^{2x} - 1)}{x^3}$$

for all values of the parameter $\alpha > 0$.

Solution. Ricordiamo che $\cosh y = 1 + \frac{y^2}{2} + o(y^2)$, ed $e^y = 1 + y + o(y)$ for $y \rightarrow 0$, hence possiamo espandere the numerator come

$$\begin{aligned} \text{Num} &= 1 + \frac{\alpha^2 x^2}{2} + o(x^2) - \cosh(2x + o(x)) \\ &= 1 + \frac{\alpha^2 x^2}{2} + o(x^2) - \left[1 + \frac{(2x + o(x))^2}{2} + o((x + o(x))^2) \right] \\ &= 1 + \frac{\alpha^2 x^2}{2} + o(x^2) - [1 + 2x^2 + o(x^2)] \\ &= \frac{(\alpha^2 - 4)x^2}{2} + o(x^2). \end{aligned}$$

Therefore

$$\lim_{x \rightarrow 0^+} \frac{\cosh(\alpha x) - \cosh(e^{2x} - 1)}{x^3} = \lim_{x \rightarrow 0^+} \frac{\frac{\alpha^2 - 4}{2} + o(1)}{x} = \begin{cases} -\infty & \text{for } 0 < \alpha < 2 \\ +\infty & \text{for } \alpha > 2. \end{cases}$$

the case $\alpha = 2$ one has più difficile perchnd is not possibile compute $\lim_{x \rightarrow 0^+} \frac{o(1)}{x}$. Dobbiamo therefore ottenere un'expansion of the numerator

all'ordine successivo (the terzo). This volta scriviamo $\cosh y = 1 + \frac{y^2}{2} + o(y^3)$, ed $e^y = 1 + y + y^2 + o(y^2)$ for $y \rightarrow 0$. In particolare

$$\begin{aligned}\cosh(e^{2x} - 1) &= \cosh(2x + 2x^2 + o(x)^2) \\ &= 1 + \frac{(2x + 2x^2 + o(x)^2)^2}{2} + o((2x + 2x^2 + o(x)^2)^3) \\ &= 1 + \frac{4x^2 + 8x^3 + o(x^3)}{2} + o(x^3) \\ &= 1 + 2x^2 + 4x^3 + o(x^3).\end{aligned}$$

Per $\alpha = 2$ we have $\cosh(\alpha x) = 1 + 2x^2 + o(x^3)$, hence

$$\text{Num} = 1 + 2x^2 + o(x^3) - (1 + 2x^2 + 4x^3 + o(x^3)) = -4x^3 + o(x^3)$$

Therefore

$$\lim_{x \rightarrow 0^+} \frac{\cosh(2x) - \cosh(e^{2x} - 1)}{x^3} = \lim_{x \rightarrow 0^+} \frac{-4x^3 + o(x^3)}{x^3} = -4.$$

Exercise . Compute

$$\lim_{x \rightarrow +\infty} \int_x^{x+e^{-x}} e^t \arctan t dt.$$

Solution. Essendo the integrand continuous in a neighbourhood of $+\infty$ (in realtà in the whole \mathbb{R}), for the mean value theorem esiste $t_x \in [x, x + e^{-x}]$ tale che

$$\int_x^{x+e^{-x}} e^t \arctan t dt = e^{-x} e^{t_x} \arctan t_x$$

and hence, siccome the integrand is increasing,

$$e^{-x} e^x \arctan x \leq e^{-x} e^{t_x} \arctan t_x \leq e^{-x} e^{x+e^{-x}} \arctan(x + e^{-x}),$$

that is,

$$\arctan x \leq \int_x^{x+e^{-x}} e^t \arctan t dt \leq e^{e^{-x}} \frac{\pi}{2}.$$

Siccome

$$\lim_{x \rightarrow +\infty} e^{e^{-x}} = 1,$$

applicando the teorema of the Carabinieri one gets

$$\lim_{x \rightarrow +\infty} \int_x^{x+e^{-x}} e^t \arctan t dt = \frac{\pi}{2}.$$

Appello of the 8.07.2019

THEME 1

Exercise 1 [6 punti] Sia

$$f(x) = e^{\frac{2}{|2+\log x|}}.$$

- a) Determine the domain D of f ; determine i limits of f at the extremes of D and study the prolongabilità for continuity di f in $x = 0$;
- b) study the derivability, compute the derivative and study the monotonicity of f ; determine the points of extreme ;
- c) draw a qualitative graph of f .

Solution (a) Essendo the domain of e^x tutto \mathbb{R} , and the domain of $\log x$ all the $x > 0$, the domain of f is determinato dalle two condizioni:

$$x > 0, \quad 2 + \log x \neq 0.$$

The seconda relazione equivale a $x \neq e^{-2}$, hence

$$D = \{x > 0 : x \neq e^{-2}\}.$$

Con three cambi of variabile one gets

$$\lim_{x \rightarrow 0^+} f(x) = \lim_{y \rightarrow -\infty} e^{\frac{2}{|2+y|}} = \lim_{s \rightarrow +\infty} e^{\frac{2}{s}} = \lim_{t \rightarrow 0^+} e^t = 1.$$

Hence f può essere estesa for continuity in 0 setting $f(0) = 1$.

Furthermore,

$$\lim_{x \rightarrow e^{-2}} f(x) = \lim_{y \rightarrow 0^+} e^{\frac{2}{y}} = \lim_{s \rightarrow +\infty} e^s = +\infty.$$

Hence f non può essere estesa for continuity in e^{-2} .

- (b) The function is differentiable at all points of its domain, essendo superposition didifferentiable functions (the function $|\cdot|$ is not differentiable only at 0, ma $2 + \log x$ is zero only at e^{-2} , which doesn't belong to the domain.)
The derivata, calcolata con the chain rule is :

$$f'(x) = e^{\frac{2}{|2+\log x|}} \left(-\frac{2}{|2+\log x|^2} \right) \frac{2+\log x}{|2+\log x|} \frac{1}{x} = -\frac{2e^{\frac{2}{|2+\log x|}}}{x|2+\log x|^3} (2+\log x).$$

Notice that the fraction of the right-hand side is always positiva in the domain D , from which:

$$f'(x) > 0 \Leftrightarrow 2 + \log x < 0 \Leftrightarrow 0 < x < e^{-2},$$

$$f'(x) < 0 \Leftrightarrow 2 + \log x > 0 \Leftrightarrow x > e^{-2},$$

ed $f'(x) \neq 0$ for every $x \in D$. In particular, f non ha punti critical .




Figure 16: the graph of f (Theme 1).

the graph of f is in the picture ??.

Exercise 2 [4 punti] Study the convergence of the series

$$\sum_{n=1}^{\infty} \frac{\sin \frac{1}{n}}{1 - 2\sqrt{n}}.$$

Solution Per $n \rightarrow \infty$ we have $\sin \frac{1}{n} \sim \frac{1}{n}$ and $1 - 2\sqrt{n} \sim -2\sqrt{n}$. Therefore the general term of the series is asymptotic to $\frac{-1}{2n^{\frac{3}{2}}}$. Since $\frac{3}{2} > 1$, the series

$$\sum_{n=1}^{\infty} \frac{-1}{2n^{\frac{3}{2}}}$$

converges, and for the principio of convergence asymptotic, also the prima series converges.

Exercise 3 [4 punti] Solve the equation

$$\frac{z}{\bar{z}} = -\frac{(\operatorname{Im} z)^2}{|iz^2|}$$

and draw the solutions on Gauss plane .

Solution Let us observe that the equation is ben definitonly one for $z \neq 0$. Therefore, assumendo $z \neq 0$ possiamo semplificare moltiplicando a sinistra

for $\frac{z}{\bar{z}}$, ottenendo

$$\frac{z}{\bar{z}} = \frac{z^2}{|z|^2} = -\frac{(\operatorname{Im} z)^2}{|z|^2},$$

where we have also usato che $|iz^2| = |z^2| = |z|^2$. Moltiplicando per $|z|^2$ we get

$$z^2 = -\operatorname{Im} z^2.$$

Scrivendo $z = x + iy$ one gets $z^2 = x^2 - y^2 + 2ixy$, $-(\operatorname{Im} z)^2 = -y^2$, and we get hence

$$x^2 - y^2 + 2ixy = -y^2. \quad (4)$$

Quest'equation può essere soddisfatto only one if $2ixy = 0$. Questo implica $xy = 0$, ovvero $x = 0$ o $y = 0$ (non entrambi perché $z \neq 0$). In the case $x = 0$, $y \neq 0$, we have a soluzione of (. In the case $y = 0$ and $x \neq 0$ instead () is not soddisfatta. Therefore the solutions they are

$$\{z = x + iy \in \mathbb{C} : x = 0, y \neq 0\},$$

ovvero the asse immaginario privato of the origine, come one may vedere in the picture

Exercise 4 [5+3+4 punti] a) Compute a primitive of the function

$$e^x \log(1 + 2e^x).$$

Per $\alpha \in \mathbb{R}$, define $f_\alpha(x) = e^{\alpha x} \log(1 + 2e^x)$:

b) study the convergence of the generalized integral

$$\int_0^{+\infty} f_\alpha(x) dx$$

as $\alpha \in \mathbb{R}$;

c) find the Taylor expansion of order 2 centered in $x_0 = 1$ of the function

$$F(x) = \int_1^x f_0(t) dt.$$

Solution a) Con the sostituzione $y = e^x$, $dy = e^x dx$ and un'integration by parts one gets

$$\int e^x \log(1 + 2e^x) dx = \int \log(1 + 2y) dy = y \log(1 + 2y) - \int \frac{2y}{1 + 2y} dy.$$

Ora, for ridurre the numerator of the integrand a destra scriviamo

$$\frac{2y}{1 + 2y} = 1 - \frac{1}{1 + 2y},$$




Figure 17: The insieme of the solutions of the exercise 3 (Theme 1).

Hence

$$\int \frac{2y}{1+2y} dy = \int \left(1 - \frac{1}{1+2y}\right) dy = y - \frac{\log(1+2y)}{2}.$$

Aggiungendo ai termini precedenti and sostituendo $y = e^x$ one gets

$$\int e^x \log(1+2e^x) dx = e^x \log(1+2e^x) - e^x + \frac{\log(1+2e^x)}{2} + c.$$

b) Per ogni $\alpha \in \mathbb{R}$ the function f_α is continuous in $[0, +\infty)$, hence for study the convergence of the suo integral, study the comportamento of f_α for $x \rightarrow \infty$. Per $\alpha \geq 0$ we have che

$$\lim_{x \rightarrow \infty} f_\alpha(x) = +\infty,$$

so that the integral $\int_0^{+\infty} f_\alpha(x) dx$ diverges.

Per $\alpha < 0$, we have $f_\alpha(x) = O(x^{-2})$ for $x \rightarrow \infty$, and for the criterion of the asymptotic comparison, the integral converges.

c) Abbiamo

$$F(x) = F(1) + F'(1)(x-1) + \frac{F''(1)}{2}(x-1)^2 + o(|x-1|^2) \quad \text{for } x \rightarrow 1.$$

Abbiamo

$$F(1) = 0, \quad F'(1) = f_0(1) = \log(1+2e), \quad f_0'(x) = \frac{2e^x}{1+2e^x}, \quad f_0'(1) = \frac{2e}{1+2e},$$

hence

$$F(x) = \log(1+2e)(x-1) + \frac{e}{1+2e}(x-1)^2 + o(|x-1|^2) \quad \text{for } x \rightarrow 1.$$

Exercise 5 [6 punti] Compute the limit

$$\lim_{x \rightarrow +\infty} x^\alpha \left(\sqrt[8]{x^2 - 2} - \sqrt[4]{x + 1} \right)$$

for all values of the parameter $\alpha > 0$.

Solution. Let us utilize the expansion $(1+y)^\alpha = 1 + \alpha y + o(y)$ for $y \rightarrow 0$ and scriviamo

$$\sqrt[8]{x^2 - 2} = (x^2 - 2)^{\frac{1}{8}} = x^{\frac{1}{4}} \left(1 - \frac{2}{x^2} \right)^{\frac{1}{8}} = x^{\frac{1}{4}} \left(1 - \frac{1}{4x^2} + o\left(\frac{1}{x^2}\right) \right), \quad \text{for } x \rightarrow \infty,$$

$$\sqrt[4]{x - 1} = (x - 1)^{\frac{1}{4}} = x^{\frac{1}{4}} \left(1 - \frac{1}{x} \right)^{\frac{1}{4}} = x^{\frac{1}{4}} \left(1 - \frac{1}{4x} + o\left(\frac{1}{x}\right) \right), \quad \text{for } x \rightarrow \infty.$$

Sottraendo we get

$$x^\alpha \left(\sqrt[8]{x^2 - 2} - \sqrt[4]{x - 1} \right) = x^\alpha \cdot x^{\frac{1}{4}} \left(-\frac{1}{4x} + o\left(\frac{1}{x}\right) \right) = -\frac{x^{\alpha - \frac{3}{4}}}{4}(1 + o(1)) \quad \text{for } x \rightarrow \infty.$$

Therefore

$$\lim_{x \rightarrow +\infty} x^\alpha \left(\sqrt[8]{x^2 - 2} - \sqrt[4]{x - 1} \right) = \lim_{x \rightarrow +\infty} \left(-\frac{x^{\alpha - \frac{3}{4}}}{4}(1 + o(1)) \right) = \begin{cases} 0 & \text{for } \alpha < \frac{3}{4} \\ -\frac{1}{4} & \text{for } \alpha = \frac{3}{4} \\ -\infty & \text{for } \alpha > \frac{3}{4}. \end{cases}$$

Appello of the 17.09.2019

Theme 1

Exercise 1. Sia

$$f(x) = \log |e^{3x} - 2|.$$

- a) Determine the domain D and study the sign of f ; determine i limits of f at the extremes of D and determine the asymptotes;
- b) study the derivability, compute the derivative and study the monotonicity of f ; determine the points of extreme relative and absolute ;
- c) draw a qualitative graph of f .

Solution. a) Clearly $D = \{x \in \mathbb{R} : |e^{3x} - 2| > 0\} = \{x \in \mathbb{R} : e^{3x} - 2 \neq 0\} = \mathbb{R} \setminus \{\frac{\log 2}{3}\}$. Segno:

$$f(x) \geq 0 \iff |e^{3x} - 2| \geq 1 \iff e^{3x} - 2 \leq -1 \text{ and } e^{3x} - 2 \geq 1 \iff x \leq 0, \text{ and } x \geq \frac{\log 3}{3}.$$

When = one has also the zeros of f . Limits and asymptotes: we have to study the function for $x \rightarrow \pm\infty, \frac{\log 2}{3}$. Easily one has $f(-\infty) = \log 2$, hence $y = \log 2$ is horizontal asymptote at $-\infty$. A $+\infty$ easily $f(+\infty) = +\infty$. Cerchiamo a oblique asymptote $y = mx + q$. As for m , we have che

$$m = \lim_{x \rightarrow +\infty} \frac{f(x)}{x} = \lim_{x \rightarrow +\infty} \frac{\log(e^{3x} - 2)}{x} = \lim_{x \rightarrow +\infty} \frac{\log(e^{3x} \cdot 1_x)}{x} = \lim_{x \rightarrow +\infty} \frac{3x + 0_x}{x} = 3.$$

As for q , we have

$$q = \lim_{x \rightarrow +\infty} (f(x) - 3x) = \lim_{x \rightarrow +\infty} (\log(e^{3x} - 2) - \log e^{3x}) = \lim_{x \rightarrow +\infty} \log \frac{e^{3x} - 2}{e^{3x}} = \log 1 = 0.$$

Conclusion: $y = 3x$ is oblique asymptote at $+\infty$. Finally ,

$$\lim_{x \rightarrow \frac{\log 2}{3}} \log |e^{3x} - 2| = \log 0+ = -\infty,$$

from which $x = \frac{\log 2}{3}$ is vertical asymptote.

- b) Clearly f is continuous sul proprio domain essendo superposition dicontinuous functions ove definite. È also differentiable poiché the unique point in which one may not applicare the chain rule is x t.c. $e^{3x} - 2 = 0$, that is, $x = \frac{\log 2}{3}$, that for ò non appartiene al domain of f . The derivative is


$$f'(x) = \frac{1}{|e^{3x} - 2|} \operatorname{sgn}(e^{3x} - 2) \cdot 3e^{3x} = \frac{3e^{3x}}{e^{3x} - 2}.$$

Da this segue che

$$f'(x) \geq 0, \iff e^{3x} - 2 > 0, \iff x > \frac{\log 2}{3}.$$

One concludes that $f \searrow$ su $]-\infty, \frac{\log 2}{3}[$ while $f \nearrow$ su $\frac{\log 2}{3}, +\infty[$. There are no , conseguenza nán minimi nán massimi (diqualsiasi natura).

- c) Grafico.



Exercise 2. Compute the limit

$$\lim_{x \rightarrow 0^+} \frac{e^{x-2x^2} - 1 - x}{\sinh x^2 + x^{7/3} \log x}.$$

Solution. Since $\lim_{x \rightarrow 0^+} x^\alpha \log x = 0$, si vede easily that the limit si presenta come a form of the tipo 0/0. Let us study the ordine di infinitesimali dinumeratore and denominator . Since

$$e^t = 1 + t + o(t) = 1 + t + \frac{t^2}{2} + o(t^2),$$

abbiamo

$$N = 1 + (x - 2x^2) + o(x - x^2) - 1 - x = -2x^2 + o(x) = o(x),$$

insufficiente for at comportamento preciso,

$$N = 1 + (x - 2x^2) + \frac{(x - 2x^2)^2}{2} + o((x - 2x^2)^2) - 1 - x = -2x^2 + \frac{x^2}{2} + o(x^2) \sim -\frac{3}{2}x^2 \text{ for } x \rightarrow 0^+.$$

Per the denominator is sufficiente ricordare che $\sinh t = t + o(t)$ so that

$$D = x^2 + o(x^2) + x^{7/3} \log x = x^2 + o(x^2) \text{ for } x \rightarrow 0^+,$$

essendo $x^{7/3} \log x = o(x^2)$ poiché $\frac{x^{7/3} \log x}{x^2} = x^{1/3} \log x \rightarrow 0$ for $x \rightarrow 0^+$.

Hence

$$\frac{N}{D} \sim \frac{-\frac{3}{2}x^2}{x^2} \rightarrow -\frac{3}{2}.$$

Exercise 3. Solve the inequality

$$\operatorname{Re} z \leq \operatorname{Re} \left(\frac{3}{z} \right)$$

and draw the solutions on Gauss plane .


Solution. Sia $z = x + iy$ con $x, y \in \mathbb{R}$. Then $z = x$ while essendo $\frac{1}{z} = \frac{1}{x+iy} = \frac{x-iy}{x^2+y^2}$,

$$\frac{3}{z} = \frac{3x}{x^2 + y^2}.$$

Therefore, $z \neq 0$ verifica la disequazione se e solo se

$$x \leq \frac{3x}{x^2 + y^2} \iff \begin{cases} x > 0, & 1 \leq \frac{3}{x^2 + y^2}, \iff x^2 + y^2 \leq 3, \\ x = 0, & \forall y \in \mathbb{R} \setminus \{0\}, \\ x < 0, & 1 \geq \frac{3}{x^2 + y^2}, \iff x^2 + y^2 \geq 3. \end{cases}$$

Figura:



Exercise 4. a) Compute the indefinite integral

$$\int \left(\tan \frac{x}{2} \right)^3 dx \quad (\text{sugg.: eseguire la sostituzione } \tan \frac{x}{2} = u).$$

b) study the convergence of the generalized integral

$$\int_0^{\frac{\pi}{6}} \frac{\tan x}{x^{\alpha+2}} dx$$

as $\alpha \in \mathbb{R}$.

Solution. a) Seguendo the hint $u = \tan x/2$, $x = 2 \arctan u$ from which $dx = \frac{2}{1+u^2}$, therefore

$$\begin{aligned} \int (\tan \frac{x}{2})^3 dx &= \int \frac{2u^3}{1+u^2} du = 2 \int \frac{u(u^2+1-1)}{1+u^2} du = \int 2u - \frac{2u}{1+u^2} du = u^2 - \log(1+u^2) \\ &= (\tan \frac{x}{2})^2 - \log \left(1 + (\tan \frac{x}{2})^2 \right). \end{aligned}$$

b) Sia $f(x) = \frac{\tan x}{x^{\alpha+2}}$. Certamente $f \in C([0, \frac{\pi}{6}])$ for every α and is continuous anche in $x = 0$ (hence integrabile sicuramente) for $\alpha + 2 \leq 0$, that is, for $\alpha \leq -2$. Per $\alpha > -2$ we have at generalized integral in $x = 0$. Since $\tan x = x + o(x) = x1_x$ for $x \rightarrow 0$,

$$f(x) \sim \frac{x}{x^{\alpha+2}} = \frac{1}{x^{\alpha+1}} \sim \frac{1}{x^{\alpha+1}} \text{ for } x \rightarrow 0^+,$$

integrabile in 0 if and only if $\alpha + 1 < 1$, that is, $\alpha < 0$ for asymptotic comparison. Morale: the generalized integral esiste finito if and solo if $\alpha < 0$.

Exercise 5. (i) Si dimostri that the sequence

$$a_n = \log(n+1) - \log \sqrt{n^2 + \alpha n + 4}$$

is infinitesimal for $n \rightarrow \infty$ (for every α) and for $\alpha = 2$ compute the order ;

(ii) study the convergence of the series

$$\sum_{n=2}^{\infty} a_n$$

as $\alpha \in \mathbb{R}$.

Solution. i) Let us observe that

$$a_n = \log \frac{n+1}{\sqrt{n^2 + \alpha n + 4}} \sim \log \frac{n}{n} \rightarrow 0.$$

In order to find the order diinfinitesimal l'occorre essere più precisi. Notiamo che, by dividing numerator and denominator by n , and usando le proprietà of the logarithms

$$a_n = \log \left(1 + \frac{1}{n} \right) - \frac{1}{2} \log \left(1 + \frac{\alpha}{n} + \frac{4}{n^2} \right).$$

Since $\log(1+t) = t + o(t) = t - \frac{t^2}{2} + o(t^2)$,

$$\begin{aligned} a_n &= \frac{1}{n} - \frac{1}{2n^2} + o\left(\frac{1}{n^2}\right) - \frac{1}{2} \left(\frac{\alpha}{n} + \frac{4}{n^2} - \frac{\left(\frac{\alpha}{n} + \frac{4}{n^2}\right)^2}{2} + o\left(\frac{1}{n^2}\right) \right) \\ &= \frac{2-\alpha}{2n} - \frac{5-\alpha^2}{2n^2} + o\left(\frac{1}{n^2}\right). \end{aligned}$$

In particolare, if $\alpha = 2$ one gets $a_n \sim -\frac{1}{2n^2}$.

ii) Per quanto visto al point i),

$$a_n \sim \begin{cases} \frac{2-\alpha}{2n} \equiv \frac{C}{n}, & \alpha \neq 2, \\ -\frac{1}{2n^2} \equiv \frac{C}{n^2}, & \alpha = 2, \end{cases}$$

from which si conclude che $\sum_n a_n$ converges if and only if $\alpha = 2$ in virtù of the criterion of the asymptotic comparison .

Appello of the 20.01.2020

THEME 1

Exercise 1 [7 punti] Consider the function

$$f(x) = \sin(2 \arctan(|x|^3))$$

- i) determine the domain D , the sign, simmetries, i limits at the extremes of D and the asymptotes;
- ii) study the derivability, calcolarne the derivata, study the monotonicity, determine the points of extreme relative and absolute ; the study of the second derivative may be skipped
- iii) draw the qualitative graph .

Solution. i) Clearly $D =]-\infty, +\infty[$. Clearly f is pari, hence basta limitarsi althe study su $[0, +\infty[$. Since $2 \arctan |x|^3 \in [0, \pi[$, f is always positiva and moreover $f = 0$ sse $x = 0$. Limits : there is only one interesting limit, $\lim_{x \rightarrow +\infty} f(x) = \sin \pi = 0$, from which the retta $y = 0$ is horizontal asymptote at $+\infty$.

ii) Essendo f superposition didifferentiable functions, eccetto for $x = 0$, one has

$$f'(x) = \cos(2 \arctan |x|^3) \frac{6x^2 \operatorname{sgn} x}{1 + x^6}, \quad \forall x \neq 0.$$

Per $x = 0$ chiaramente f is continuous and siccome

$$\lim_{x \rightarrow 0} f'(x) = 0,$$

for the test diderivability it follows that $\exists f'(0) = 0$. Per the monotonicity, study the sign of f' : for $x > 0$,

$$f'(x) \geq 0, \iff \cos(2 \arctan x^3) \geq 0, \iff 2 \arctan x^3 \leq \frac{\pi}{2}, \iff \arctan x^3 \leq \frac{\pi}{4}, \iff x^3 \leq 1,$$

grafici/1app1920_disigntema1.pdf

that is, for $x \leq 1$. Hence f is increasing su $[0, 1]$ and decreasing su $[1, +\infty[$. Si deduce easily the monotonicity su D and che $x = 0$ è punto di minimo globale while $x = \pm 1$ they are massimi globali.

Exercise 2 [6 punti] Compute the limit

$$\lim_{x \rightarrow 0^+} (1 + \sin x)^{x^a}$$

as $a \in \mathbb{R}$, usando the form “ $\exp\{\log \dots\}$ ”.

Solution. Per $x \rightarrow 0^+$, $1 + \sin x \rightarrow 1$ while

$$x^a \rightarrow \begin{cases} 0, & \text{if } a > 0, \\ 1, & \text{if } a = 0, \\ +\infty, & \text{if } a < 0. \end{cases}$$

Since $1^0 = 1$ and $1^1 = 1$ si deduce that the limit it is 1 for every $a \geq 0$. Per $a < 0$, $1^{+\infty}$ is indeterminate form. Since

$$(1 + \sin x)^{x^a} = e^{x^a \log(1 + \sin x)},$$

ricordato che $\log(1+t) = t1_t$ and che $\sin x = x1_x$ abbiamo

$$(1 + \sin x)^{x^a} = e^{x^a \sin x \cdot 1_x} = e^{x^{a+1} 1_x} \longrightarrow \begin{cases} e^0 = 1, & \text{if } -1 < a < 0, \\ e^1 = e, & \text{if } a = -1, \\ e^{+\infty} = +\infty, & \text{if } a < -1. \end{cases}$$

Exercise 3 [4 punti] Trovare the zeros in \mathbb{C} di

$$(z^3 + 5)(z^2 + z + 1) = 0.$$

Solution. Clearly

$$(z^3 + 5)(z^2 + z + 1) = 0, \iff z^3 = -5, \vee z^2 + z + 1 = 0.$$

In the first case , si tratta di compute le radici terze of -5 . Premesso che $-5 = 5u(\pi)$ ($u(0a) = \cos 0a + i \sin 0a$), for the formula di De Moivre, $z = \rho u(0a)$ is t.c.

$$z^3 = -5, \iff \begin{cases} \rho^3 = 5, \\ 0a = \frac{\pi}{3} + k\frac{2\pi}{3}, \quad k = 0, 1, 2, \end{cases} \iff z = \sqrt[3]{5} \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right), -\sqrt[3]{5}, \sqrt[3]{5} \left(\frac{1}{2} - i \frac{\sqrt{3}}{2} \right).$$

In the second case,

$$z_{1,2} = \frac{-1 \pm \sqrt{-3}}{2} = \frac{-1 \pm i\sqrt{3}}{2} = -\frac{1}{2} \pm i\frac{\sqrt{3}}{2}.$$

Exercise 4 [4+3 punti] Siano $\alpha \in \mathbb{R}$ fissato and

$$f_\alpha(t) := \frac{e^{2t} + 2e^t}{(e^t - 1)^\alpha}.$$

- i) Compute a primitive of f_α con $\alpha = 1$.
- ii) Determine for which $\alpha \in \mathbb{R}$ esiste finito $\int_0^1 f_\alpha(t) dt$.

Solution. i) Abbiamo che

$$\begin{aligned} \int \frac{e^{2t} + 2e^t}{e^t - 1} dt &\stackrel{u=e^t, t=\log u, dt=du/u}{=} \int \frac{u^2 + 2u}{(u-1)} \frac{du}{u} = \int \frac{u+2}{u-1} du = \int \left(1 + \frac{3}{u-1}\right) du \\ &= u + 3 \log|u-1| = e^t + 3 \log|e^t - 1|. \end{aligned}$$

ii) Considerato che $f_\alpha \in C([0, 1])$, the integral $\int_0^1 f_\alpha(t) dt$ is generalizzato in 0. Essendo $f_\alpha \geq 0$ su $[0, 1]$, possiamo applicare the test of the asymptotic comparison for stabilire the convergence of the integral. Notiamo che

$$f_\alpha(t) = \frac{3_t}{(e^t - 1)^\alpha} = \frac{3_t}{(t1_t)^\alpha} \sim_{0+} \frac{3}{t^\alpha},$$

so that esiste $\int_0^1 f_\alpha$ sse esiste $\int_0^1 \frac{1}{t^\alpha} dt$, sse $\alpha < 1$ come ben noto.

Exercise 5 [6 punti] Study the convergence semplice and assoluta of the series

$$\sum_{n=1}^{+\infty} \frac{(3 \sin x)^n n}{n^2 + \sqrt{n}}$$

as $x \in [-\frac{\pi}{2}, \frac{\pi}{2}]$.

Solution. Let us study the absolute convergence, that is, the convergence of the series

$$\sum_n |a_n| = \sum_n \frac{n 3^n |\sin x|^n}{n^2 + \sqrt{n}}.$$

A tal fine, let us apply the Root Test: essendo

$$|a_n|^{1/n} = \frac{n^{1/n} 3 |\sin x|}{n^{2/n} 1_n} \longrightarrow 3 |\sin x|, \quad \forall x \in [-\pi/2, \pi/2],$$

(ricordiamo che $n^{1/n} \longrightarrow 1$) we have che:

- if $3 |\sin x| < 1$ (that is, $|\sin x| < \frac{1}{3}$ ovvero, essendo $x \in [-\pi/2, \pi/2]$, sse $x \in]-\arcsin 1/3, \arcsin 1/3[$), the series absolutely converges (hence also semplicemente);
- if $3 |\sin x| > 1$ (that is, for $[-\pi/2, \pi/2] \setminus [-\arcsin 1/3, \arcsin 1/3]$), the series diverges assolutamente and poichnd the test dice in questo caso che $|a_n| \longrightarrow +\infty$, the condizione necessaria of convergence is not verificata, so that the series does not converge nemmeno semplicemente.

Rimangono the casi $\sin x = \pm \frac{1}{3}$, nei quali the test precedente fallisce. Per $\sin x = 1/3$, the series diventa

$$\sum_n \frac{n}{n^2 + \sqrt{n}} \sim \sum_n \frac{1}{n}, \text{ divergente.}$$

Since the terms have constant sign, convergence semplice and assoluta coincidono (hence there is no kind of convergence). Finally, for $\sin x = -1/3$,

$$\sum_n (-1)^n \frac{n}{n^2 + \sqrt{n}},$$

that is a series a termini disign alternato. The absolute convergence ritorna al case precedente (hence is esclusa). Per the convergence semplice possiamo applicare the test di Leibniz purch

$$\frac{n}{n^2 + \sqrt{n}} \searrow 0.$$

The convergence at 0 is evident. For the monotonicity we can proceed directly or introduce the auxiliary function $f(x) := \frac{x}{x^2 + \sqrt{x}}$ and observe that

$$f'(x) = \frac{x^2 + \sqrt{x} - x(2x + \frac{1}{2\sqrt{x}})}{(x^2 + \sqrt{x})^2} = \frac{-x^2 + \frac{\sqrt{x}}{2}}{(x^2 + \sqrt{x})^2}.$$

Since $f' \leq 0$ and $-x^2 + \frac{\sqrt{x}}{2} \leq 0$ whenever $x^{3/2} \geq \frac{1}{2}$, in particular for $n \geq 1$ one has $f(n) \searrow$, from which the conclusion: the series simply converges (but not absolutely) for the Leibniz test.

Exercise Sia $\{a_n\}$ a sequence such that $a_n > 0$ and $\frac{a_{n+1}}{a_n} \geq \frac{n}{n+1}$ for every $n \in \mathbb{N}$. Show that $\sum_{n=1}^{\infty} a_n$ diverges.

Solution. From the hypothesis it follows that $(n+1)a_{n+1} \geq na_n$, that is (na_n) is increasing; therefore $na_n \geq a_1 > 0$, from which $a_n \geq \frac{a_1}{n}$ for every $n \geq 1$. But then the series diverges by comparison with the harmonic series.

Time available: 2 hours and 45 minutes.

Appello of the 10.02.2020

THEME 1

Exercise 1 [7 points] Consider the function

$$f(x) = \exp \left\{ \left| \frac{x}{x+1} \right| \right\}.$$

- i) Find the domain D , the limits at the extremes of D and the asymptotes;
- ii) study the derivability, calculate the derivative, study the monotonicity, determine the points of extreme relative and absolute;
- iii) draw the qualitative graph.

Solution. i) Clearly $D = \mathbb{R} \setminus \{-1\}$. The limits at the extremes of D are

$$\lim_{x \rightarrow \pm\infty} f(x) = \lim_{t \rightarrow 1} e^t = e, \quad \lim_{x \rightarrow -1} f(x) = \lim_{t \rightarrow +\infty} e^t = +\infty.$$

Therefore, f has a horizontal asymptote of equation $y = e$ for $x \rightarrow \pm\infty$, and a vertical asymptote of equation $x = -1$ for $x \rightarrow -1$.

ii) f is composed of differentiable functions except where the denominator is zero, that is, f is surely differentiable in every $x \in D \setminus \{0\} = \mathbb{R} \setminus \{0, -1\}$. The point $x = 0$ is studied separately. We distinguish between the case in which $\frac{x}{x+1} > 0$, that is, $x > 0$ or $x < -1$, and the case in which $\frac{x}{x+1} < 0$, that is, $-1 < x < 0$.

- if $x \in]-\infty, -1[\cup]0, +\infty[$

$$f(x) = \exp \left\{ \frac{x}{x+1} \right\}$$

$$f'(x) = \exp \left\{ \frac{x}{x+1} \right\} \frac{d}{dx} \left(\frac{x}{x+1} \right) = \frac{1}{(x+1)^2} \exp \left\{ \frac{x}{x+1} \right\},$$

that is strictly positiva, Therefore therefore f is increasing su $]-\infty, -1[\cup]0, +\infty[$.

- If $x \in]-1, 0[$ one has

$$f(x) = \exp \left\{ -\frac{x}{x+1} \right\}$$

$$f'(x) = \exp \left\{ -\frac{x}{x+1} \right\} \frac{d}{dx} \left(-\frac{x}{x+1} \right) = -\frac{1}{(x+1)^2} \exp \left\{ -\frac{x}{x+1} \right\},$$

that is strictly negativa, Therefore therefore f is decreasing su $] -1, 0 [$.

Si vede che $\lim_{x \rightarrow 0^+} f'(x) = 1e^0 = 1$, while $\lim_{x \rightarrow 0^-} f'(x) = -1e^0 = -1$. Therefore f is not differentiable in $x = 0$, that is a angular point . Essendo D a union of intervals aperti, f può avere extremes locali solo where Therefore derivative si annulla and in points of non derivability. Come osservato sopra, $f'(x) \neq 0$, and the unique extreme si trova in $x = 0$, dove f ha the suo absolute minimum con $f(0) = 1$.

iii) Grafico:

Exercise 2 [5 punti] Study the convergence of the series

$$\sum_{k=1}^{\infty} 3^k \frac{k!}{k^k}.$$

Solution. The series has positive terms . let us apply the criterio of the rapporto asymptotic . One has

$$\frac{a_{k+1}}{a_k} = \frac{3^{k+1}(k+1)!}{(k+1)^{k+1}} \frac{k^k}{3^k k!} = \frac{3(k+1)k^k}{(k+1)(k+1)^k} = \frac{3}{(1+\frac{1}{k})^k} \rightarrow \frac{3}{e} \text{ for } k \rightarrow \infty.$$

Essendo $\frac{3}{e} > 1$, the series diverges for the criterio of the rapporto asymptotic

Exercise 3 [5 punti] Solve in \mathbb{C} nella form preferita (algebrica, esponenziale, trigonometrica):

$$z^3 = \frac{1}{-\frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2}i}.$$

Solution. Essendo $-\frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2}i = e^{\frac{5\pi}{4}i}$, the equation da risolvere diventa

$$z^3 = \frac{1}{e^{\frac{5\pi}{4}i}} = e^{-\frac{5\pi}{4}i} = e^{\frac{3\pi}{4}i}.$$

Per the teorema di De Moivre the solutions they are

$$z_0 = e^{\frac{\pi}{4}i} = \frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}i, \quad z_1 = e^{(\frac{\pi}{4} + \frac{2\pi}{3})i} = e^{\frac{11\pi}{12}i} = e^{-\frac{\pi}{12}i}, \quad z_2 = e^{(\frac{\pi}{4} + \frac{4\pi}{3})i} = e^{\frac{19\pi}{12}i} = e^{-\frac{5\pi}{12}i}$$

Applicando le formule di bisezione, one has

$$\cos\left(-\frac{\pi}{12}\right) = \sqrt{\frac{1 + \cos\left(-\frac{\pi}{6}\right)}{2}} = \sqrt{\frac{1 + \sqrt{3}/2}{2}},$$

$$\sin\left(-\frac{\pi}{12}\right) = -\sqrt{\frac{1 - \cos\left(-\frac{\pi}{6}\right)}{2}} = -\sqrt{\frac{1 - \sqrt{3}/2}{2}},$$

from which anche

$$\cos\left(-\frac{5\pi}{12}\right) = -\sqrt{\frac{1 + \sqrt{3}/2}{2}},$$

$$\sin\left(-\frac{5\pi}{12}\right) = -\sqrt{\frac{1 - \sqrt{3}/2}{2}},$$

cosicché

$$z_1 = \sqrt{\frac{1 + \sqrt{3}/2}{2}} - \sqrt{\frac{1 - \sqrt{3}/2}{2}}i \quad z_2 = -\sqrt{\frac{1 + \sqrt{3}/2}{2}} - \sqrt{\frac{1 - \sqrt{3}/2}{2}}i.$$

Exercise 4 [4+3 punti] Siano $\alpha \in \mathbb{R}$ and

$$f_\alpha(t) := \frac{e^{-2/t}}{3t^\alpha}.$$

- i) Compute a primitive of f_α con $\alpha = 3$.
- ii) Determine for which $\alpha \in \mathbb{R}$ esiste finito $\int_0^{+\infty} f_\alpha(t) dt$.

Solution. i) Con the sostituzione $y = -2/t$ one has $t = -2/y$, $dt = \frac{2}{y^2}dy$, and hence

$$\int f_3(t)dt = \int \frac{e^{-2/t}}{3t^3} = \int \frac{e^y}{3} \frac{-y^3}{8} \frac{2}{y^2} dy = -\frac{1}{12} \int ye^y dy.$$

Integrando by parts, one gets

$$\int f_3(t)dt = -\frac{1}{12} \int ye^y dy = -\frac{1}{12} \left(ye^y - \int e^y dy \right) = \frac{1}{12} (1-y)e^y = \frac{1}{12} \left(1 + \frac{2}{t} \right) e^{-2/t}.$$

ii) f_α is continuous su $(0, \infty)$. Per qualsiasi $\alpha \in \mathbb{R}$ one has (for the gerarchia degli infiniti) $\lim_{x \rightarrow 0^+} \frac{e^{-2/t}}{3t^\alpha} = 0$. Therefore, the function f_α può essere prolungata for continuity in $t = 0$, so that is, always integrabile in $[0, c]$, for qualsiasi $c > 0$. Per $t \rightarrow +\infty$, da $\frac{2}{t} \rightarrow 0$ one gets $e^{-2/t} \sim 1$ so that

$$f_\alpha(t) \sim \frac{1}{3t^\alpha},$$

and essendo f_α a sign constant, in virtù of the test of the asymptotic comparison, the integral esiste if and only if $\alpha > 1$.

Exercise 5 [6 punti] Compute the seguente limit

$$\lim_{x \rightarrow 0^+} \frac{\sin(x - x^3) - \log(1 + \sinh x) + \alpha x^2}{x^3}$$

as $\alpha \in \mathbb{R}$.

Solution. the limit is a indeterminate form 0/0. Analizziamo the numerator. Ricordando che (for $t \rightarrow 0$)

$$\sin t = t + o(t) = t - \frac{t^3}{6} + o(t^3), \quad \log(1+t) = t + o(t) = t - \frac{t^2}{2} + \frac{t^3}{3} + o(t^3), \quad \sinh t = t + o(t) = t + \frac{t^3}{6} + o(t^3),$$

si vede che (for $x \rightarrow 0^+$)

$$\begin{aligned} \text{Numerator} &= (x - x^3) - \frac{(x-x^3)^3}{6} + o((x-x^3)^3) \\ &\quad - \left(x + \frac{x^3}{6} + o(x^3) - \frac{1}{2} \left(x + \frac{x^3}{6} + o(x^3) \right)^2 + \frac{1}{3} \left(x + \frac{x^3}{6} + o(x^3) \right)^3 + o \left(\left(x + \frac{x^3}{6} + o(x^3) \right)^3 \right) \right) + \alpha x^2 \\ &= \left(\alpha + \frac{1}{2} \right) x^2 + \left(-1 - \frac{1}{6} - \frac{1}{6} - \frac{1}{3} \right) x^3 + o(x^3) = \left(\alpha + \frac{1}{2} \right) x^2 - \frac{5}{3} x^3 + o(x^3) \sim \left(\alpha + \frac{1}{2} \right) x^2 - \frac{5}{3} x^3. \end{aligned}$$

Si conclude allora che

$$\lim_{x \rightarrow 0^+} \frac{\text{Numerator}}{x^3} = \lim_{x \rightarrow 0^+} \left(\frac{\alpha + \frac{1}{2}}{x} - \frac{5}{3} \right) = \begin{cases} \infty, & \alpha > -\frac{1}{2}, \\ -\infty, & \alpha < -\frac{1}{2}, \\ -\frac{5}{3}, & \alpha = -\frac{1}{2}. \end{cases}$$

Exercise Sia $\alpha \in [0, +\infty[$ and define

$$F_\alpha(x) := \int_0^x t^\alpha e^{-t^2} dt, \quad x \geq 0.$$

Establish for which values of α one has che F_α is concave sull'interval $[1, +\infty[$.

There are values $\alpha > 0$ so that F_α sia concave su $[0, +\infty[$?

Solution. The function F_α is a function integral of $f_\alpha(t) := t^\alpha e^{-t^2}$. Essendo this ben defined and continuous su $[0, +\infty[$ (si ricorda $\alpha \geq 0$), anche F_α is ben defined , continuous and differentiable (for the teorema fondamentale of the calcolo) e

$$F'_\alpha(x) = f_\alpha(x) = x^\alpha e^{-x^2}.$$

Da this,

$$F''_\alpha(x) = e^{-x^2} (\alpha x^{\alpha-1} + x^\alpha (-2x)) = x^{\alpha-1} e^{-x^2} (\alpha - 2x^2).$$

Siccome F_α is twice differentiable, for a noto result

$$F_\alpha \text{ concave su } [1, +\infty[, \iff F''_\alpha(x) \leq 0, \quad \forall x \in [1, +\infty[.$$

Essendo

$$F''_\alpha(x) \leq 0, \iff \alpha - 2x^2 \leq 0, \stackrel{x \geq 0}{\iff} x \geq \sqrt{\frac{\alpha}{2}},$$

F_α is concave su $[1, +\infty[$ if and only if $\sqrt{\frac{\alpha}{2}} \leq 1$, that is, $\alpha \leq 2$. Lo stesso calcolo mostra che, for every $\alpha > 0$ one has $F''_\alpha(x) > 0$ for every $x \in [0, \sqrt{\frac{\alpha}{2}}[$, so that F_α non può essere concave su $[0, +\infty[$ for alcun value of $\alpha > 0$. Per $\alpha = 0$, one has che

$$F''_0(x) = -2xe^{-x^2} < 0 \quad \forall x > 0,$$

hence F_0 is concave su $[0, +\infty[$.

Tempo a disposizione: 2 ore and 45 minuti.

Appello of the 06.07.2020 - Modalità telematica (causa COVID)

THEME 1

Exercise 1 [6 punti] Consider the function

$$f(x) = |(x+3)\log(x+3)|, \quad x \in D =]-3, +\infty[.$$

(i) Compute

$$\lim_{x \rightarrow -3^+} f(x), \quad \lim_{x \rightarrow +\infty} f(x).$$

Solution.

$$\begin{aligned} \lim_{x \rightarrow -3^+} |(x+3) \log(x+3)| &= \lim_{x \rightarrow -3^+} -(x+3) \log(x+3) = \lim_{x \rightarrow -3^+} -\frac{\log(x+3)}{\frac{1}{x+3}} \quad (\text{De l'Hôpital}) \\ &= \lim_{x \rightarrow -3^+} x+3 = 0 \end{aligned}$$

$$\lim_{x \rightarrow +\infty} |(x+3) \log(x+3)| = \lim_{x \rightarrow +\infty} (x+3) \lim_{x \rightarrow +\infty} \log(x+3) = +\infty$$

(ii) Compute the first derivative of the function f , study the monotonicity intervals and draw the graph of f .

Solution. Per ogni x tale che $f(x) \neq 0$, that is,, for every $x \in D \setminus \{-2\}$,

$$\begin{aligned} f'(x) &= \operatorname{sgn}((x+3) \log(x+3)) (\log(x+3) + 1) \\ f'(x) \geq 0 &\iff \\ x \in \left\{ x \in D, (x+3) \log(x+3) > 0, \log(x+3) + 1 \geq 0 \right\} \cup \\ &\quad \cup \left\{ x \in D, (x+3) \log(x+3) < 0, \log(x+3) + 1 \leq 0 \right\} \\ &\iff \\ x \in \left\{ x > -2 \mid x \geq -3 + \frac{1}{e} \right\} \cup \\ &\quad \cup \left\{ -3 < x < -2, x \leq -3 + \frac{1}{e} \right\} = \\ &[-3, -3 + \frac{1}{e}] \cup [-2, +\infty[\end{aligned}$$

Therefore f is monotonic increasing in each of the $[-3, -3 + \frac{1}{e}]$ and $[-2, +\infty[$, while is monotonic decreasing in $-3 + \frac{1}{e}, -2[$. Therefore the function ha a localmaximum at the point $x = -3 + \frac{1}{e}$, where it is $f(-3 + \frac{1}{e}) = \frac{1}{e}$, and a minimum locale at the point $x = -2$, where it is $f(-2) = 0$. From the theorem of the right and left limit of the derivative one gets

$$f'_+(-2) = \lim_{x \rightarrow -2^+} f'(x) = 1 \quad f'_-(-2) = \lim_{x \rightarrow -2^-} f'(x) = -1.$$

Hence $x = -2$ is a angular point con tangent sinistra of equation $y = -x - 2$ and tangent destra of equation $y = x + 2$.

Exercise 2 [6 punti] Find the solutions of the equation

$$z^3 = 8i$$

in algebraic and exponential form (or trigonometric), and draw them on the Gauss plane .

Solution. Let us begin by writing $8i$ in trigonometric form :

$$8i = 8e^{i\frac{\pi}{2}}$$

Therefore $8i$ ha modulo $\rho = 8$ and argument $0a = \frac{\pi}{2}$. Solve l' equation means trovare le third roots of $8i$, that noi sappiamo essere in numero of three . Let us call them z_0, z_1, z_2 . One has

$$z_0 = \rho^{\frac{1}{3}} e^{i\frac{0a}{3}} = 2e^{i\frac{\pi}{6}} = 2 \left(\cos \frac{\pi}{6} + i \sin \frac{\pi}{6} \right) = \sqrt{3} + i$$

$$z_1 = \rho^{\frac{1}{3}} e^{i\left(\frac{0a}{3} + \frac{2\pi}{3}\right)} = 2e^{i\left(\frac{\pi}{6} + \frac{2\pi}{3}\right)} = 2e^{i\frac{5\pi}{6}} = 2 \left(\cos \frac{5\pi}{6} + i \sin \frac{5\pi}{6} \right) = -\sqrt{3} + i$$

$$z_2 = \rho^{\frac{1}{3}} e^{i\left(\frac{0a}{3} + \frac{4\pi}{3}\right)} = 2e^{i\left(\frac{\pi}{6} + \frac{4\pi}{3}\right)} = 2e^{i\frac{3\pi}{2}} = 2 \left(\cos \frac{3\pi}{2} + i \sin \frac{3\pi}{2} \right) = -2i$$

Come sapevamo già dalla teoria, the solutions on the Gauss plane are the vertexes dian equilateral triangle inscribed in a circle diradius 2. More precisely, one of the three vertexes si trova in $(0, -2)$ and an edge is a subset of the line $y = 1$.

Exercise 3 [6 punti] Study the convergence of the series

$$\sum_{n=1}^{\infty} \frac{(1+n^2) \log n}{n^4}.$$

Solution.

Si tratta of a series a termini positivi. Proviamo ad applicare the criterio of the rapporto:

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{\frac{(1+(n+1)^2) \log(n+1)}{(n+1)^4}}{\frac{(1+n^2) \log n}{n^4}} &= \lim_{n \rightarrow \infty} \frac{n^4}{(n+1)^4} \frac{(2+n^2+2n)}{(1+n^2)} \frac{\log(n+1)}{\log n} = \lim_{n \rightarrow \infty} \frac{\log(n+1)}{\log n} = \\ &= \lim_{n \rightarrow \infty} \frac{\log(n(1+1/n))}{\log n} = \lim_{n \rightarrow \infty} \frac{\log n + \log(1+1/n)}{\log n} = 1 \end{aligned}$$

Purtroppo siamo in the case in which the criterio of the rapporto non gives alc a informazione.

Tentiamo allora the strada of the confronto (asymptotic).

the factor $\frac{(1+n^2)}{n^4}$ is asymptotic to $\frac{1}{n^2}$, that fornirebbe a series converging. there is the factor $\log n$, that peggiora the situazione. Però noi sappiamo che, for $x \rightarrow \infty \log x = o(x^\alpha)$ for qualsiasi $\alpha > 0$: infatti

$$\lim_{x \rightarrow \infty} \frac{\log x}{x^\alpha} \stackrel{\text{(De the Hôpital)}}{=} \lim_{x \rightarrow \infty} \left(-\frac{1}{x} x^{-\alpha+1} \right) = \lim_{x \rightarrow \infty} x^{-\alpha} = 0.$$

Therefore, scegliendo ad esempio $\alpha = 1/2$, one has che

$$\frac{(1+n^2)\log n}{n^4} = o\left(\frac{(1+n^2)n^{\frac{1}{2}}}{n^4}\right) = o\left(\frac{1}{n^{\frac{3}{2}}}\right)$$

Since the series $\sum_{n=1}^{\infty} \frac{1}{n^{\frac{3}{2}}}$ is converging, for the criterio of the rapporto asymptotic si conclude that also the series data is converging .

Osservazione. Si sarebbe potuto scegliere a qualsiasi $\alpha \in]0, 1]$ al posto of $\alpha = \frac{1}{2}$. Invece the $\alpha \geq 1$ sarebbero stati inservibili, in quanto the series $\sum_{n=1}^{\infty} \frac{1}{n^{2-\alpha}}$ is divergente for $\alpha \geq 1$.

Exercice 4 [6 punti] Compute

$$\int_0^{+\infty} e^{-\sqrt{2x}} dx.$$

Solution. Per ogni $r > 0$, calcoliamo l' integral $\int_0^r e^{-\sqrt{2x}} dx$. Con the sostituzione $y(x) = \sqrt{2x}$, that is, $x(y) = \frac{y^2}{2}$ one gets

$$\int_0^r e^{-\sqrt{2x}} dx = \int_0^{\frac{r^2}{2}} e^{-y} \frac{d}{dy} \left(\frac{y^2}{2} \right) dy = \int_0^{\frac{r^2}{2}} e^{-y} y dy \stackrel{\text{(by parts)}}{=} [-e^{-y} y]_0^{\frac{r^2}{2}} + \int_0^{\frac{r^2}{2}} e^{-y} dy = -\frac{r^2 e^{-\frac{r^2}{2}}}{2} - e^{-\frac{r^2}{2}} + 1.$$

Hence

$$\int_0^{+\infty} e^{-\sqrt{2x}} dx = \lim_{r \rightarrow +\infty} \int_0^r e^{-\sqrt{2x}} dx = \lim_{r \rightarrow +\infty} \left(-\frac{r^2 e^{-\frac{r^2}{2}}}{2} - e^{-\frac{r^2}{2}} + 1 \right) = 1$$

$$(\text{perché } \lim_{r \rightarrow +\infty} \frac{r^2 e^{-\frac{r^2}{2}}}{2} = \lim_{r \rightarrow +\infty} \frac{r^2}{2 e^{\frac{r^2}{2}}} = 0)$$

Exercice 5 [6 punti] Compute the limit

$$\lim_{x \rightarrow +\infty} x^{\frac{4}{3}} \left(\sqrt[3]{x+2} - \sqrt[6]{x^2-1} \right)^2.$$

Solution.

$$\begin{aligned} \lim_{x \rightarrow +\infty} x^{\frac{4}{3}} \left(\sqrt[3]{x+2} - \sqrt[6]{x^2-1} \right)^2 &= \lim_{x \rightarrow +\infty} x^{\frac{4}{3}} \left(\sqrt[3]{x} \sqrt[3]{1 + \frac{2}{x}} - \sqrt[6]{x} \sqrt[6]{1 - \frac{1}{x^2}} \right)^2 = \\ &= \lim_{x \rightarrow +\infty} x^{\frac{5}{3}} \left(\sqrt[3]{1 + \frac{2}{x}} - \sqrt[6]{1 - \frac{1}{x^2}} \right)^2 = \lim_{x \rightarrow +\infty} x^{\frac{5}{3}} \left(\sqrt[3]{1 + \frac{2}{x}} - \sqrt[6]{1 - \frac{1}{x^2}} \right)^2. \end{aligned}$$

By making use lo sviluppo di Taylor

$$(1+y)^\alpha = 1 + \alpha y + o(y) \quad y \rightarrow 0$$

(valido study ogni $\alpha \in \mathbb{R}$), one gets

$$\begin{aligned} \lim_{x \rightarrow +\infty} x^{\frac{4}{3}} \left(\sqrt[3]{x+2} - \sqrt[6]{x^2-1} \right)^2 &= \lim_{x \rightarrow +\infty} x^{\frac{5}{3}} \left(1 + \frac{2}{3x} + o\left(\frac{1}{x}\right) - 1 + \frac{1}{6x^2} + o\left(\frac{1}{x^2}\right) \right)^2 = \\ &= \lim_{x \rightarrow +\infty} x^{\frac{5}{3}} \left(\frac{2}{3x} + o\left(\frac{1}{x}\right) \right)^2 = \lim_{x \rightarrow +\infty} x^{\frac{5}{3}} \left(\frac{2}{3x} \right)^2 \stackrel{(P.S.I.)}{=} \lim_{x \rightarrow +\infty} \frac{4}{9} x^{-\frac{1}{3}} = 0 \end{aligned}$$

Tempo a disposizione: 1 ore and 30 minuti.

Appello of the 14.09.2020 - Modalità telematica (causa COVID)

THEME 1

Exercise 1 [6 punti] Consider the function

$$f(x) = \arctan \left(\frac{x+1}{x-1} \right), \quad x \in (1, +\infty).$$

- (i) Individuarne the asymptotes.
- (ii) If ne determini the monotonicity .

Solution.

(i) the function is defined and continuous in the whole the domain $(1, +\infty)$, therefore the asymptotes riguardano solo $x \rightarrow 1+$ and $x \rightarrow +\infty$. Da

$$\lim_{x \rightarrow 1^+} \arctan \left(\frac{x+1}{x-1} \right) \underset{y=\frac{x+1}{x-1}}{=} \lim_{y \rightarrow +\infty} \arctan y = \frac{\pi}{2}$$

$$\lim_{x \rightarrow +\infty} \arctan \left(\frac{x+1}{x-1} \right) \underset{y=\frac{x+1}{x-1}}{=} \lim_{y \rightarrow 1} \arctan y = \frac{\pi}{4}$$

one gets the function ha a horizontal asymptote study $y \rightarrow +\infty$ of equation $y = \frac{\pi}{4}$.

- (ii) Calcoliamo the derivative of f :

$$\frac{df}{dx}(x) = \frac{1}{1 + \frac{(x+1)^2}{(x-1)^2}} \cdot \frac{-2}{(x-1)^2} = -\frac{2}{(x-1)^2 + (x+1)^2}.$$

Therefore $\frac{df}{dx}(x) < 0$ study ogni $x \in (1, +\infty)$, from which segue that the function is strictly decreasing in the domain $(1, +\infty)$.

Exercise 2 [6 punti] Consider the complex number $z = \sqrt{3} - i$.

- (i) Scrivere in esponenziale form .
- (ii) Compute the real part of z^6 .

Solution.

- (i) One has $\rho := |z| = \sqrt{3+1} = 2$, from which

$$z = 2 \left(\frac{\sqrt{3}}{2} - i \frac{1}{2} \right) = 2e^{-i\frac{\pi}{6}}$$

(ii)

$$\operatorname{Re}(z^6) = \operatorname{Re}(2^6 e^{-i\pi}) = -64 \quad (= z^6)$$

Exercise 3 [6 punti] Establish the convergence semplice and assoluta of the series

$$\sum_{n=1}^{\infty} (-1)^n \frac{n}{n^2 + 1}.$$

Solution. (i) In virtù of the criterio di Leibniz the series simply converges

:

- ha segni alterni;
- $\frac{n}{n^2+1}$ is decreasing infatti,

$$\frac{n_1}{n_1^2 + 1} \geq \frac{n_2}{n_2^2 + 1} \iff n_2(n_1^2 + 1) \leq n_1(n_2^2 + 1) \iff (n_2 - n_1)(1 - n_2 n_1) \leq 0 \iff n_2 \geq n_1,$$

(the ultimo passaggio dovuto al fatto che $(1 - n_2 n_1) \leq 0$); oppure si calcola the derivative

$$\left(\frac{x}{x^2 + 1} \right)' = \frac{-x^2 + 1}{(x^2 + 1)^2} \leq 0 \iff |x| \geq 1 \quad \text{if } x \geq 1$$

- one has $\lim_{n \rightarrow +\infty} (-1)^n \frac{n}{n^2 + 1} = 0$.

- (ii) The series does not converge assolutamente, perchànd the series

$$\sum_{n=1}^{\infty} \left| (-1)^n \frac{n}{n^2 + 1} \right| = \sum_{n=1}^{\infty} \frac{n}{n^2 + 1}$$

ànd asintotica alla series armonica.

Exercise 4 [6 punti] Compute the limit

$$\lim_{x \rightarrow 0^+} \frac{\log(1 + \sinh x) - \sin x}{x^2}.$$

Solution. da

$$\log(1 + \sinh x) - \sin x = \sinh x - \frac{(\sinh x)^2}{2} + o((\sinh x)^2) - \sin x = \frac{x^3}{3} + o(x^3) - \frac{(x)^2}{2} + o(x^2)$$

one has

$$\lim_{x \rightarrow 0^+} \frac{\log(1 + \sinh x) - \sin x}{x^2} = \lim_{x \rightarrow 0^+} \frac{-\frac{(x)^2}{2} + o(x^2)}{x^2} = -\frac{1}{2}.$$

Oppure si applica De l'Hôpital twice.

Exercise 5 [6 punti] Consider the generalized integral

$$\int_1^\infty \log\left(\frac{x^\alpha}{x^\alpha + 1}\right) dx.$$

- (i) Compute the integral for $\alpha = 2$.
- (ii) Establish study quali $\alpha \in [0, \infty)$ it converges.

Solution.

(i)

$$\begin{aligned} \lim_{k \rightarrow +\infty} \int_1^k \log\left(\frac{x^2}{x^2 + 1}\right) dx &= \lim_{k \rightarrow +\infty} \left(\left[x \log\left(\frac{x^2}{x^2 + 1}\right) \right]_1^k - \int_1^k \frac{2}{(1 + x^2)} dx \right) = \\ &\lim_{k \rightarrow +\infty} \left(k \log\left(\frac{k^2}{k^2 + 1}\right) - \log\left(\frac{1}{2}\right) - 2 \arctan k + 2 \arctan 1 \right) \\ &\lim_{k \rightarrow +\infty} \left(k \log\left(1 - \frac{1}{k^2 + 1}\right) - \log\left(\frac{1}{2}\right) - 2 \arctan k + 2 \arctan 1 \right) \\ &= \log 2 - \pi + \frac{\pi}{2} = \log 2 - \frac{\pi}{2} \end{aligned}$$

(ii)

$$\log\left(\frac{x^\alpha}{x^\alpha + 1}\right) = \log\left(1 - \frac{1}{x^\alpha + 1}\right) = -\frac{1}{x^\alpha + 1} + o\left(\frac{1}{x^\alpha + 1}\right)$$

study asymptotic comparison con $-\frac{1}{x^\alpha}$ converges if and only if $\alpha > 1$.

NB: con log si indica the logarithm in base e .

Tempo a disposizione: 1 ore and 30 minuti.

THEME 1

Exercise 1 [8 punti] Consider the function

$$f(x) = \arctan\left(\frac{x}{x^2 + x + 1}\right);$$

- (i) find the domain, study the sign, compute the limits at the extremes of the domain;
- (ii) calcolare the first derivative, study the monotonicity intervals and find the punti estremanti;
- (iii) draw the graph of f .

Solution. (i). Iniziamo dallo studio del dominio. Il denominatore $x^2 + x + 1 > 0$ è sempre positivo, in quanto il discriminante $\Delta = -3$ è negativo. Considerando anche che il dominio della funzione arctan è tutto \mathbb{R} , otteniamo $D = \mathbb{R}$.

Per studiare il segno di f : dato che $x^2 + x + 1 > 0$ per ogni $x \in \mathbb{R}$, abbiamo

$$f(x) = \arctan\left(\frac{x}{x^2 + x + 1}\right) \geq 0 \iff \frac{x}{x^2 + x + 1} \geq 0 \iff x \geq 0$$

e

$$f(x) = 0 \iff x = 0.$$

Per studiare i limiti ai punti estremi del dominio; abbiamo

$$\lim_{x \rightarrow +\infty} \arctan\left(\frac{x}{x^2 + x + 1}\right) = \arctan\left(\lim_{x \rightarrow +\infty} \left(\frac{x}{x^2 + x + 1}\right)\right) = 0$$

$$\lim_{x \rightarrow -\infty} \arctan\left(\frac{x}{x^2 + x + 1}\right) = \arctan\left(\lim_{x \rightarrow -\infty} \left(\frac{x}{x^2 + x + 1}\right)\right) = 0.$$

Hence $y = 0$ è l'assento orizzontale a $+\infty$ e a $-\infty$.

(ii). La derivata di f è

$$f'(x) = \frac{1}{1 + \left(\frac{x}{x^2 + x + 1}\right)^2} \cdot \frac{-2x^2 - x + x^2 + x + 1}{(x^2 + x + 1)^2} = \frac{1}{\left(1 + \left(\frac{x}{x^2 + x + 1}\right)^2\right)(x^2 + x + 1)^2} (-x^2 + 1).$$

Hence

$$f'(x) \geq 0 \iff 1 - x^2 \geq 0 \iff x \in [-1, 1]$$

e

$$f'(x) = 0 \iff 1 - x^2 = 0 \iff x \in \{-1, 1\}.$$

We deduce che f is crescente in the interval $[-1, 1]$, decreasing in $]-\infty, -1]$ and in $[1, +\infty[$, hence $x = -1$ is point of minimum globale while $x = 1$ is point of global maximum.

(iii). the graph of the function is sketched in the picture .

Exercise 2 [8 punti] Find in \mathbb{C} the solutions of the equation

$$z^4 + (-1 + i)z^2 - i = 0.$$

Suggerimento: sostituire $w = z^2$.

Solution. Con the sostituzione $w = z^2$ one gets

$$w^2 + (-1 + i)w - i = 0$$

whose solutions are

$$w_{1,2} = \frac{1 - i + \sqrt{2i}}{2} = \frac{1 - i \pm (1 + i)}{2} = \{1, -i\}.$$

Therefore, the solutions they are 4 and coincidono con the union of the solutions of $z^2 = 1$ and $z^2 = -i$, it is a dire

$$z_1 = 1, \quad z_2 = -1, \quad z_3 = -\frac{\sqrt{2}}{2} + i\frac{\sqrt{2}}{2}, \quad z_4 = \frac{\sqrt{2}}{2} - i\frac{\sqrt{2}}{2}.$$

the solutions **Exercise 3 [8 punti]**

(i) Compute

$$\lim_{n \rightarrow \infty} \frac{n^{2n}}{(n+1)^{2n}}.$$

(ii) Study the convergence of the series

$$\sum_{n=1}^{\infty} \frac{(2n)!}{n^{2n}}.$$

Solution. (i).

$$\lim_{n \rightarrow \infty} \frac{n^{2n}}{(n+1)^{2n}} = \lim_{n \rightarrow \infty} \frac{1}{(1+1/n)^{2n}} = \left(\lim_{n \rightarrow \infty} \frac{1}{(1+1/n)^n} \right)^2 = e^{-2}.$$

(ii). Since the series has positive terms, let us apply the criterio of the rapporto asymptotic and the result of (i), ottenendo

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{(2n+2)!n^{2n}}{(2n)!(n+1)^{2n+2}} &= \lim_{n \rightarrow \infty} \frac{(2n+2)(2n+1)n^{2n}}{(n+1)^2(n+1)^{2n}} = \lim_{n \rightarrow \infty} \frac{(4n^2+6n+2)n^{2n}}{(n^2+2n+1)(n+1)^{2n}} \\ &= 4 \lim_{n \rightarrow \infty} \frac{n^{2n}}{(n+1)^{2n}} = 4e^{-2}. \end{aligned}$$

Vale $4e^{-2} < 1$; the series is converging .

Exercise 4 [8 punti] study $\alpha \in \mathbb{R}$, si consideri

$$f_\alpha(x) = \frac{1}{\sinh x + x^\alpha}.$$

(a) Study as $\alpha \in \mathbb{R}$ the convergence

$$\int_0^{\log 2} f_\alpha(x) dx.$$

(b) Compute

$$\int_0^{\log 2} f_0(x) dx.$$

Solution. (a). Si tratta of an integrand a values positivi hence possiamo sfruttare the criterion of the asymptotic comparison . Da

$$f_\alpha(x) = \frac{1}{\sinh x + x^\alpha} = \frac{1}{x + o(x) + x^\alpha}$$

we get that, study $x \rightarrow 0$ the function is asintotica a $\frac{1}{x}$ if $\alpha > 1$, a $\frac{2}{x}$ if $\alpha = 1$ and a $\frac{1}{x^\alpha}$ if $\alpha < 1$. Therefore the integral converges $\Leftrightarrow \alpha < 1$.

(b) Con the sostituzione $t = e^x$ (that is, $x = \log t$) one gets

$$\begin{aligned} \int_0^{\log 2} f_0(x) dx &= \int_0^{\log 2} \frac{1}{\sinh x + 1} dx = \int_0^{\log 2} \frac{2}{e^x - e^{-x} + 2} dx = \int_1^2 \frac{2}{(t - t^{-1} + 2)t} dt \\ &= \int_1^2 \frac{2}{t^2 + 2t - 1} dt. \end{aligned}$$

Le radici of $t^2 + 2t - 1 = 0$ they are $-1 \pm \sqrt{2}$, hence

$$\frac{1}{t^2 + 2t - 1} = \frac{A}{t + 1 - \sqrt{2}} + \frac{B}{t + 1 + \sqrt{2}}$$

study suitable $A, B \in \mathbb{R}$. One has $1 = A(t + 1 + \sqrt{2}) + B(t + 1 - \sqrt{2})$, from which

$$\begin{cases} A + B = 0 \\ A(1 + \sqrt{2}) + B(1 - \sqrt{2}) = 1 \end{cases}$$

hence

$$\begin{cases} A + B = 0 \\ -B(1 + \sqrt{2}) + B(1 - \sqrt{2}) = 1 \end{cases}$$

and therefore $A = \frac{\sqrt{2}}{4}$, $B = -\frac{\sqrt{2}}{4}$. We deduce

$$\begin{aligned}\int_0^{\log 2} f_0(x) dx &= \left(\frac{\sqrt{2}}{2} \int_1^2 \frac{1}{t+1-\sqrt{2}} dt - \frac{\sqrt{2}}{2} \int_1^2 \frac{1}{t+1+\sqrt{2}} dt \right) \\ &= \left[\frac{\sqrt{2}}{2} \log |t+1-\sqrt{2}| - \frac{\sqrt{2}}{2} \log |t+1+\sqrt{2}| \right]_1^2 \\ &= \frac{\sqrt{2}}{2} \log \left(\frac{3-\sqrt{2}}{3+\sqrt{2}} \right) - \frac{\sqrt{2}}{2} \log \left(\frac{2-\sqrt{2}}{2+\sqrt{2}} \right).\end{aligned}$$

Tempo a disposizione: 1 ore and 30 minuti.

Appello of the 08.02.2021 - Modalità telematica (causa COVID)

THEME 1

Exercise 1 [8 punti] Consider the function

$$f(x) = \sqrt{\frac{|x|}{x^2 + 1}}.$$

- (i) Determine the domain of f , study the sign and the simmetria of f and compute limits and asymptotes at the extremes of the domain;
- (ii) Study the derivability of f and compute the first derivative, study the monotonicity intervals and find the points of maximum/ absolute minimum/relativo;
- (iii) draw the graph of f .

Solution. (i). Iniziamo dal domain. The denominator $x^2 + 1 > 0$ is always strictly positive . the numerator $|x|$ is always maggiorearctangent uguale dizerlo. Considerando that the domain of the function Root Test is $[0, \infty)$, we get $D = \mathbb{R}$.

Let us study the sign and le simmetries of f . The function is pari: $f(x) = f(-x), \forall x \in \mathbb{R}$. Furthermore, it ha always values non negativi:

$$f(x) = \sqrt{\frac{|x|}{x^2 + 1}} \geq 0 \iff x \in \mathbb{R},$$

e

$$f(x) = 0 \iff x = 0.$$

Let us study the limits at the extremes of the domain; abbiamo:

$$\lim_{x \rightarrow +\infty} f(x) = \lim_{x \rightarrow -\infty} f(x) = 0.$$

Hence $y = 0$ is horizontal asymptote a $+\infty$ and a $-\infty$.

(ii) Let us study the derivability of f . One has che $f \in C^{(1)}(\mathbb{R} \setminus \{0\})$ in quanto superposition difunctions $C^{(1)}(\mathbb{R})$, esclusa the function $g(x) = |x|$ that stonly one in $C^{(1)}(\mathbb{R} \setminus \{0\})$. study ogni $x \neq 0$ one has

$$f'(x) = \frac{1}{2\sqrt{\frac{|x|}{x^2+1}}} \frac{\operatorname{sgn}(x)(x^2+1) - |x|2x}{(x+1)^2}.$$

Hence,

$$\{x > 0 \text{ e } f'(x) > 0\} \iff (x^2+1) - 2x^2 > 0 \iff 1 - x^2 > 0 \iff x \in]0, 1[$$

e

$$\{x > 0 \text{ and } f'(x) = 0\} \iff x = 1$$

study simmetria, one has

$$\{x < 0 \text{ and } f'(x) > 0\} \iff x \in]-\infty, -1[$$

e

$$\{x < 0 \text{ and } f'(x) = 0\} \iff x = -1.$$

Furthermore,, study the teorema of the limit of the derivata,

$$\begin{aligned} \lim_{x \rightarrow 0^+} f'(x) &= \lim_{x \rightarrow 0^+} \frac{1 - x^2}{2(x^2 + 1)^2 \sqrt{\frac{|x|}{x^2+1}}} = \lim_{x \rightarrow 0^+} \frac{1}{2\sqrt{|x|}} = +\infty \\ \lim_{x \rightarrow 0^-} f'(x) &= \lim_{x \rightarrow 0^-} \frac{(-1)(1 - x^2)}{2(x^2 + 1)^2 \sqrt{\frac{|x|}{x^2+1}}} = \lim_{x \rightarrow 0^-} \frac{-1}{2\sqrt{|x|}} = -\infty \end{aligned}$$

therefore, the function is not differentiable in $x = 0$, where ha a cuspid.

Dalla precedente analisi and dalla continuity of the function one has that the function is increasing in each of the two intervals $[0, 1]$ and $] -\infty, -1[$ and is decreasing in each of the two intervals $[-1, 0]$ and $[1, +\infty[$.

Furthermore, vi is a maximum (resp. minimum) globale in $x = 1$ (resp. $x = -1$).

(iii). the graph of the function is sketched in the picture .

Exercice 2 [8 punti] Find the complex solutions of the equation

$$\frac{8}{z^3} = \frac{1+i}{1-i},$$

in algebraic and exponential form (or trigonometric), and draw them on the Gauss plane .

Solution. Da

$$\frac{1+i}{1-i} = \frac{1+i}{1-i} \cdot \frac{1+i}{1+i} = \frac{2i}{2} = i$$



grafici/2app2021T1E1.pdf

Figure 18: the graph of f .

we get

$$z^3 = \frac{8}{i} = -8i.$$

Solutions of equation they are le (three) radici terze of $-8i = 8e^{i\frac{3}{2}\pi}$, that is,

$$z_1 = 2e^{i\frac{1}{2}\pi} = 2i, \quad z_2 = 2e^{i\frac{7}{6}\pi} = -\sqrt{3} - i, \quad z_3 = 2e^{i\frac{11}{6}\pi} = \sqrt{3} - i$$

and they are disegnate nella figura seguente

Exercise 3 [8 punti]

(i) Compute

$$\int \log(t+1) dt.$$

(ii) Dedurre the value of

$$\int_0^1 \frac{\log(\sqrt{x} + 1)}{\sqrt{x}} dx.$$

grafici/2app2021T1E2.pdf

Figure 19: Solutions of exercise 2.

Solution. (i) study parti:

$$\begin{aligned}\int \log(t+1) dt &= \log(t+1)t - \int \frac{t}{t+1} dt \\ &= t \log(t+1) - \int \left(1 - \frac{1}{t+1}\right) dt \\ &= t \log(t+1) - t + \log|t+1| + c,\end{aligned}$$

con $c \in \mathbb{R}$.

(ii) Utilizzando the sostituzione $t = \sqrt{x}$,

$$\begin{aligned}\int_0^1 \frac{\log(\sqrt{x}+1)}{\sqrt{x}} dx &= \lim_{c \rightarrow 0+} \int_c^1 \frac{\log(\sqrt{x}+1)}{\sqrt{x}} dx = \lim_{c \rightarrow 0+} \int_{\sqrt{c}}^1 \frac{\log(t+1)}{t} 2t dt \\ &= \lim_{c \rightarrow 0+} 2 \int_{\sqrt{c}}^1 \log(t+1) dt \\ &= \lim_{c \rightarrow 0+} 2 [t \log(t+1) - t + \log(t+1)]_{\sqrt{c}}^1 \\ &= 2(2 \log 2 - 1)\end{aligned}$$

Exercise 4 [8 punti]

(i) Individuare as $\alpha \in \mathbb{R}$ the order di infinitesimal 1 of

$$n(\cos(1/n) - 1) + \frac{\alpha}{n}$$

(ii) Study as $\alpha \in \mathbb{R}$ the convergence of

$$\sum_{n=1}^{+\infty} \left| n(\cos(1/n) - 1) + \frac{\alpha}{n} \right|.$$

Solution. (i)

$$n(\cos(1/n) - 1) + \frac{\alpha}{n} = n \left(-\frac{1}{2n^2} + \frac{1}{24n^4} + o\left(\frac{1}{n^4}\right) \right) + \frac{\alpha}{n} = \frac{-1/2 + \alpha}{n} + \frac{1}{24n^3} + o\left(\frac{1}{n^3}\right)$$

and di order 1 study ogni $\alpha \neq 1/2$ and di order 3 study $\alpha = 1/2$.

(ii) The terms of this series have constant sign. Da quanto visto at the previous point, the general term of the series verifica le seguenti asintoticità

$$a_n \sim \begin{cases} \frac{-1/2 + \alpha}{n} & \text{if } \alpha \neq 1/2 \\ \frac{1}{24n^3} & \text{if } \alpha = 1/2. \end{cases}$$

Applicando the teorema of the asymptotic comparison con the series armonica generalizzata, we get that the series converges if $\alpha = 1/2$ and diverges if $\alpha \neq 1/2$.

Tempo a disposizione: 1 ore and 30 minuti.

Appello of the 05.07.2021 - Modalità telematica (causa COVID)

THEME 1

Exercise 1 [8 punti] Consider the function

$$f(x) = \log \left(1 + \sqrt{1 - x^2} \right).$$

- (i) Determine the domain of f , study the sign and the simmetria of f and compute the limits at the extremes of the domain;
- (ii) Study the derivability of f and compute the first derivative, study the monotonicity intervals and find the points of maximum/ absolute minimum/relativo;
- (iii) draw the graph of f .

Solution. (i). In order to determine the domain bisogna imporre that the radicando sia nonnegativo and the argument of the logarithm sia positive. The diseguaglianza $1 - x^2 \geq 0$ ha come solutions $x \in [-1, 1]$. study

Figure 20: graph of the exercise 1

questi valori di x , è ovvio che l'argomento del logaritmo sia positivo. Therefore

$$\text{dom}(f) = [-1, 1].$$

To single out the simmetries, let us observe that it is

$$f(-x) = \log\left(1 + \sqrt{1 - (-x)^2}\right) = f(x);$$

the function is pari.

Let us study the sign of the function: $f(x) \geq 0$ equivale a

$$1 + \sqrt{1 - x^2} \geq 1 \quad \text{that is,} \quad \sqrt{1 - x^2} \geq 0.$$

Since $\sqrt{\dots}$ is sicuramente nonnegativo, deduce that the function is always nonnegativa and si annulla only one in $x = \pm 1$ that they are therefore points of absolute minimum (con $f(\pm 1) = 0$).

From the theorem on the algebra of continuous functions and the theorem on superposition of discontinuous functions, $f \in C^0(\text{dom}(f))$. We henc

$$\lim_{x \rightarrow 1^-} f(x) = f(1) = 0$$

and analogamente, for simmetria, $\lim_{x \rightarrow -1^+} f(x) = 0$.

(ii). In $(-1, 1)$, for the teorema sull'algebra of the derivate and quello sulla derivata della funzione composta, we get that the f is differentiable. The derivability in ± 1 va studiata separatamente. Abbiamo

$$f'(x) = \frac{1}{1 + \sqrt{1 - x^2}} \cdot \frac{-x}{\sqrt{1 - x^2}}.$$

Since it is $\lim_{x \rightarrow 1^-} f'(x) = -\infty$ (and for simmetria $\lim_{x \rightarrow -1^+} f'(x) = +\infty$), concludiamo che f is not differentiable in $x = \pm 1$. Furthermore, the intervals di crescenza they are determinati da $f' \geq 0$ that is, $x \leq 0$. Therefore che

- f is increasing in $[-1, 0]$
- f is decreasing in $[0, 1]$
- $x = 0$ is the unique point di absolute maximum
- $x = \pm 1$ they are points of absolute minimum (già lo sapevamo).

Figure 21: graph of the exercise 2

(iii). Si veda the graph in the picture 1.

Exercise 2 [8 punti] Find the complex solutions of the equation

$$\operatorname{Im}(z^2) + |z|^2 \operatorname{Re}\left(\frac{1}{z}\right) = 0,$$

and draw them on the Gauss plane .

Solution. Innanzitutto notiamo that the equation ha senso solo for $z \neq 0$. Per tali values of z risolviamo the equation usando the algebraic form of the numeri complessi: $z = x + iy$ con $x, y \in \mathbb{R}$. Abbiamo

$$z^2 = (x^2 - y^2) + 2ixy, \quad |z|^2 = x^2 + y^2, \quad \frac{1}{z} = \frac{x - iy}{x^2 + y^2}.$$

The equation iniziale diventa

$$2xy + x = 0 \quad \text{that is,} \quad x(2y + 1) = 0$$

that ha solutions

$$x = 0 \quad \text{e} \quad y = -\frac{1}{2}$$

that formano le two rette (for $z \neq 0$) in the graph in Figura 2.

Exercise 3 [8 punti]

Sia

$$f_\alpha(x) := \frac{\arctan x}{1 + x^{2\alpha}}.$$

(i) Compute

$$\int f_1(x) dx = \int \arctan x \left(\frac{1}{1 + x^2} \right) dx.$$

(ii) Study as $\alpha \in [0, \infty)$ the convergence of

$$\int_1^{+\infty} f_\alpha(x) dx.$$

Solution. (i). By making use the sostituzione $\arctan x = t$ (ricordarsi: $(\arctan x)' = \frac{1}{1+x^2}$) we get

$$\int \arctan x \left(\frac{1}{1 + x^2} \right) dx = \int t dt = \frac{t^2}{2} + c = \frac{\arctan^2 x}{2} + c, \quad c \in \mathbb{R}.$$

(ii). Osserviamo $f \in C^0([1, +\infty))$ (e $f > 0$ su $[1, +\infty)$); hence the integral is improper solo for $x \rightarrow +\infty$. Let us study the asymptoticity of f_α for $x \rightarrow +\infty$:

$$f_\alpha(x) \sim \frac{\pi}{2} \cdot \frac{1}{1+x^{2\alpha}} \sim \frac{\pi}{2} \cdot \frac{1}{x^{2\alpha}} \quad \text{for } x \rightarrow +\infty.$$

Applicando the criterion of the asymptotic comparison for the integral the improper (and ricordando che $\int_1^{+\infty} x^a dx$ converges if and only if $a < -1$) we get that the integral dipartenza is converging if and only if $\alpha > 1/2$.

Exercise 4 [8 punti]

(i) Compute as $\alpha \in \mathbb{R}$ the limit

$$\lim_{n \rightarrow \infty} \frac{2 \log[\cos(1/n)] + \alpha[\sin(1/n)]^2}{(1/n)^2}.$$

(ii) Dedurre the comportamento of the series

$$\sum_{n=1}^{\infty} \{2 \log[\cos(1/n)] + [\sin(1/n)]^2\}.$$

Solution. (i). By making use the sviluppi di Mc Laurin of $\cos x$ and of $\log(1+x)$, for $n \rightarrow +\infty$ abbiamo

$$\begin{aligned} \log[\cos(1/n)] &= \log \left[1 + \left(-\frac{1}{2} \frac{1}{n^2} + o\left(\frac{1}{n^3}\right) \right) \right] \\ &= -\frac{1}{2} \frac{1}{n^2} + o\left(\frac{1}{n^3}\right) + o\left(-\frac{1}{2} \frac{1}{n^2} + o\left(\frac{1}{n^3}\right)\right) \\ &= -\frac{1}{2} \frac{1}{n^2} + o\left(\frac{1}{n^2}\right). \end{aligned}$$

Furthermore,, usando lo sviluppo di Mc Laurin of $\sin x$, for $n \rightarrow +\infty$ abbiamo

$$[\sin(1/n)]^2 = \left[\frac{1}{n} + o\left(\frac{1}{n^2}\right) \right]^2 = \frac{1}{n^2} + o\left(\frac{1}{n^3}\right).$$

Deduciamo that the numerator verifica

$$\text{num.} = (\alpha - 1) \frac{1}{n^2} + o\left(\frac{1}{n^2}\right);$$

conseguentemente vale

$$\lim_{n \rightarrow \infty} \frac{2 \log[\cos(1/n)] + \alpha[\sin(1/n)]^2}{(1/n)^2} = \alpha - 1 \quad \forall \alpha \in \mathbb{R}.$$

(ii). Let us observe that the point precedente con $\alpha = 1$ dà

$$\lim_{n \rightarrow +\infty} \frac{2 \log[\cos(1/n)] + [\sin(1/n)]^2}{(1/n)^2} = 0$$

that is,

$$2 \log[\cos(1/n)] + [\sin(1/n)]^2 = o[(1/n)^2] \quad \text{for } n \rightarrow +\infty.$$

We deduce in particolare that the termine of the nostra series is definitively positive . Furthermore,, applicando the criterion of the asymptotic comparison and ricordando che $\sum(1/n)^2$ is converging, we get that the series is converging .

Tempo a disposizione: 1 ore and 30 minuti.

Appello of the 13.09.2021 - Modalità telematica (causa COVID)

THEME 1

Exercise 1 [8 punti] Consider the function

$$f(x) = \frac{|\sin x|}{1 - 2 \cos x} .$$

- (i) Find the domain; study the periodicity , the sign and the simmetria of f ;
- (ii) study the derivability and calcolarne the first derivative ; study the monotonicity intervals and find the points of maximum/ absolute minimum/relativo;
- (iii) draw the graph.

Solution. (i). The function is defined for every $x \in \mathbb{R}$ tale che

$$1 - 2 \cos x \neq 0 \iff \cos x \neq \frac{1}{2} \iff x \in \mathbb{R} \setminus \left\{ \pm \frac{\pi}{3} + 2k\pi, \ k \in \mathbb{Z} \right\}$$

Clearly the function is periodica con periodo 2π . Furthermore,

$$f(x) = \frac{|\sin x|}{1 - 2 \cos x} = \frac{|\sin(-x)|}{1 - 2 \cos(-x)} = f(-x)$$

therefore the function is pari, that is, the suo graphis simmetrico rispetto all'asse of the ordinate.

Limits the study al domain $[-\pi, \pi] \setminus \{\pm \frac{\pi}{3}\}$; calcolo the limit at the extremes

$$\lim_{x \rightarrow \pi/3^-} f(x) = -\infty, \quad \lim_{x \rightarrow \pi/3^+} f(x) = +\infty.$$

(ii). Per ogni point of the domain tale che $|\sin x| \neq 0$, cioé $x \neq k\pi$, $k \in \mathbb{Z}$, one has

$$f'(x) = \frac{\cos x \frac{|\sin x|}{\sin x} (1 - 2 \cos x) - 2 \sin x |\sin x|}{(1 - 2 \cos x)^2} = \frac{|\sin x|}{(1 - 2 \cos x)^2} \left(\frac{\cos x}{\sin x} (1 - 2 \cos x) - 2 \sin x \right)$$

$$f'(x) \geq 0 \iff \frac{1}{\sin x} (\cos x - 2 \cos^2 x + 2 \sin^2 x) = \frac{1}{\sin x} (\cos x - 2) \geq 0$$

In $]0, \pi[\setminus \{\pi/3\}$ one has $\sin x > 0$ and $\cos x - 2 < 0$, hence $f'(x) < 0$, therefore le restrictions to the intervals $]0, \pi/3[,]\pi/3, \pi[$ they are strictly decreasing. By symmetry, le restrictions to the intervals $]-\pi/3, 0[,]-\pi, -\pi/3[$ they are strictly crescenti. e, the function ha a minimum locale in π , with value $f(\pi) = 0$ and hence in ogni point $\pi + 2k\pi$, $k \in \mathbb{Z}$. One has moreover

$$\lim_{x \rightarrow 0^-} f'(x) = 1 \quad \lim_{x \rightarrow 0^+} f'(x) = -1 \quad \lim_{x \rightarrow \pi^-} f'(x) = -\frac{1}{3} \quad \lim_{x \rightarrow -\pi^+} f'(x) = \frac{1}{3};$$

hence the funziona presenta punti angolosi in $k\pi$, $\forall k \in \mathbb{Z}$.

(iii). Vedi figura.

Figure 22: the graph of f .

Exercise 2 [8 punti] Find the solutions $z \in \mathbb{C}$ of the inequality

$$\left| \frac{z+1}{z} \right| \geq 1$$

and draw them on the Gauss plane .

Solution. Innanzitutto Let us observe that the campo diesistenza of the disuguaglianza is dato da $|z| \neq 0$ that is, da $z \neq 0$. Forniamo two metodi disoluzione.

1. Eleviamo al quadrato entrambi the membri:

$$\frac{|z+1|^2}{|z|^2} \geq 1 \iff (x+1)^2 + y^2 \geq x^2 + y^2 \iff 1 + 2x \geq 0 \iff x \geq -1/2$$

- 2.

$$\left| \frac{z+1}{z} \right| \geq 1 \iff \left| 1 + \frac{x-iy}{x^2+y^2} \right| \geq 1 \iff |x^2 + y^2 + x - iy| \geq x^2 + y^2$$

if and only if

$$x^4 + y^4 + x^2 + 2x^2y^2 + 2x^3 + 2xy^2 + y^2 \geq x^4 + y^4 + 2x^2y^2$$

if and only if

$$x^2 + 2x^3 + 2xy^2 + y^2 \geq 0 \iff (2x+1)(x^2+y^2) \geq 0 \iff 2x+1 \geq 0 \iff x \geq -1/2.$$

Solutions they are the numeri complessi $z = x + iy$ (con $x, y \in \mathbb{R}$) tali che: $z \neq 0$ and $x \geq -1/2$.

Exercise 3 [8 punti]

Study the convergence of the series

$$\sum_{n=1}^{\infty} n^{\alpha} \left(\frac{1}{n} - \sin \frac{1}{n} \right)$$

as $\alpha \in \mathbb{R}$.

Solution. By making use lo sviluppo di McLaurin of $\sin x$ one gets is asintotica alla series

$$\sum_{n=1}^{\infty} \frac{1}{6} n^{\alpha-3},$$

in particolare is a series a termini positivi. Possiamo therefore applicare the criterion of the asymptotic comparison and dedurre that it is converging if and only if $\alpha - 3 < -1$, that is, $\alpha < 2$, and is divergente per $\alpha \geq 2$.

Exercise 4 [8 punti]

Compute the integral

$$\int_{-1}^0 \frac{x}{x^2 + 2x + 2} dx.$$

Solution. Abbiamo

$$\begin{aligned} \int_{-1}^0 \frac{x}{x^2 + 2x + 2} dx &= \frac{1}{2} \int_{-1}^0 \frac{2x + 2 - 2}{x^2 + 2x + 2} dx \\ &= \frac{1}{2} \int_{-1}^0 \frac{2x + 2}{x^2 + 2x + 2} dx - \int_{-1}^0 \frac{1}{x^2 + 2x + 2} dx \\ &= \frac{1}{2} \log(x^2 + 2x + 2) \Big|_{-1}^0 - \int_{-1}^0 \frac{1}{(x+1)^2 + 1} dx \\ &= \frac{1}{2} \log(x^2 + 2x + 2) \Big|_{-1}^0 - \arctan(x+1) \Big|_{-1}^0 \\ &= \frac{1}{2} \log(2) - \frac{\pi}{4}. \end{aligned}$$

Appello of the 17.01.2022 - Modalità telematica (causa COVID)

THEME 1

Exercise 1 [10 punti] Given the function

$$f(x) = \arctan\left(\frac{|x+1|}{x^2+4}\right),$$

(i) find the domain:

$$\text{Domain} = \mathbb{R};$$

study the sign:

$$f(x) \geq 0 \iff \frac{|x+1|}{x^2+4} \geq 0 \quad \text{Hence } f(x) \geq 0 \ \forall x \in \mathbb{R}. \text{ Furthermore, } f(x) = 0 \iff x = -1;$$

compute the limits at the extremes of the domain:

$$\lim_{x \rightarrow \pm\infty} \arctan\left(\frac{|x+1|}{x^2+4}\right) \underset{y=\frac{|x+1|}{x^2+4}}{=} \lim_{y \rightarrow 0} \arctan y = 0,$$

hence $y = 0$ is horizontal asymptote at $-\infty$ and at $+\infty$.

(ii) Study the derivability of f sul suo domain, compute the first derivative :

Let us study f separatamente nelle regioni

$$x > -1 \iff |x+1| = x+1 \text{ and}$$

$$x < -1 \iff |x+1| = -(x+1), \text{ so that one has:}$$

$$f(x) = \arctan\left(\mp \frac{(x+1)}{x^2+4}\right) \quad x \leq -1,$$

and hence

$$f'(x) = \frac{\mp \frac{x^2+4-2x(x+1)}{(x^2+4)^2}}{1 + \frac{(x+1)^2}{(x^2+4)^2}} = \pm \frac{x^2+2x-4}{(x^2+4)^2 + (x+1)^2} \quad \text{if } x \leq -1,$$

$$\lim_{x \rightarrow -1^-} f'(x) = -\frac{1}{5} \quad \lim_{x \rightarrow -1^+} f'(x) = \frac{1}{5} \implies f'_-(-1) = -\frac{1}{5}, \quad f'_+(-1) = \frac{1}{5}.$$

Hence the function is differentiable for every $x \in \mathbb{R} \setminus \{-1\}$ while in $x = -1$ vi is a angular point .

Study the intervals dimonotonicity :

$$\begin{cases} f'(x) \geq 0 \\ x < -1 \end{cases} \iff \begin{cases} x^2+2x-4 \geq 0 \\ x < -1 \end{cases} \iff x \leq -1 - \sqrt{5}$$

Hence the function is strictly increasing in $]-\infty, -1 - \sqrt{5}]$ and strictly decreasing in $[-1 - \sqrt{5}, -1[$.

Furthermore,

$$\begin{cases} f'(x) \geq 0 \\ x > -1 \end{cases} \iff \begin{cases} x^2+2x-4 \leq 0 \\ x > -1 \end{cases} \iff -1 < x \leq -1 + \sqrt{5}.$$

Hence the function is strictly increasing in $] -1, -1 + \sqrt{5} [$ and strictly decreasing in $] -1 + \sqrt{5}, +\infty [$. Finally ,

$$f'(-1 + \sqrt{5}) = f'(-1 - \sqrt{5}) = 0$$

and the punti $-1 - \sqrt{5}, -1 + \sqrt{5}$ they are direlative maximum . Da $f(-1) = 0$, the point $x = -1$ is point of absolute minimum.

(iii) draw the graph of f .



Figure 23: Grafico of the function.

Exercise 2 [7 punti] Determine the solutions in \mathbb{C} of the equation

$$\left(\frac{z}{i}\right)^3 = -8 \iff z^3 = 8i = 8 \left(\cos\left(\frac{1}{2}\pi\right) + i \sin\left(\frac{1}{2}\pi\right) \right)$$

Dobbiamo that is, trovare le radici terze of $8i$, that is,, con the formula di De Moivre,

$$z_0 = 2 \left(\cos\left(\frac{1}{6}\pi\right) + i \sin\left(\frac{1}{6}\pi\right) \right) = \sqrt{3} + i$$

$$z_1 = 2 \left(\cos\left(\frac{5}{6}\pi\right) + i \sin\left(\frac{5}{6}\pi\right) \right) = -\sqrt{3} + i$$

$$z_2 = 2 \left(\cos\left(\frac{3}{2}\pi\right) + i \sin\left(\frac{3}{2}\pi\right) \right) = -2i.$$

(Stanno sui verteces of a triangolo equilatero inscritto in the cerchio diraggio
2 con a vertex in $= -2i$)

Exercise 3 [7 punti]

(i) Mediante suitable sviluppi di Taylor, determine, as $\alpha \in \mathbb{R}$, a sviluppo of the sequenze

$$a_n = \frac{1}{n} - \sin\left(\frac{1}{n}\right) - \alpha \log\left(1 + \frac{1}{n^3}\right) \quad \text{for } n \rightarrow +\infty.$$

One has

$$\begin{aligned} a_n &= \frac{1}{n} - \sin\left(\frac{1}{n}\right) - \alpha \log\left(1 + \frac{1}{n^3}\right) = \frac{1}{n} - \frac{1}{n} + \frac{1}{6n^3} - \frac{1}{120n^5} + o\left(\frac{1}{n^5}\right) - \alpha \frac{1}{n^3} + \alpha \frac{1}{2n^6} + o\left(\frac{\alpha}{n^6}\right) = \\ &= \frac{1 - 6\alpha}{6n^3} - \frac{1}{120n^5} + o\left(\frac{1}{n^5}\right) \end{aligned}$$

(ii) Study the convergence of the series

$$\sum_{n=1}^{\infty} n^2 a_n.$$

Da

$$\sum_{n=1}^{\infty} n^2 a_n = \sum_{n=1}^{\infty} \left(\frac{1 - 6\alpha}{6n} - \frac{1}{120n^3} + o\left(\frac{1}{n^3}\right) \right)$$

segue che, for $\alpha = \frac{1}{6}$, the termine generico of the series is always negative for n sufficientemente grande and is asymptotic to $\frac{1}{n^3}$. Therefore the series converges. If instead $\alpha \neq \frac{1}{6}$, the termine generico of the series is with constant sign for n sufficientemente grande and is asymptotic to $\frac{1}{n}$. Therefore the series diverges for $\alpha \neq \frac{1}{6}$.

Exercise 4 [8 punti]

(i) By making use the definizione, compute the integral generalizzato

$$\int_0^{+\infty} \frac{\arctan t}{(1+t^2)(\arctan^2 t + 8 \arctan t + 17)} dt;$$

Consideriamo the sostituzione $y = \arctan t$, that implica $dy = \frac{1}{1+t^2} dt$:

$$\begin{aligned} \lim_{c \rightarrow \infty} \int_0^c \frac{\arctan t}{(1+t^2)(\arctan^2 t + 8 \arctan t + 17)} dt &= \lim_{c \rightarrow +\infty} \int_0^{\arctan c} \frac{y}{y^2 + 8y + 17} dy = \\ &= \lim_{c \rightarrow +\infty} \left[\frac{1}{2} \log((y+4)^2 + 1) - 4 \arctan(y+4) \right]_0^{\arctan c} \\ &= \frac{1}{2} \log((\pi/2 + 4)^2 + 1) - 4 \arctan(\pi/2 + 1) - \frac{1}{2} \log 17 + 4 \arctan(4) \end{aligned}$$

where we have usato:

$$\begin{aligned} \int \frac{y}{y^2 + 8y + 17} dy &= \int \frac{y}{y^2 + 8y + 16 + 1} dy = \frac{1}{2} \int \frac{2(y+4)}{(y+4)^2 + 1} dy - 4 \int \frac{1}{(y+4)^2 + 1} dy = \\ &= \frac{1}{2} \log((y+4)^2 + 1) - 4 \arctan(y+4) + c, \quad c \in \mathbb{R} \end{aligned}$$

discuss the convergence of the generalized integral

$$\int_0^{+\infty} \frac{\arctan t}{(1+t^2)^{2\alpha} (\arctan^2 t + 8 \arctan t + 17)} dt;$$

for every $\alpha \in \mathbb{R}$.

The integrand, for $t \rightarrow +\infty$, is asymptotic to

$$\frac{C}{t^{4\alpha}}$$

for a suitable constant $C > 0$, hence the integral converges for

$$4\alpha > 1 \iff \alpha > \frac{1}{4}.$$

Appello of the 07.02.2022

Exercise 1 [10 punti] Given the function

$$f(x) = \log(|x| - x^2 + 2),$$

(i) determine the domain; determine the simmetria and the sign.

$$x \in \text{Domain} \iff |x| - x^2 + 2 > 0 \iff |x|^2 - |x| - 2 < 0 \quad (\text{da } x^2 = |x|^2)$$

inequality that is risolta da

$$|x| \in]-1, 2[\iff |x| \in [0, 2[\iff x \in]-2, 2[$$

Hence Domain = $] -2, 2 [$

The function is chiaramente pari.

The function is continuous perchand composta di continuopus functions .

In alternativa si sarebbe potuto also argomentare come segue.

$$f(x) = \begin{cases} \log(x - x^2 + 2) & \forall x \geq 0 \\ \log(-x - x^2 + 2) & \forall x < 0 \end{cases}$$

Si osserva che f è pari, and si limita lo studio a $x \geq 0$. Hence $x \in \text{Domain}$ and $x \geq 0$ if and only if $x - x^2 + 2 > 0$ and $x \geq 0$, that is, $x \in [0, 2[$. Since f è pari, one has

$$\text{Domain} =]-2, 2[$$

Furthermore,

$$\begin{cases} f(x) \geq 0 \\ x \geq 0 \end{cases} \iff \begin{cases} x - x^2 + 2 \geq 1 \\ x \geq 0 \end{cases} \iff x \in \left[0, \frac{1+\sqrt{5}}{2}\right].$$

By symmetry, si conclude che

$$f(x) \geq 0 \iff x \in \left[-\frac{1+\sqrt{5}}{2}, \frac{1+\sqrt{5}}{2}\right]$$

Compute the limits and asymptotes at the extremes of the domain:

$$\lim_{x \rightarrow 2} f(x) = -\infty \quad \text{that for simmetria implica} \quad \lim_{x \rightarrow -2} f(x) = -\infty$$

così in 2 and -2 ci sono due asintoti verticali.

(ii) study the derivability and calcolare the first derivative ; study the monotonicity intervals individuando the points of maximum and of minimum sia relativi che assoluti :

$$\begin{cases} x > 0 \\ f'(x) = \frac{1-2x}{x-x^2+2} > 0 \end{cases} \iff x \in \left]0, \frac{1}{2}\right[.$$

Furthermore, $f'(x) = 0$, $x > 0$ if and only if $x = \frac{1}{2}$. Since f è continua in its domain, if we deduce che f è strictly increasing in $[0, \frac{1}{2}]$, strictly decreasing in $[\frac{1}{2}, 2[$, and ha un punto di relativo massimo in $x = \frac{1}{2}$.

By symmetry, one has also che f è strictly decreasing in $[-\frac{1}{2}, 0]$, strictly increasing in $] -2, -\frac{1}{2} [$, and ha un punto di relativo massimo in $x = -\frac{1}{2}$.

In particolare $x = \frac{1}{2}, -\frac{1}{2}$ sono punti di massimo assoluto.

Per $x = 0$ (the function is continuous): $f(0) = \log 2$. $x = 0$ è hence punto di relativo minimo (ma non assoluto perché f tende a $-\infty$ at the extremes). One has moreover $\lim_{x \rightarrow 0^+} f'(x) = 1/2 = f'_+(0)$, that for simmetria implica $f'_-(0) = -1/2$. Hence 0 is a point of inflection.

(iii) draw the graph.

Exercise 2 [7 punti] Determine the insieme A of the numeri complessi $z \in \mathbb{C}$ tali che

$$\frac{|z + i\operatorname{Im}(z)|^2}{|z|^2 + \operatorname{Re}(z)^2} \geq 1$$

and disegnarlo in the Gauss plane .

If scriviamo $z = x + iy$, the inequality diventa

$$\frac{|x + 2yi|^2}{2x^2 + y^2} = \frac{x^2 + 4y^2}{2x^2 + y^2} \geq 1$$

the numerator is 0 if and solo if $(x, y) = (0, 0)$. Negli altri punti is positive, therefore for $(x, y) \neq (0, 0)$ the inequality is equivalente a

$$\begin{aligned} x^2 + 4y^2 &\geq 2x^2 + y^2 \iff \\ 3y^2 - x^2 &= (\sqrt{3}y - x)(\sqrt{3}y + x) \geq 0 \iff \\ x+iy \in \left\{ x+iy, \ y \leq x/\sqrt{3}, \ y \leq -x/\sqrt{3} \right\} \cup \left\{ x+iy, \ y \geq x/\sqrt{3}, \ y \geq -x/\sqrt{3} \right\}, \quad (x, y) \neq (0, 0). \end{aligned}$$

Exercise 3 [7 punti]

Study the convergence of the series

$$\sum_{n=1}^{\infty} n \left\{ \alpha \sinh \left(\frac{1}{n^2} \right) + \log \left[\cosh \left(\frac{1}{n} \right) \right] \right\}$$

as $\alpha \in \mathbb{R}$.

Da

$$\begin{aligned} n \left\{ \alpha \sinh \left(\frac{1}{n^2} \right) + \log \left[\cosh \left(\frac{1}{n} \right) \right] \right\} &= \\ = n \left\{ \alpha \frac{1}{n^2} + \alpha \cdot o \left(\frac{1}{n^4} \right) + \log \left[1 + \frac{1}{2n^2} + \frac{1}{24n^4} + o \left(\frac{1}{n^5} \right) \right] \right\} &= \\ = n \left\{ \alpha \frac{1}{n^2} + \alpha \cdot o \left(\frac{1}{n^4} \right) + \frac{1}{2n^2} - \frac{1}{12n^4} + o \left(\frac{1}{n^4} \right) \right\} &= \\ = (2\alpha + 1) \frac{1}{2n} - \frac{1}{12n^3} + o \left(\frac{1}{n^3} \right) & \end{aligned}$$

deduce that it is a series a sign definitively constant e, applicando the criterion of the asymptotic comparison, che converges if and only if $2\alpha + 1 = 0$, i.e. $\alpha = -1/2$.

Exercise 4 [8 punti]

By making use the integration by parts, compute

$$\int \arctan \left(\frac{2}{x} \right) dx.$$

Abbiamo

$$\begin{aligned}
\int \arctan\left(\frac{2}{x}\right) dx &= \int \arctan\left(\frac{2}{x}\right) dx \\
&= x \arctan\left(\frac{2}{x}\right) + \int \frac{2}{x(1+4/x^2)} dx \\
&= x \arctan\left(\frac{2}{x}\right) + \int \frac{2x}{x^2+4} dx \\
&= x \arctan\left(\frac{2}{x}\right) + \log(x^2+4) + c, \quad c \in \mathbb{R}.
\end{aligned}$$

. Study the convergence of the integral improprio

$$\int_0^{+\infty} \arctan\left(\frac{x^3+1}{x^\alpha}\right) dx$$

as $\alpha > 0$.

Let us observe that the function integrand is always $C^{(0)}((0, +\infty))$ and non-negativa. All'extreme $x = 0$ the integrand tende a $\pi/2$ hence

$$\int_0^1 \arctan\left(\frac{x^3+1}{x^\alpha}\right) dx$$

è and integrable for ogni $\alpha > 0$. Let us study the integrabilità of

$$\int_1^{+\infty} \arctan\left(\frac{x^3+1}{x^\alpha}\right) dx$$

If $\alpha \leq 3$ the argument of the arctangent is always > 1 so that $\arctan\left(\frac{x^3+1}{x^\alpha}\right) > \pi/4$. It follows that the integral diverges. If $\alpha > 3$, the argument of arctangent tende a zero for $x \rightarrow +\infty$, and the integrand is asymptotic to $1/x^{\alpha-3}$. Hence the ultimo integral converges if and only if $\alpha - 3 > 1$, that is, $\alpha > 4$.

In conclusion

$$\int_0^{+\infty} \arctan\left(\frac{x^3+1}{x^\alpha}\right) dx$$

converges if and only if $\alpha > 4$.

Appello of the 01.07.2022

Exercise 1 [9 punti] Consider the function

$$f(x) = |x-2| e^{\frac{1}{(x-2)^2}}.$$

(i) determine the domain of f and the sign of f ;

$$Domain = \mathbb{R} \setminus \{2\}$$

e

$$f(x) > 0$$

for every $x \in Domain$, perché prodotto di due funzioni positive.

(ii) compute the main limits of f ;

$$\lim_{x \rightarrow \pm\infty} |x - 2| e^{\frac{1}{(x-2)^2}} = +\infty \cdot 1 = +\infty$$

$$\lim_{x \rightarrow 2^\pm} |x - 2| e^{\frac{1}{(x-2)^2}} = \lim_{y \substack{\rightarrow \\ = \frac{1}{(x-2)^2}}} y^{-\frac{1}{2}} e^y = +\infty$$

(iii) compute the derivative of f , discuss the monotonicity of f and determine the infimum and the supremum of f and relative and absolute minimum and maximum points; Per ogni $x > 2$

$$\frac{df}{dx}(x) = e^{\frac{1}{(x-2)^2}} - 2(x-2)e^{\frac{1}{(x-2)^2}} \frac{1}{(x-2)^3} = e^{\frac{1}{(x-2)^2}} \left(1 - \frac{2}{(x-2)^2}\right) = e^{\frac{1}{(x-2)^2}} \cdot \frac{x^2 - 4x + 2}{(x-2)^2}$$

Analogamente, Per ogni $x < 2$

$$\frac{df}{dx}(x) = -\left(e^{\frac{1}{(x-2)^2}} - 2(x-2)e^{\frac{1}{(x-2)^2}} \frac{1}{(x-2)^3}\right) = -e^{\frac{1}{(x-2)^2}} \left(1 - \frac{2}{(x-2)^2}\right) = -e^{\frac{1}{(x-2)^2}} \cdot \frac{x^2 - 4x + 2}{(x-2)^2}$$

Hence, poiché $x^2 - 4x + 2 > 0$ if and only if $x > 2 + \sqrt{2}$ o $x < 2 - \sqrt{2}$,

one has che $\frac{df}{dx}(x) > 0$ if and only if $x > 2 + \sqrt{2}$ o $2 - \sqrt{2} < x < 2$, while

$$\frac{df}{dx}(2 + \sqrt{2}) = \frac{df}{dx}(2 - \sqrt{2}) = 0.$$

Furthermore, $f(2 + \sqrt{2}) = f(2 - \sqrt{2}) = \sqrt{2}$.

Hence the function is strictly increasing in $[2 - \sqrt{2}, 2[$ and in $]2 + \sqrt{2}, +\infty[$, is strictly decreasing in $] -\infty, 2 - \sqrt{2}[$ and in $]2, 2 + \sqrt{2}]$, cosicché in $2 + \sqrt{2}$ and in $2 - \sqrt{2}$ it ha two minimi relative che sono assoluti. Furthermore, la funzione è limitata superiormente.

(iv) compute asymptotes of f ;

$$\lim_{x \rightarrow +\infty} \frac{f(x)}{x} = \lim_{x \rightarrow +\infty} \frac{(x-2)e^{\frac{1}{(x-2)^2}}}{x} = 1 \cdot 1 = 1$$

$$\lim_{x \rightarrow -\infty} \frac{f(x)}{x} = \lim_{x \rightarrow -\infty} \frac{(2-x)e^{\frac{1}{(x-2)^2}}}{x} = -1 \cdot 1 = -1$$

$$\lim_{x \rightarrow +\infty} (f(x) - x) = (x-2)e^{\frac{1}{(x-2)^2}} - x =$$

Utilizzo lo sviluppo of e^y for $y \rightarrow 0$, con $y = \frac{1}{(x-2)^2}$

$$= \lim_{x \rightarrow +\infty} \left(x - 2 + \frac{x-2}{(x-2)^2} + o\left(\frac{1}{(x-2)}\right) - x \right) = -2$$

Analogamente

$$\lim_{x \rightarrow -\infty} (f(x)+x) = (2-x)e^{\frac{1}{(x-2)^2}} - x = \lim_{x \rightarrow +\infty} \left(2 - x + \frac{2-x}{(x-2)^2} + o\left(\frac{1}{(x-2)}\right) + x \right) = 2$$

In conclusion, for $x \rightarrow +\infty$, one has l' asintoto $y = -2+x$ and for $x \rightarrow -\infty$, one has l' asintoto $y = 2-x$

(v) draw a qualitative graph of f .

Exercise 2 [8 punti] Determine in algebraic form the solutions in \mathbb{C} of the equation

$$z^4 + (-2 - 2i)z^2 + 4i = 0.$$

Pongo $w := z^2$. The equation for w is

$$w^2 + (-2 - 2i)w + 4i = 0$$

whose solutions are

$$w_1 = 1 + i + r_1, \quad w_2 = 1 + i + r_2$$

dove r_1, r_2 they are le radici quadrate of $(1+i)^2 - 4i = 1 - 2i - 1 = -2i$. That is, $r_1 = -1 + i$ and $r_2 = 1 - i$, from which $w_1 = 2i, w_2 = 2$ so that the solutions si trovano unendo the solutions $z^2 = 2i$ a quelle of $z^2 = 2$. Ne segue (con the solito de Moivre) che the solutions they are

$$z_1 = 1 + i, z_2 = -1 - i, z_3 = \sqrt{2}, \quad z_4 = -\sqrt{2}$$

Exercise 3 [7 punti]

(i) Determine, as $\alpha \in \mathbb{R}$, the limit

$$\lim_{x \rightarrow 0^+} \frac{(1+x)^{\alpha x} - 1}{x^2}.$$

Utilizzando the principio of sostituzione in the prodotto/quoziente of limits con functions asintotiche, osservando che, for $x \rightarrow 0^+$, $e^{\alpha x \log(1+x)} - 1 \sim \alpha x \log(1+x) \sim \alpha x^2$, one gets

$$\lim_{x \rightarrow 0^+} \frac{(1+x)^{\alpha x} - 1}{x^2} = \lim_{x \rightarrow 0^+} \frac{e^{\log(1+x)^{\alpha x}} - 1}{x^2} = \lim_{x \rightarrow 0^+} \frac{\alpha x \log(1+x)}{x} = \lim_{x \rightarrow 0^+} \frac{\alpha x^2}{x} = \alpha.$$

Exercice 4 [8 punti] (i) Compute the seguente indefinite integral

$$\int \frac{\sqrt{t}}{1+t} dt.$$

Pongo $y := \sqrt{t}$, so that $dy = \frac{1}{2}(\sqrt{t})^{-1}dt = \frac{1}{2}y^{-1}dt$, that is, $dt = 2ydy$. One has hence

$$\begin{aligned} \int \frac{\sqrt{t}}{1+t} dt &= 2 \int \frac{y^2}{1+y^2} dy = 2 \left[\int \frac{1+y^2}{1+y^2} dy - \int \frac{1}{1+y^2} dy \right] = \\ &2(y - \arctan(y)) + c = 2\left(\sqrt{t} - \arctan(\sqrt{t})\right) + c, \quad \forall c \in \mathbb{R}. \end{aligned}$$

(ii) Discutere the convergence of the generalized integral

$$\int_0^{+\infty} \frac{\sqrt{t}}{1+t^\alpha} dt$$

as $\alpha \in \mathbb{R}$.

Per $t \rightarrow 0+$, if $\alpha \geq 0$ the function integrand is continuous nello 0. If instead $\alpha < 0$ the function is prolungabile for continuità, uguale a 0 nello 0. Hence non ci they are problemi diintegrabilità in a right neighbourhood of 0.

Per $t \rightarrow +\infty$, if $\alpha < 0$, $\frac{\sqrt{t}}{1+t^\alpha} \sim \sqrt{t}$ that is not integrable for $t \rightarrow +\infty$. If $\alpha = 0$ $\frac{\sqrt{t}}{1+t^\alpha} \sim \frac{\sqrt{t}}{2}$ che, similmente, is not integrable for $t \rightarrow +\infty$. If $\alpha > 0$, $\frac{\sqrt{t}}{1+t^\alpha} \sim \frac{1}{t^{\alpha-\frac{1}{2}}}$, that for $t \rightarrow +\infty$ is integrabile if and only if $\alpha - \frac{1}{2} > 1$ cioè, if and only if $\alpha > \frac{3}{2}$.