


2<sup>nd</sup> RECURSION THEOREM

Let  $f: \mathbb{N} \rightarrow \mathbb{N}$  computable total extensional



$$\forall e, e' \in \mathbb{N} \quad \varphi_e = \varphi_{e'}$$

$$\leadsto \varphi_{f(e)} = \varphi_{f(e')}$$

by Myhill-Shepherdson's theorem there a (unique)

recursive functional  $\bar{\Phi}: \mathcal{F}(\mathbb{N}) \rightarrow \mathcal{F}(\mathbb{N})$

$$\forall e \in \mathbb{N} \quad \bar{\Phi}(\varphi_e) = \varphi_{f(e)}$$

By 1<sup>st</sup> recursion theorem  $\bar{\Phi}$  has a least fixed point  $f_{\bar{\Phi}}: \mathbb{N} \rightarrow \mathbb{N}$   
computable

$$\begin{cases} \bar{\Phi}(f_{\bar{\Phi}}) = f_{\bar{\Phi}} \\ \exists e_0 \in \mathbb{N} \text{ s.t. } f_{\bar{\Phi}} = \varphi_{e_0} \end{cases}$$

$$\underline{\varphi_{e_0}} = f_{\bar{\Phi}} = \bar{\Phi}(f_{\bar{\Phi}}) = \bar{\Phi}(\varphi_{e_0}) = \underline{\varphi_{f(e_0)}}$$

In summary

Given  $f: \mathbb{N} \rightarrow \mathbb{N}$  computable total ~~extensional~~

there is  $e_0 \in \mathbb{N}$  s.t.  $\varphi_{e_0} = \varphi_{f(e_0)}$

2<sup>nd</sup> recursion theorem

## 2<sup>nd</sup> RECURSION THEOREM

Let  $f: \mathbb{N} \rightarrow \mathbb{N}$  be total computable function

Then there exists  $e_0 \in \mathbb{N}$  s.t.  $\varphi_{e_0} = \varphi_{f(e_0)}$

proof

Let  $f: \mathbb{N} \rightarrow \mathbb{N}$  be total computable

observe  $x \mapsto \varphi_x(x)$  computable

"  
 $\psi_v(x, x)$

$x \mapsto f(\varphi_x(x))$  computable

define

$$g(x, y) = \varphi_{f(\varphi_x(x))}(y) \quad \text{convention } \varphi_{\uparrow} = \uparrow$$

$$= \psi_v(f(\varphi_x(x)), y)$$

$$= \psi_v(f(\psi_v(x, x)), y) \quad \text{computable}$$

By smm theorem there  $s: \mathbb{N} \rightarrow \mathbb{N}$  total computable s.t.  $\forall x, y$

$$\varphi_{s(x)}(y) = g(x, y) = \varphi_{f(\varphi_x(x))}(y) \quad (*)$$

Since  $s$  is computable there is  $m \in \mathbb{N}$  s.t.  $s = \varphi_m$ .

Substituting in  $(*)$

$$\varphi_{\varphi_m(x)}(y) = \varphi_{f(\varphi_x(x))}(y) \quad \forall x, y$$

In particular, for  $x = m$

$$\varphi_{\varphi_m(m)}(y) = \varphi_{f(\varphi_m(m))}(y) \quad \forall y$$

Hence

$$\varphi_{\varphi_m(m)} = \varphi_{f(\varphi_m(m))}$$

If we let  $e_0 = \varphi_m(m)$  we conclude

$$\varphi_{e_0} = \varphi_{f(e_0)}$$

(note that  $\varphi_m = \Sigma$  total, hence  $\varphi_m(m) \downarrow$ )

□

Idea :

if  $h: \mathbb{N} \rightarrow \mathbb{N}$  computable

$$h \begin{matrix} \varphi_0 & \varphi_1 & \varphi_2 & \varphi_3 & \dots \\ \varphi_{h(0)} & \varphi_{h(1)} & \varphi_{h(2)} & \varphi_{h(3)} & \dots \end{matrix}$$

you can do the above for  $h = \varphi_i \quad i = 0, 1, 2, \dots$

$$\begin{array}{l} E_0 \quad \varphi_{\varphi_0(0)} \quad \varphi_{\varphi_0(1)} \quad \varphi_{\varphi_0(2)} \quad \dots \\ E_1 \quad \varphi_{\varphi_1(0)} \quad \varphi_{\varphi_1(1)} \quad \varphi_{\varphi_1(2)} \quad \dots \\ E_2 \quad \varphi_{\varphi_2(0)} \quad \varphi_{\varphi_2(1)} \quad \varphi_{\varphi_2(2)} \quad \dots \end{array}$$

$\leftarrow h(x) = \varphi_x(x)$

in the proof we took the diagonal transformed by  $f$

$$h(x) = f(\varphi_x(x)) = f(\varphi_{\varphi_m(x)}(x)) = \varphi_m(x)$$

$$\begin{array}{l} E_0 \quad \varphi_{\varphi_0(0)} \quad \varphi_{\varphi_0(1)} \quad \varphi_{\varphi_0(2)} \quad \dots \\ E_1 \quad \varphi_{\varphi_1(0)} \quad \varphi_{\varphi_1(1)} \quad \varphi_{\varphi_1(2)} \quad \dots \\ E_2 \quad \varphi_{\varphi_2(0)} \quad \varphi_{\varphi_2(1)} \quad \varphi_{\varphi_2(2)} \quad \dots \\ E_m \quad \varphi_{f(\varphi_0(0))} \quad \varphi_{f(\varphi_1(1))} \quad \varphi_{f(\varphi_2(2))} \quad \dots \end{array}$$

$\varphi_x(x)$

$\varphi_{\varphi_m(m)}$


$\varphi_{f(\varphi_m(m))}$

## Rice's Theorem

Let  $A \subseteq \mathbb{N}$  saturated  $A \neq \emptyset$   $A \neq \mathbb{N}$  then  $A$  not recursive

proof (alternative proof using 2<sup>nd</sup> recursion theorem)

Let  $A \subseteq \mathbb{N}$   $A \neq \emptyset$ ,  $A \neq \mathbb{N}$  saturated



$$A \neq \emptyset \rightsquigarrow \exists e_1 \in A$$

$$A \neq \mathbb{N} \rightsquigarrow \exists e_0 \in \bar{A}$$

Assume by contradiction  $A$  recursive and defined:

$$f: \mathbb{N} \rightarrow \mathbb{N}$$

$$f(x) = \begin{cases} e_0 & \text{if } x \in A \\ e_1 & \text{if } x \notin A \end{cases}$$

$$= e_0 \cdot \chi_A(x) + e_1 \cdot \chi_{\bar{A}}(x)$$

$$\left( \begin{array}{lll} \text{if } x \in A & \rightsquigarrow & \chi_A(x) = 1 \quad \chi_{\bar{A}}(x) = 0 \quad e_0 \cdot 1 + e_1 \cdot 0 = e_0 \\ \text{if } x \notin A & \rightsquigarrow & \chi_A(x) = 0 \quad \chi_{\bar{A}}(x) = 1 \quad e_0 \cdot 0 + e_1 \cdot 1 = e_1 \end{array} \right)$$

if  $A$  recursive,  $f$  computable total

but for all  $e \in \mathbb{N}$   $\varphi_e \neq \varphi_{f(e)}$

•  $e \in A \Rightarrow f(e) = e_0 \notin A$  and since  $A$  saturated

$$\varphi_e \neq \varphi_{f(e)}$$

•  $e \notin A \Rightarrow f(e) = e_1 \in A$  thus, since  $A$  saturated


$$\varphi_e \neq \varphi_{f(e)}$$

This contradicts the 2<sup>nd</sup> recursion theorem.  $\rightsquigarrow A$  not recursive

□

Proposition : The halting set  $K = \{ x \in \mathbb{N} \mid \varphi_x(x) \downarrow \}$  is not recursive.

proof (alternative proof using 2<sup>nd</sup> recursion theorem)



if  $e_0 \in \mathbb{N}$  s.t.  $\varphi_{e_0}(x) \uparrow \forall x$   
we have  $e_0 \in \overline{K}$

if  $e_1 \in \mathbb{N}$  s.t.  $\varphi_{e_1}(x) = 1 \forall x$   
we have  $e_1 \in K$

define  $f: \mathbb{N} \rightarrow \mathbb{N}$  s.t.

$$f(x) = \begin{cases} e_0 & \text{if } x \in K \\ e_1 & \text{if } x \notin K \end{cases}$$

$$= e_0 \cdot \chi_K(x) + e_1 \cdot \chi_{\overline{K}}(x)$$

if  $K$  were recursive then  $\chi_K, \chi_{\overline{K}}$  would be computable and thus  $f$  would be computable.

Since  $f$  is total, by 2<sup>nd</sup> recursion theorem there is  $e \in \mathbb{N}$  s.t.  $\varphi_e = \varphi_{f(e)}$

$$\rightarrow e \in K \quad \rightsquigarrow \quad f(e) = e_0$$

$$\varphi_e(e) \downarrow \neq \varphi_{f(e)}(e) = \varphi_{e_0}(e) \uparrow$$

$$\rightarrow e \notin K \quad \rightsquigarrow \quad f(e) = e_1$$

$$\varphi_e(e) \uparrow \neq \varphi_{f(e)}(e) = \varphi_{e_1}(e) = 1 \downarrow$$

contradiction!

Hence  $K$  is not recursive.

□

\* K is not saturated

$$K = \{x \in \mathbb{N} \mid \varphi_x(x) \downarrow\}$$

We want to show that there are  $e, e' \in \mathbb{N}$  s.t.

$$\begin{array}{l} \varphi_e = \varphi_{e'} \\ e \in K \quad e' \notin K \end{array}$$

\* Assume that there is  $e \in \mathbb{N}$  s.t.

$$\varphi_e(x) = \begin{cases} 0 & \text{if } x = e \\ \uparrow & \text{otherwise} \end{cases}$$

then

- $e \in K$  since  $\varphi_e(e) = 0 \downarrow$
- there  $e' \in \mathbb{N}$   $e' \neq e$  s.t.  $\varphi_e = \varphi_{e'}$
- $e' \notin K$   $\varphi_{e'}(e') = \varphi_e(e') \uparrow \sim e \neq e'$

\* We need to show that there exists  $e \in \mathbb{N}$  s.t.

$$\varphi_e(x) = \begin{cases} 0 & \text{if } x = e \\ \uparrow & \text{otherwise} \end{cases}$$

intuition

kleeemy.py

def P (x) :

if x = "

then return 0

else loop

program we are defining

read ("kleeemy.py")

formally

$$g(e, x) = \begin{cases} 0 & \text{if } x = e \\ \uparrow & \text{otherwise} \end{cases}$$

$$= \mu z. |x - e| \quad \text{computable}$$

by smm theorem there is  $s: \mathbb{N} \rightarrow \mathbb{N}$  total computable s.t.

$$\varphi_{s(e)}(x) = g(e, x) = \begin{cases} 0 & \text{if } x = e \\ \uparrow & \text{otherwise} \end{cases}$$

since  $s$  is total computable, by 2nd recursion theorem

there is  $e_0 \in \mathbb{N}$  s.t.  $\varphi_{e_0} = \varphi_{s(e_0)}$ . Hence

$$\varphi_{e_0}(x) = \varphi_{s(e_0)}(x) = g(e_0, x) = \begin{cases} 0 & \text{if } x = e_0 \\ \uparrow & \text{otherwise} \end{cases}$$

Hence  $e_0$  is the desired program. Thus  $K$  not saturated.

### EXERCISE: RANDOM NUMBERS (from 1<sup>st</sup> lesson)

$\rightarrow m \in \mathbb{N}$  is random if all programs producing  $m$  in output are "larger" than  $m$

two questions:

$\rightarrow$  there are infinitely many random numbers

$\rightarrow$  the property of being random is not decidable

Try again:

$\rightarrow$  size of a program?  $|P_e| = e$

$\rightarrow$  define  $m \in \mathbb{N}$  random if

for all  $e \in \mathbb{N}$  s.t.  $\varphi_e(0) = m$  it holds  $e > m$

### EXERCISE :

Let  $f: \mathbb{N} \rightarrow \mathbb{N}$  be a function

and consider

$$B_f = \{ e \in \mathbb{N} \mid \varphi_e = f \}$$

Are  $B_f, \overline{B_f}$  recursive / r.e. ?

(1)  $f$  not computable

$$B_f = \emptyset, \quad \overline{B_f} = \mathbb{N}$$

recursive  
(hence r.e.)

(2)  $f$  computable

$B_f$  is saturated

$$B_f \neq \emptyset \quad B_f \neq \mathbb{N}$$

Rice  
 $\Rightarrow$

$B_f$  not recursive

$\overline{B_f}$  not recursive

can it be r.e. ? yes it can!

$$f = \emptyset \quad (f(x) \uparrow \forall x)$$

$$\begin{aligned} \overline{B_f} &= \{ e \mid \varphi_e \neq \emptyset \} \\ &= \{ e \mid \underbrace{\exists x. \varphi_e(x) \downarrow} \} \end{aligned}$$

$$SC_{\overline{B_f}}(x) = \exists (\mu \omega. H(x, (\omega)_1, (\omega)_2))$$

complete the exercise!