Euclidean Norm

Throughout this course we will work with functs of "several vars"

$$f(x_1, x_2, \dots, x_d)$$

$$\mathbb{R}^d := \{ \underbrace{\overset{\text{vector}}{(x_1, \dots, x_d)}}_{\vec{x}} : x_1, \dots, x_d \in \mathbb{R} \}$$

On $\mathbb{R}^d$ we have two main operations:

sum: $\vec{x}, \vec{y} \in \mathbb{R}^d$

$\vec{x} = (x_1, \dots, x_d)$

$\vec{y} = (y_1, \dots, y_d)$

$\vec{x} + \vec{y} := (x_1+y_1, x_2+y_2, \dots, x_d+y_d)$

product by scalars

$\vec{x} \in \mathbb{R}^d$, $\lambda \in \mathbb{R}$

$\lambda \vec{x} = (\lambda x_1, \lambda x_2, \dots, \lambda x_d)$

Warning: there is not a prod between vectors.

Rmk: You can sum

$(1,2,-5) + (0,-5,+3) = (1,-3,-2)$

You can't sum vectors with different number of components:

$(1,2,-5) + (0,1)$ does not make sense ∎

Ex: $3(1,0,1) = (3,0,3)$

$-1(1,0,1) = (-1,0,-1)$

$\mathbb{R}^d$, equipped with sum and prod by scalars is a vector space

This means that: the sum is

- (associative) $\vec{x} + (\vec{y} + \vec{z}) = (\vec{x} + \vec{y}) + \vec{z}$
- (commutative) $\vec{x} + \vec{y} = \vec{y} + \vec{x}$
- (zero) $\exists \vec{0} = (0, \dots, 0)$ s.t. $\vec{x} + \vec{0} = \vec{x} \quad \forall \vec{x} \in \mathbb{R}^d$
- (opposite) $\forall \vec{x} \in \mathbb{R}^d \ \exists \vec{y} \in \mathbb{R}^d \ \vec{x} + \vec{y} = \vec{0}$
  (if $\vec{x} = (x_1, \dots, x_d)$, $\vec{y} = (-x_1, \dots, -x_d) = -\vec{x}$)

the product fulfils

- (ass.) $\lambda(\mu \vec{x}) = (\lambda\mu)\vec{x}$
- (unit) $1\vec{x} = \vec{x}$
- (distrib. prop) $\lambda(\vec{x} + \vec{y}) = \lambda\vec{x} + \lambda\vec{y}$
  $(\lambda + \mu)\vec{x} = \lambda\vec{x} + \mu\vec{x}$

The goal is to def the op. of limit.

Let's refresh the def of limit for a seq of real numbers:

Def: (finite limit)

Let $(x_n) \subset \mathbb{R}$ $(x_0, x_1, x_2, \dots)$

We say that $\exists \lim_{n \to +\infty} x_n = \ell \in \mathbb{R}$

if

$\forall \varepsilon > 0 \ \exists N : |x_n - \ell| \le \varepsilon \quad \forall n \ge N$

The goal is to extend this def to

$$\lim_{n \to \infty} \vec{x}_n = \vec{\ell}$$

where $(\vec{x}_n) \subset \mathbb{R}^d$, $\vec{\ell} \in \mathbb{R}^d$.

The key point is to understand what should be

$$\underbrace{|\vec{x}_n - \vec{\ell}|}_{\text{distance between } \vec{x}_n \text{ and } \vec{\ell}}$$

On $\mathbb{R}^2$ $(d=2)$ there's a nat way to do this: $\mathbb{R}^2 = \{(a,b) : a,b \in \mathbb{R}\}$



$\vec{x} = (a_1, b_1)$

$\vec{y} = (a_2, b_2)$

$\ell = \sqrt{|a_2-a_1|^2 + |b_2-b_1|^2}$



$$|\vec{x} - \vec{y}| = \sqrt{(a_2-a_1)^2 + (b_2-b_1)^2}$$



$\ell = \sqrt{(a_2-a_1)^2 + (b_2-b_1)^2 + (c_2-c_1)^2}$

In gen

$\vec{x} = (x_1 \dots x_d)$

$\vec{y} = (y_1 \dots y_d)$

$\text{dist}(\vec{x}, \vec{y}) = \sqrt{(y_1-x_1)^2 + (y_2-x_2)^2 + \dots + (y_d-x_d)^2}$

Def: Given $\vec{x} \in \mathbb{R}^d$, we call euclidean norm of $\vec{x} = (x_1 \dots x_d)$ the quantity

$$\|\vec{x}\| := \sqrt{x_1^2 + x_2^2 + \dots + x_d^2}$$

$$\|\vec{x}\| := \sqrt{\sum_{j=1}^d x_j^2}$$

Warnings: $\|\vec{x}\|$ is a positive (≥ 0) number

Ex: On $\mathbb{R}^2$ $\|\underbrace{(x,y)}_{\text{vect}}\| = \sqrt{x^2+y^2}$ numbers/scalars

On $\mathbb{R}^3$ $\|\underbrace{(x,y,z)}_{\text{vect}}\| = \sqrt{x^2+y^2+z^2}$.

$\|(1,1)\| = \sqrt{1^2+1^2} = \sqrt{2}$

$\|(1,-1)\| = \sqrt{1^2+(-1)^2} = \sqrt{2}$ ← correct

$= \sqrt{1^2-1^2} = 0$ ← wrong

$\|(1,-2)\| = \sqrt{1^2-2^2} = \sqrt{1-4} = \sqrt{-3} = i\sqrt{3} \dots$

$\|(1,0,3)\| = \sqrt{1^2+0^2+3^2} = \sqrt{10}$. ∎

Rmk: What $\|\vec{x}\|$ represents?

$\mathbb{R}^2$: $\|(x,y)\| = \sqrt{x^2+y^2}$



$\|\vec{x}\|$ is the Euclidean dist between $\vec{x}$ and $\vec{0}$. ∎



Prop: The Eucl. norm verifies the following properties:

1. (positivity) $\|\vec{x}\| \ge 0 \quad \forall \vec{x} \in \mathbb{R}^d$
2. (vanishing) $\|\vec{x}\| = 0 \iff \vec{x} = \vec{0}$
3. (homogeneity) $\|\lambda\vec{x}\| = |\lambda| \, \|\vec{x}\|$ $\quad \lambda \in \mathbb{R}$, $\vec{x} \in \mathbb{R}^d$
4. (triang ineq.) $\|\vec{x} + \vec{y}\| \le \|\vec{x}\| + \|\vec{y}\|$

Rmk: (on triang ineq)

We can have $\|\vec{x} + \vec{y}\| < \|\vec{x}\| + \|\vec{y}\|$

$\vec{x}$, $\vec{y} = -\vec{x}$

$\|\vec{x} + \vec{y}\| = \|\vec{0}\| = 0$

$\|\vec{x}\| + \|\vec{y}\| = 2\|\vec{x}\|$

$\|-\vec{x}\| = |-1| \, \|\vec{x}\| = \|\vec{x}\|$ ∎

Proof:

1. true by def: $\|\vec{x}\| = \sqrt{\text{pos. numb}} \ge 0$

2. $\|\vec{x}\| = 0 \iff \sqrt{x_1^2 + \dots + x_d^2} = 0$

$\iff \underset{\overset{\vee}{0}}{x_1^2} + \dots + \underset{\overset{\vee}{0}}{x_d^2} = 0$

$\iff x_1^2 = x_2^2 = \dots = x_d^2 = 0$

$\iff x_1 = x_2 = \dots = x_d = 0$

$\iff \vec{x} = (0, 0, \dots, 0) = \vec{0}$

3. $\|\lambda\vec{x}\| = \sqrt{(\lambda x_1)^2 + (\lambda x_2)^2 + \dots + (\lambda x_d)^2}$
$\quad \lambda^2(x_1^2 + x_2^2 + \dots + x_d^2)$

$\vec{x} = (x_1 \dots x_d)$

$\lambda\vec{x} = (\lambda x_1, \lambda x_2, \dots, \lambda x_d)$

$= \underbrace{\sqrt{\lambda^2}}_{|\lambda|} \underbrace{\sqrt{x_1^2 + \dots + x_d^2}}_{\|\vec{x}\|}$

$= |\lambda| \, \|\vec{x}\|$.

4. $\vec{x} = (x_1 \dots x_d)$

$\vec{y} = (y_1 \dots y_d) \Rightarrow \vec{x} + \vec{y} = (x_1+y_1, \dots, x_d+y_d)$

$\|\vec{x} + \vec{y}\| = \sqrt{\sum_{j=1}^d (x_j+y_j)^2}$

⇓

$\|\vec{x} + \vec{y}\|^2 = \sum_{j=1}^d (x_j+y_j)^2$

$\quad \| \quad (x_j^2 + y_j^2 + 2x_jy_j)$

$= \underbrace{\sum_{j=1}^d x_j^2}_{\|\vec{x}\|^2} + \underbrace{\sum_{j=1}^d y_j^2}_{\|\vec{y}\|^2} + 2\sum_{j=1}^d x_jy_j$

$$\|\vec{x} + \vec{y}\|^2 = \|\vec{x}\|^2 + \|\vec{y}\|^2 + 2\sum_{j=1}^d x_jy_j$$





$c^2 = a^2 + b^2 + 2ab\cos\theta$

Cauchy - Schwarz Inequality:

$$\text{(CS)} \quad \left| \sum_{j=1}^d x_jy_j \right| \le \|\vec{x}\| \, \|\vec{y}\| \quad \forall \vec{x}, \vec{y} \in \mathbb{R}^d$$

(and = holds ⟺ $\vec{x} = \alpha\vec{y}$ or $\vec{y} = \alpha\vec{x}$)

With the (CS) ineq we have:

$\|\vec{x} + \vec{y}\|^2 = \|\vec{x}\|^2 + \|\vec{y}\|^2 + 2\left(\sum x_jy_j\right)$

$-\|\vec{x}\| \, \|\vec{y}\| \le \boxed{\sum x_jy_j \le \|\vec{x}\| \, \|\vec{y}\|}$

$(\ |x| \le \alpha \iff -\alpha \le x \le \alpha \ )$

$\le \|\vec{x}\|^2 + \|\vec{y}\|^2 + 2\|\vec{x}\| \, \|\vec{y}\|$

$= (\|\vec{x}\| + \|\vec{y}\|)^2$

$\Rightarrow \|\vec{x} + \vec{y}\|^2 \le (\|\vec{x}\| + \|\vec{y}\|)^2$

$\Rightarrow \|\vec{x} + \vec{y}\| \le \|\vec{x}\| + \|\vec{y}\|$. ∎

Proof of CS ineq:

$\left| \sum x_jy_j \right| \le \|\vec{x}\| \, \|\vec{y}\|$

$\le \sum |x_jy_j| = \sum |x_j| \, |y_j|$

triang ineq for abs val.

So we prove

$(*) \quad \sum |x_j| \, |y_j| \le \|\vec{x}\| \, \|\vec{y}\| \quad \forall \vec{x}, \vec{y}$.

If $\vec{x} = \vec{0}$ or $\vec{y} = \vec{0}$ ⇒ ineq true. $(0 \le 0)$

So we can assume $\vec{x}, \vec{y} \neq \vec{0} \Rightarrow \|\vec{x}\|, \|\vec{y}\| \neq 0$

$(*) \iff \sum \overset{a}{\boxed{\frac{|x_j|}{\|\vec{x}\|}}} \underset{b}{\boxed{\frac{|y_j|}{\|\vec{y}\|}}} \le 1$

Trick: $(a-b)^2 \ge 0$

$a^2 + b^2 - 2ab \ge 0 \iff 2ab \le a^2 + b^2$

$\boxed{ab \le \frac{a^2+b^2}{2}}$

$\overset{a}{\frac{|x_j|}{\|\vec{x}\|}} \cdot \overset{b}{\frac{|y_j|}{\|\vec{y}\|}} \le \frac{1}{2}\left(\frac{x_j^2}{\|\vec{x}\|^2} + \frac{y_j^2}{\|\vec{y}\|^2}\right)$

$\sum \quad // \quad \le \frac{1}{2}\sum_j \left(\frac{x_j^2}{\|\vec{x}\|^2} + \frac{y_j^2}{\|\vec{y}\|^2}\right)$

$= \frac{1}{2}\left(\underbrace{\frac{1}{\|\vec{x}\|^2} \cdot \sum x_j^2}_{=1} + \underbrace{\frac{1}{\|\vec{y}\|^2} \sum y_j^2}_{=1}\right)$

$= 1$ ∎

Def: (Limit for a seq of vects)

Let $(\vec{x}_n)_n \subset \mathbb{R}^d$. We say that

$$\exists \lim_{n \to +\infty} \vec{x}_n = \vec{\ell} \in \mathbb{R}^d$$

if

$\forall \varepsilon > 0 \ \exists N \in \mathbb{N} : \|\vec{x}_n - \vec{\ell}\| \le \varepsilon \quad \forall n \ge N$.