Friday, 3 October 2025 10:35

Ex 1.8.1

$$\|\vec x+\vec y\|^2 + \|\vec x-\vec y\|^2 = 2\left(\|\vec x\|^2+\|\vec y\|^2\right)$$



Sol: $\to \|\vec x+\vec y\|^2 = \underbrace{\|\vec x\|^2} + \underbrace{\|\vec y\|^2} + \underbrace{\sum x_j y_j}_{(=:\, \vec x \bullet \vec y \text{ scalar prod of } \vec x \text{ and } \vec y)}$

+

$\to \|\vec x-\vec y\|^2 = \|\vec x\|^2 + \|-\vec y\|^2 + \sum x_j(-y_j)$

$= \underbrace{\|\vec x\|^2} + \underbrace{\|\vec y\|^2} - \underbrace{\sum x_j y_j}$

$\|\vec x+\vec y\|^2 + \|\vec x-\vec y\|^2 = 2\|\vec x\|^2 + 2\|\vec y\|^2$ ■

1.8.2 $\|\vec x\|_1 := \sum_{j=1}^{d} |x_j| \qquad \vec x \in \mathbb R^d$

(Manhattan norm)

$\|\vec x\|_\infty := \max_{j=1\ldots d} |x_j|$



Check that they verify the same props of the Eucl. norm.

Let's do for $\|\cdot\|_1$ norm:

- positivity : $\|\vec x\|_1 \geq 0 \quad \forall \vec x \in \mathbb R^d$. Since $\|\vec x\|_1 = \sum |x_j| \geq 0$
- vanishing
- homog.
- triang ineq

Vanishing $\|\vec x\|_1 = 0 \overset{?}{\Leftrightarrow} \vec x = \vec 0$

$\|\vec x\|_1 = 0 \Leftrightarrow \sum_{j=1}^{d} \underset{\substack{\vee\\ 0}}{|x_j|} = 0$

$\Leftrightarrow |x_j| = 0 \ \forall j$

$\Leftrightarrow x_j = 0 \ \forall j$

$\Leftrightarrow \vec x = \vec 0$.

Homog: $\|\lambda \vec x\|_1 = |\lambda|\, \|\vec x\|_1, \quad \forall \lambda \in \mathbb R, \forall \vec x \in \mathbb R^d$

$\|\lambda \vec x\|_1 = \sum_j \underset{\substack{\| \\ \boxed{|\lambda|}\,|x_j|}}{|\lambda x_j|} = |\lambda| \underbrace{\sum_j |x_j|}_{\|\vec x\|_1}$

Triang ineq: $\boxed{\|\vec x+\vec y\|_1} \leq \|\vec x\|_1 + \|\vec y\|_1$ (goal)

$\|\vec x+\vec y\|_1 = \underbrace{\sum_j |x_j+y_j|}$

$|a+b| \leq |a|+|b|$

$|x_j+y_j| \leq |x_j|+|y_j|$

$\Downarrow$

$\sum |x_j+y_j| \leq \sum (|x_j|+|y_j|)$

$\|$

$\|\vec x+\vec y\|_1 \ \textcircled{\leq}\ \underbrace{\sum |x_j|}_{\|\vec x\|_1} + \underbrace{\sum |y_j|}_{\|\vec y\|_1}$ ■

We introduced the def of lim. of a seq. of vectors:

$(\vec x_n) \subset \mathbb R^d, \quad \exists \lim_{n\to+\infty} \vec x_n = \vec\ell \in \mathbb R^d$

$\Leftrightarrow \boxed{\forall \varepsilon>0\ \exists N : \underbrace{\|\vec x_n - \vec\ell\|}_{\delta_n} \leq \varepsilon \ \forall n \geq N.}$

Rmk: $\delta_n := \|\vec x_n - \vec \ell\| \geq 0$

(δ_n) is a seq of positive numbers:

$\lim_n \vec x_n = \vec\ell \Leftrightarrow \forall\varepsilon>0\ \exists N : 0 \leq \delta_n \leq \varepsilon \ \forall n \geq N$

$\Downarrow$

$\boxed{\forall \varepsilon>0\ \exists N : |\delta_n| \leq \varepsilon \ \forall n \geq N}$

$|\delta_n - 0| \leq \varepsilon$

$\Leftrightarrow \lim_n \delta_n = 0$

$\Leftrightarrow \lim_n \|\vec x_n - \vec\ell\| = 0$

$\lim_n \vec x_n = \vec\ell \in \mathbb R^d \Longleftrightarrow \lim_n \|\vec x_n - \vec \ell\| = 0$

Ex (*)

Q. Show that $\lim_n \underbrace{\left(\frac1n, \frac{n+1}{n}\right)}_{\vec x_n} = \overset{\vec\ell}{(0,1)} \quad n \geq 1$



$\vec x_1 = (1, 2)$

$\vec x_2 = (1/2, 3/2)$

$\vec x_3 = (1/3, 4/3)$

Sol: $\vec x_n - \vec \ell = \left(\frac1n, \frac{n+1}{n}\right) - (0,1)$

$= \left(\frac1n - 0, \frac{n+1}{n} - \cancel{1}\right)$

$\cancel{1} + \frac1n$

$= \left(\frac1n, \frac1n\right)$

$\|\vec x_n - \vec\ell\| = \left\|\left(\frac1n, \frac1n\right)\right\|$

$= \sqrt{\frac{1}{n^2} + \frac{1}{n^2}} = \sqrt{\frac{2}{n^2}}$

$= \frac{\sqrt 2}{n}$

$\|\vec x_n - \vec\ell\| = \frac{\sqrt2}{n}$

$\lim_n \|\vec x_n - \vec \ell\| = \lim_n \frac{\sqrt 2}{n} = 0$

$\Rightarrow \exists \lim_{n\to+\infty} \left(\frac1n, \frac{n+1}{n}\right) = (0,1)$. ■

Rmk: $\vec x_n = \underbrace{(x_{n,1}, \ldots, x_{n,d})}_{\text{components of } \vec x_n}$

$\vec\ell = (\ell_1, \ldots, \ell_d)$

Can we say that

$\vec x_n \to \vec\ell \in \mathbb R^d \Leftrightarrow \begin{matrix} x_{n,1} \to \ell_1 \in \mathbb R \\ x_{n,2} \to \ell_2 \in \mathbb R \\ \vdots \\ x_{n,d} \to \ell_d \in \mathbb R \end{matrix}$

$(\lim_n \vec x_n = \vec \ell)$

Ex 1.8.3 (*)

i) $\vec x_n = \left(\boxed{e^{-n}}, 1\right) \in \mathbb R^2$

$\exists \lim \vec x_n = ?$

Here

$x_{n,1} = e^{-n} \longrightarrow 0$

$= \frac{1}{e^n} \to 0$, $e^n \to +\infty$

$x_{n,2} \equiv 1 \longrightarrow 1$

$\Rightarrow \vec x_n \longrightarrow (0,1)$. ■

iii) $\vec x_n = \left(\frac1n, \frac{1}{n^2}, \sin\frac1n\right) \in \mathbb R^3$

Here

$x_{n,1} = \frac1n \longrightarrow 0$

$x_{n,2} = \frac{1}{n^2} \longrightarrow 0 \Rightarrow \vec x_n \to \vec 0$

$x_{n,3} = \sin\frac1n \to 0$

■

Ex: $\vec x_n = \left(\frac1n, (-1)^n\right)$

$x_{n,1} = \frac1n \longrightarrow 0$

$x_{n,2} = (-1)^n \not\to \quad \Rightarrow \vec x_n$ has no limit in $\mathbb R^2$.



■

Def: (infinite limit)

We say that $(\vec x_n) \subset \mathbb R^d$?

has infinite limit

$\left(\lim_{n\to\infty} \vec x_n = \infty_d\right)$

if $\boxed{\lim_n \|\vec x_n\| = +\infty}$

Ex: $\left(\frac1n, n\right) =: \vec x_n$

$\downarrow\ \downarrow$
$0\ +\infty$

Since $x_{n,1} = \frac1n \longrightarrow 0$ but

$x_{n,2} = n \to +\infty$

we can say that $\nexists \lim \vec x_n \in \mathbb R^2$

However since

$\|\vec x_n\| = \sqrt{\underbrace{\underset{\substack{\downarrow\\0}}{\frac{1}{n^2}} + \underset{\substack{\downarrow\\+\infty}}{n^2}}_{\to +\infty}} \longrightarrow +\infty$

$\Rightarrow \vec x_n \longrightarrow \infty_2$.



$\vec x_1 = (1,1)$

$\vec x_2 = \left(\frac12, 2\right)$

$\vec x_3 = \left(\frac13, 3\right)$

■

Ex: $\vec x_n = (n\cos n, n \sin n) = n(\cos n, \sin n)$



$\vec x_1 = (\cos 1, \sin 1)$

$\vec x_2 = 2(\cos 2, \sin 2)$

$\vec x_3 = 3(\cos 3, \sin 3)$

Now notice that

$\|\vec x_n\| = \sqrt{(n\cos n)^2 + (n \sin n)^2}$

$= \sqrt{n^2 \underbrace{(\cos^2 n + \sin^2 n)}_{=1}}$

$= n \longrightarrow +\infty. \quad \Rightarrow \vec x_n \longrightarrow \infty_2$

Notice that

$x_{n,1} = n\cos n \not\to$

$x_{n,2} = n \sin n \not\to$

Warning: it is clear that if one of the components of $\vec x_n$ goes to $\pm\infty \Rightarrow \vec x_n \to \infty_d$

$(\vec x_n = (x_{n,1}, x_{n,2}, \ldots, x_{n,d}))$

$x_{n,2} \longrightarrow +\infty$

$\|\vec x_n\| = \sqrt{\sum_j x_{n,j}^2} \geq \sqrt{x_{n,2}^2} = |x_{n,2}|$

$\underset{x_{n,2}^2}{\vee}$ $\qquad \downarrow$ $+\infty$

$\Rightarrow \|\vec x_n\| \longrightarrow +\infty$

BUT this is not an IFF $\Longleftrightarrow$

The example $\vec x_n = (n\cos n, n \sin n)$ shows that $\vec x_n \to \infty_2$ but none of the components have a lim!

Do 1.8.3/4/5

Limit of a function

In this course we consider several types of functs.

$\underset{\in\mathbb R^2}{f(x,y)} = \underset{\in \mathbb R}{xy^2} \qquad f : \mathbb R^2 \longrightarrow \mathbb R$

$f(\underset{}{1,-5}) = 1\cdot(-5)^2 = \textcircled{25}$

$\underset{\in\mathbb R^3}{f(x,y,z)} = \underset{\in\mathbb R}{x + y \sin z} \qquad f : \mathbb R^3 \longrightarrow \mathbb R$

In general:

$f(x_1, \ldots, x_d) = f(\vec x)$

$f : D \subset \mathbb R^d \longrightarrow \mathbb R$

f is a numerical funct of vect var.

$\vec F(\underset{\in\mathbb R^2}{x,y}) = \underset{\in\mathbb R^2}{(x+y, 2y)} \qquad \vec F : \mathbb R^2 \to \mathbb R^2$



$\vec F(\underset{\mathbb R^2}{x,y}) = \underset{\mathbb R^3}{(x+y, xy, x/y)}$

In general, we consider functs

$\vec F(\underset{\mathbb R^d}{\vec x}) \in \mathbb R^m$

$\boxed{\vec F = \vec F(\vec x) : D \subset \mathbb R^d \longrightarrow \mathbb R^m}$

We want to def

$\lim_{\vec x \to \vec p} \vec F(\vec x) = \vec \ell$

Let's refresh a poss. def of lim for $f = f(x) \quad x \in \mathbb R$:

Def: (Calc 1)

$f : D \subset \mathbb R \longrightarrow \mathbb R$, $p \in \mathrm{Acc}(D)$. We say

$\lim_{x\to p} f(x) = \ell \in \mathbb R$

if

$\forall (x_n) \subset D,\ x_n \neq p,\ x_n \to p \Rightarrow f(x_n) \to \ell$.

Def (Calc 2)

Let $\vec F = \vec F(\vec x) : D \subset \mathbb R^d \longrightarrow \mathbb R^m$, $\vec p \in \mathrm{Acc}\, D$. We say

$\exists \lim_{\vec x \to \vec p} \vec F(\vec x) = \vec \ell \in \mathbb R^m \cup \{\infty_m\}$

if

$\forall (\vec x_n) \subset D,\ \vec x_n \neq \vec p,\ \vec x_n \to \vec p \Rightarrow \vec F(\vec x_n) \to \vec \ell$