Wednesday, 8 October 2025 12:26

Ex 1.8.7

Check that the following lims do not ∃

1. $\lim_{(x,y)\to(0,0)} \frac{x^2-y^2}{x^2+y^2}$.

Sol: $f(x,y) = \frac{x^2-y^2}{x^2+y^2}$ defd on

$D = \mathbb{R}^2 \setminus \{(0,0)\}$, $(0,0) \in \mathrm{Acc}\, D$.

$f(x,0) = \frac{x^2}{x^2} \equiv 1 \xrightarrow[x\to 0]{} \boxed{1}$



$f(0,y) = \frac{-y^2}{y^2} \equiv -1 \xrightarrow[y\to 0]{} \boxed{-1}$

$\Rightarrow \nexists \lim_{(x,y)\to\vec{0}} f$. ∎

2. $\lim_{(x,y)\to(0,0)} \underbrace{\frac{x^2+y^3}{x^2+y^2}}_{f(x,y)}$.

f is defd on
$D = \mathbb{R}^2 \setminus \{(0,0)\}$

$f(x,0) = \frac{x^2}{x^2} \equiv 1 \xrightarrow[x\to 0]{} \boxed{1}$

$f(0,y) = y^3/y^2 = y \xrightarrow[y\to 0]{} \boxed{0}$

$\Rightarrow \nexists \lim_{(0,0)} f$.

4. $\lim_{(x,y,z)\to\vec{0}} \frac{x+y^2+z^3}{\sqrt{\underbrace{x^2+y^2+z^2}_{\geq 0}}}$.

Here $f(x,y,z)$ is defd

$D = \{(x,y,z) \in \mathbb{R}^3 : x^2+y^2+z^2 \neq 0\}$

$= \mathbb{R}^3 \setminus \{(0,0,0)\}$



$f(x,0,0) = \frac{x}{\sqrt{x^2}} = \frac{x}{|x|} = \mathrm{sgn}(x)$.

$\xrightarrow[x\to 0+]{} \boxed{+1}$

$\xrightarrow[x\to 0-]{} \boxed{-1}$ $\Rightarrow \nexists \lim_{(0,0,0)} f$.

∎

6. (**)

$\lim_{(x,y,z)\to\vec{0}} \underbrace{\frac{xz}{x^4+y^2+z^2}}_{f}$

f is defd on $D = \{(x,y,z) : x^4+y^2+z^2 \neq 0\}$

$= \mathbb{R}^3 \setminus \{(0,0,0)\}$

$f(x,0,0) = \frac{0}{x^4} \equiv 0 \xrightarrow[x\to 0]{} \boxed{0}$

$f(x,0,z) = \frac{xz}{x^4+z^2}$

$f(x,0,x^2) = \frac{x \cdot x^2}{x^4+x^4} = \frac{x^3}{2x^4} = \frac{1}{2}\frac{1}{x}$ $\xrightarrow[x\to 0+]{} +\infty$, $\xrightarrow[x\to 0-]{} -\infty$



$(x,0,x^2) \to (0,0,0)$
⇕
$x \to 0$

$\Rightarrow \nexists \lim_{(x,y,z)\to\vec{0}} f$. ∎

⌈ $f(x,0,mx) = \frac{mx^2}{x^4+m^2x^2}$ $m \neq 0$

$= \frac{m\cancel{x^2}}{\cancel{x^2}(m^2+x^2)} \xrightarrow[x\to 0]{} \frac{m}{m^2} = 1/m$ ⌋

Ex 1.8.8

1. (*) $\lim_{(x,y)\to\vec{0}} \underbrace{\frac{xy}{\sqrt{x^2+y^2}}}_{f(x,y)}$. (numerator → 0, $x^2+y^2 \to 0$)

Here f is defd

on $D = \{(x,y) : x^2+y^2 = 0\} = \mathbb{R}^2 \setminus \{(0,0)\}$.



$\boxed{\begin{matrix}(x,y) \to (0,0) \\ \Updownarrow \\ x \to 0 \\ y \to 0\end{matrix}}$
$\Updownarrow$
$\boxed{\rho \to 0 ,\ \theta \in [0,2\pi[}$





$\underset{\begin{cases}x=\rho\cos\theta\\ y=\rho\sin\theta\end{cases}}{f(x,y) =} \frac{(\rho\cos\theta)(\rho\sin\theta)}{\rho}$

$= \frac{\rho^{\cancel{2}}}{\cancel{\rho}} \cos\theta \sin\theta$

$|f(x,y) - 0| = |f(x,y)|$

$= |\rho \cos\theta \sin\theta|$

$= \rho \underbrace{|\cos\theta||\sin\theta|}_{\leq 1}$

$0 \leq |f(x,y) - 0| \leq \rho = \|(x,y)\| \to 0$ (↓ $\vec{0}$)

$\Rightarrow |f(x,y) - 0| \to 0 \Rightarrow f(x,y) \to 0$ ∎

Do #2.

#3. $\lim_{(x,y)\to(0,0)} \frac{x^3-y^3}{x^2+y^2}$.

Here f is defd on $D = \mathbb{R}^2 \setminus \{(0,0)\}$

$\underset{\substack{x=\rho\cos\theta\\ y=\rho\sin\theta\\ x^2+y^2=\rho^2}}{f(x,y) =} \frac{\rho^3\cos^3\theta - \rho^3\sin^3\theta}{\rho^2}$

$= \rho \overbrace{\left(\cos^3\theta - \sin^3\theta\right)}^{\text{bounded}}$ ($\rho \to 0$; $-2 \leq \ldots \leq 2$)

$|f(x,y) - 0| = |f(x,y)| = \rho\, |\cos^3\theta - \sin^3\theta|$ (≤ 2)
↓ 0
$\leq 2\rho = 2\|(x,y)\| \xrightarrow[(x,y)\to(0,0)]{} 0$

$\Rightarrow \exists \lim_{(x,y)\to(0,0)} f = 0$. ∎

We learnt that, to prove

$\lim_{(x,y)\to(0,0)} f(x,y) = 0$

we have to obtain a bound

$|f(x,y)| \leq g(\|(x,y)\|)$

s.t. $g(\rho) \to 0$ when $\rho \to 0$

To prove $\lim f = \ell \in \mathbb{R}$ we need to bound

$|f(x,y) - \ell| \leq g(\|(x,y)\|)$ s.t $g(\rho) \xrightarrow[\rho\to 0]{}$
⇕
$|f(\vec{x}) - \ell| \leq g(\|\vec{x}\|)$ with g s.t $g(\rho) \xrightarrow[\rho\to 0]{} 0$

Prop: If

$\underbrace{|f(\vec{x}) - \ell|}_{\text{dist between } f(\vec{x}) \text{ and } \ell} \leq \underbrace{g(\|\vec{x}\|)}_{\text{a funct of } \|\vec{x}\|}$

s.t $g(\rho) \to 0$ when $\rho \to 0$

$\Rightarrow \exists \lim_{\vec{x}\to\vec{0}} f(\vec{x}) = \ell$.

Ex 1.8.8.

#5. $\lim_{(x,y)\to\vec{0}} \frac{xy}{|x|+|y|}$.

Sol: $f(x,y)$ defd on

$D = \{(x,y) : |x|+|y| \neq 0\} = \mathbb{R}^2 \setminus \{(0,0)\}$.

$|x| + |y| = 0 \iff |x| = |y| = 0$ ($|x|, |y| \geq 0$)

$\iff x = y = 0$

$\underset{\substack{x=\rho\cos\theta\\ y=\rho\sin\theta}}{f(x,y) =} \frac{\rho^{\cancel{2}}\cos\theta\sin\theta}{\cancel{\rho}(|\cos\theta| + |\sin\theta|)}$

$= \rho \frac{\cos\theta\sin\theta}{|\cos\theta| + |\sin\theta|}$. (*)

$\varphi(\theta) = |\cos\theta| + |\sin\theta| \geq K > 0 \quad \forall \theta \in [0,2\pi]$

φ is a cont funct of θ

⇒ (Weierstrass thm) ∃ min φ

$\exists\, \theta_{min} \in [0,2\pi] : \varphi(\theta) \geq \underbrace{\varphi(\theta_{min})}_{K}$

Now I claim $\varphi(\theta_{min}) > 0$

(otherwise, if $\boxed{\varphi(\theta_{min}) = 0}$
⇓
$|\cos\theta_{min}| + |\sin\theta_{min}| = 0$
⇓
$\begin{cases}\sin\theta_{min} = 0\\ \cos\theta_{min} = 0\end{cases}$ imposs!)

This yields to guess $\exists \lim_{(x,y)\to\vec{0}} f = 0$

$|f(x,y) - 0| = |f(x,y)|$

$= \left| \rho \frac{\cos\theta \sin\theta}{|\cos\theta| + |\sin\theta|} \right|$

$\overset{\rho\geq 0}{=} \rho \cdot \frac{\overset{\leq 1}{|\cos\theta|}\,\overset{\leq 1}{|\sin\theta|}}{|\cos\theta| + |\sin\theta| \geq K}$ ≤ 1

$\leq \rho \cdot \frac{1}{K}$ $=: C$

$|f(x,y) - 0| \leq C\rho = g(\|(x,y)\|)$

$g(\rho) \to 0 \quad \rho \to 0$

⇓

$\exists \lim_{(x,y)\to\vec{0}} f = 0$. ∎

Do #4 (similar)

Ex 1.8.9

#2 $\lim_{(x,y,z)\to\vec{0}} \underbrace{\frac{xyz}{x^2+y^2+z^2}}_{f(x,y,z)}$.

f is defd on $\mathbb{R}^3 \setminus \{(0,0,0)\}$.



Spherical Coords.

$\rho = \sqrt{x^2+y^2+z^2}$
$= \|(x,y,z)\|$

(ρ, φ, θ)

$\rho \geq 0, \quad 0 \leq \varphi \leq \pi$
$0 \leq \theta \leq 2\pi$







$x = \rho\cos\theta$
$y = \rho\sin\theta$

$\begin{cases} x = \rho(\sin\varphi)(\cos\theta) \\ y = \rho(\sin\varphi)(\sin\theta) \\ z = \rho\cos\varphi \end{cases}$ $\boxed{x^2+y^2+z^2 = \rho^2}$

Let's write f in spher. coords:

$f(x,y,z) = \frac{\overset{x}{\rho\sin\varphi\cos\theta} \cdot \overset{y}{\rho\sin\varphi\sin\theta}\ \overset{z}{\rho\cos\varphi}}{\rho^2}$

$= \frac{\rho^{\cancel{3}}}{\cancel{\rho^2}} \left(\sin^2\varphi \tfrac{1}{2} 2 \underbrace{\cos\theta \sin\theta}_{\frac{1}{2}\sin(2\theta)} \cos\varphi\right)$

$f(x,y,z) = \rho \cdot \overset{-1\leq \qquad\qquad \leq 1}{\left(\sin^2\varphi \cos\theta \sin\theta \cos\varphi\right)}$

$(x,y,z) \to \vec{0} \iff \underbrace{\|(x,y,z)\|}_{\rho} \to 0$

$|f(x,y,z) - 0| = |f(x,y,z)|$

$= \rho\, \underbrace{|\sin^2\varphi||\cos\theta||\sin\theta||\cos\varphi|}_{\leq 1}$

$\leq \rho = \|(x,y,z)\|$
‖
$g(\rho) \to 0 \quad \rho \to 0$

$\Rightarrow \exists \lim_{(x,y,z)\to\vec{0}} f = 0$ ∎

Do #3

#4 $\lim_{(x,y)\to\vec{0}} \underbrace{\frac{\log(1+2x^3)}{\mathrm{Sh}(x^2+y^2)}}_{f(x,y)}$ $\left(\frac{0}{0}\right)$

Sol:

$D = \{(x,y) : 1+2x^3 > 0, \ \mathrm{Sh}(x^2+y^2) \neq 0\}$

$1 + 2x^3 > 0 \iff x^{\cancel{3}} > -\sqrt[3]{1/2}$



$x > -\sqrt[3]{1/2}$.

$\mathrm{Sh}(x^2+y^2) = 0 \iff x^2+y^2 = 0 \iff x = y = 0$
$\mathrm{Sh}(x^2+y^2) \neq 0 \iff (x,y) \neq (0,0)$

$\underset{\substack{x=\rho\cos\theta\\ y=\rho\sin\theta}}{\log(1+2x^3) =} \log(1+\underbrace{2\rho^3\cos^3\theta}_{\downarrow 0})$

$\log(1+t) = t + o(t) \qquad t \to 0$

$= \boxed{2}\rho^3\cos^3\theta + o(\rho^3 \cancel{\cos^3\theta})$

$= 2\rho^3\cos^3\theta + o(\rho^3)$

$\mathrm{Sh}(x^2+y^2) \underset{\text{pol coords.}}{=} \mathrm{Sh}\,\rho^2 = \rho^2 + o(\rho^2)$

$\mathrm{Sh}\, t = t + o(t)$

$f(x,y) = \frac{2\rho^3\cos^3\theta + o(\rho^3)}{\rho^2 + o(\rho^2)}$

$= \frac{\rho^{\cancel{3}}\left(2\cos^3\theta + \frac{o(\rho^3)}{\rho^3}\right)}{\cancel{\rho^2}\left(1 + \frac{o(\rho^2)}{\rho^2}\right)}$ (→ 0, → 0)

$= \rho \frac{2\cos^3\theta + \varepsilon_\rho}{1 + \varepsilon_\rho} \overset{?}{\longrightarrow} 0$ ($\varepsilon_\rho \to 0$)

$|f - 0| \leq$