Corrected 24 July 2024. See full text.

CONSERVATION

The positive impact of conservation action

Penny F. Langhammer[1,2]*, Joseph W. Bull[3,4,5], Jake E. Bicknell[4], Joseph L. Oakley[6], Mary H. Brown[1], Michael W. Bruford[7,8]†, Stuart H. M. Butchart[9,10], Jamie A. Carr[11,12,13], Don Church[1], Rosie Cooney[14,15]‡, Simone Cutajar[16], Wendy Foden[13,17,18,19], Matthew N. Foster[1], Claude Gascon[20], Jonas Geldmann[10,21], Piero Genovesi[22,23,24], Michael Hoffmann[25,26], Jo Howard-McCombe[27,28], Tiffany Lewis[29], Nicholas B. W. Macfarlane[30], Zoe E. Melvin[27,31], Rossana Stoltz Merizalde[32], Meredith G. Morehouse[33], Shyama Pagad[34], Beth Polidoro[25,35], Wes Sechrest[1], Gernot Segelbacher[8,36], Kevin G. Smith[37], Janna Steadman[4], Kyle Strongin[38], Jake Williams[39], Stephen Woodley[40], Thomas M. Brooks[41,42,43]

Governments recently adopted new global targets to halt and reverse the loss of biodiversity. It is therefore crucial to understand the outcomes of conservation actions. We conducted a global meta-analysis of 186 studies (including 665 trials) that measured biodiversity over time and compared outcomes under conservation action with a suitable counterfactual of no action. We find that in two-thirds of cases, conservation either improved the state of biodiversity or at least slowed declines. Specifically, we find that interventions targeted at species and ecosystems, such as invasive species control, habitat loss reduction and restoration, protected areas, and sustainable management, are highly effective and have large effect sizes. This provides the strongest evidence to date that conservation actions are successful but require transformational scaling up to meet global targets.

Ongoing and substantial declines in global biodiversity as well as the associated negative consequences for human well-being are among the most pressing contemporary risks to society (*1*). Governments have thus adopted goals to tackle biodiversity loss and its drivers: 20 Aichi Biodiversity Targets in the Strategic Plan for Biodiversity 2010–2020 through the Convention on Biological Diversity and now 4 goals for 2050 and 23 targets for 2030 in the 2022 Kunming-Montreal Global Biodiversity Framework (GBF) (*2*). Similar targets are echoed in the 17 sustainable development goals presented in the 2030 Agenda for Sustainable Development (*3*). More than $121 billion is invested annually into biodiversity conservation worldwide (*4*). However, despite this, none of the Aichi targets were fully met (*1*). It could be concluded that responses to the ongoing biodiversity crisis are insufficient, ineffective, or both (*5*, *6*) and that the targets established in the new GBF will also likely not be achieved. However, such conclusions are premature: Conservation interventions could represent progress and be at least partly effective even if global policy targets have not yet been achieved in full. There is consequently a need for a robust evaluation of policy targets and an assessment of whether conservation interventions are working—that is, having positive impacts and providing better outcomes than the absence of interventions—as governments start committing resources and implementing the GBF.

Robust impact assessment using a counterfactual approach (*7*) reveals that conservation action has prevented extinctions (*8*) and reduced extinction risk (*9*) for species across taxonomic groups compared with an absence of conservation action. There has been an increase over the last decade in studies evaluating the impact of specific conservation actions from global to local scales using counterfactual comparisons, including effects of protected areas (*10*), payments for environmental services (*11*), invasive alien species (IAS) eradications (*12*), and sustainable management of ecosystems (*13*). Other studies have undertaken meta-analyses or systematic reviews, but only for individual conservation actions (*14*–*16*). Similarly, the Conservation Evidence website (*17*) provides a compendium of evidence for the effectiveness of a wide range of individual conservation interventions. However, since conservation action started over a century ago there has been no comprehensive meta-analysis of the impact of conservation across the full suite of conservation actions and intervention types, multiple levels and metrics of biodiversity, and over time. Such an assessment is critically needed to inform implementation of the GBF.

We conduct a meta-analysis of the impact of a wide range of conservation interventions globally. Meta-analysis provides a powerful and informative method to summarize results from multiple studies and accounts for unequal precision among studies in the calculation of effect sizes. We evaluate the impact of conservation actions that address direct pressures on biodiversity, promote restoration and recovery of populations and habitat, and aim to safeguard the environment across different levels of biological organization, compared with outcomes expected without intervention. Specifically, we consider seven intervention types aiming to tackle the direct drivers of environmental degradation: (i) establishment and management of protected areas; (ii) other measures to reduce habitat loss and degradation such as policy and restoration; (iii) sustainable use of species; (iv) sustainable management of ecosystems; (v) control of pollution; (vi) eradication and control of invasive alien (and problematic native) species; and finally (vii) climate change adaptation. These classes of intervention were identified based on the strategies of intergovernmental environmental agreements,

[1]Re:wild, PO Box 129, Austin, TX 78767, USA. [2]Arizona State University, School of Life Sciences, PO Box 874501, Tempe, AZ 85287, USA. [3]Department of Biology, University of Oxford, 11a Mansfield Rd, Oxford OX1 3SZ, UK. [4]Durrell Institute of Conservation and Ecology (DICE), School of Anthropology and Conservation, University of Kent, Canterbury CT2 7NR, UK. [5]Wild Business Ltd, London, UK. [6]44 Hillary Road, Maidstone ME14 2JT, UK. [7]School of Biosciences and Sustainable Places Research Institute, Cathays Park, Cardiff CF10 3AX, UK. [8]IUCN SSC Conservation Genetics Specialist Group, 28 rue Mauverney, 1196 Gland, Switzerland. [9]BirdLife International, David Attenborough Building, Pembroke Street, Cambridge CB2 3QZ, UK. [10]Department of Zoology, University of Cambridge, Downing St., Cambridge CB2 3EJ, UK. [11]Leverhulme Centre for Anthropocene Biodiversity, University of York, York YO10 15DD, UK. [12]Department of Environment and Geography, University of York, York YO10 5DD, UK. [13]IUCN SSC Climate Change Specialist Group, 28 rue Mauverney, 1196 Gland, Switzerland. [14]CEESP/SSC IUCN Sustainable Use and Livelihoods Specialist Group, 28 rue Mauverney, 1196 Gland, Switzerland. [15]Fenner School of Environment and Society, Australian National University, ACT 2601, Australia. [16]University of Malta, Msida MSD 2080, Malta. [17]South African National Parks, Cape Research Centre, Tokai, Cape Town, 7966, South Africa. [18]FitzPatrick Institute of African Ornithology, Rondebosch, Cape Town, 7701, South Africa. [19]Global Change Biology Group, Department of Botany and Zoology, Stellenbosch University, Stellenbosch, South Africa. [20]The Global Environment Facility, 1818 H Street NW, Washington, DC 20433, USA. [21]Center for Macroecology, Evolution and Climate, GLOBE Institute, University of Copenhagen, Universitetsparken 15, 2100 Copenhagen E, Denmark. [22]Institute for Environmental Protection and Research, Via Vitaliano Brancati 48, 00144 Rome, Italy. [23]IUCN SSC Invasive Species Specialist Group, 00144 Rome, Italy. [24]Centre for Invasion Biology, Department of Botany and Zoology, Stellenbosch University, Stellenbosch, South Africa. [25]IUCN Species Survival Commission, 28 rue Mauverney, 1196 Gland, Switzerland. [26]Zoological Society of London, Regent's Park, London NW1 4RY, UK. [27]School of Biosciences, Cardiff University, The Sir Martin Evans Building, Museum Avenue, Cardiff, CF10 3AX, UK. [28]RZSS WildGenes, Conservation Department, Royal Zoological Society of Scotland, Edinburgh EH12 6TS, UK. [29]Arizona State University, 427 E. Tyler Mall, Tempe, AZ 85281, USA. [30]IUCN, 1630 Connecticut Ave. NW, Washington, DC 20009, USA. [31]Bangor University, School of Natural Sciences, Deiniol Road, Bangor, Gwynedd, Wales LL57 2UW, UK. [32]Conservation International, 2011 Crystal Drive #600, Arlington, VA 22202, USA. [33]LLaves: Keys to Bilingual Conservation, LLC, 346 Mayberry Hill Road, Casco, Maine 04015, USA. [34]University of Auckland, Auckland 1072, New Zealand. [35]Arizona State University, 4701 W. Thunderbird Rd, Glendale, AZ 85382, USA. [36]University Freiburg, Tennenbacher Str. 4, 79106 Freiburg, Germany. [37]IUCN, The David Attenborough Building, Pembroke St, Cambridge CB2 3QZ, UK. [38]Arizona State University, 800 S. Cady Mall, Tempe, AZ 85281, USA. [39]Imperial College London, Silwood Park, Ascot SL5 7PY, UK. [40]IUCN World Commission on Protected Areas, 64 Juniper Road, Chelsea, Quebec J9B 1T3, Canada. [41]IUCN, 28 rue Mauverney, 1196 Gland, Switzerland. [42]World Agroforestry Center, University of The Philippines Los Baños, Laguna, Philippines. [43]Institute for Marine & Antarctic Studies, University of Tasmania, Hobart, Australia.

***Corresponding author. Email: planghammer@rewild.org**

†Deceased.

‡Present address: ACT Government Office of Nature Conservation, 480 Northbourne Ave, Dickson, ACT 2602, Australia.

Downloaded from https://www.science.org at University of Nevada Reno on September 26, 2026

Corrected 24 July 2024. See full text.

especially the nine Aichi Targets in Strategic Goals B and C of the Strategic Plan for Biodiversity 2011–2020 (table S1), aligned in turn to the targets of the GBF and Sustainable Development Goals 14 (Life Below Water) and 15 (Life on Land). We consider impacts on biodiversity at the ecosystem, species, and genetic level.

We conducted a rapid evidence assessment (see Methods) and meta-analysis of studies published in English that present a counterfactual-based analysis of the impact of conservation actions over time (*18*). Our literature search yielded 1445 studies (published papers) spanning spatial scales from local to continental and more than a century (1890 to 2019) (table S2 and fig. S1). Of these, we retained only studies that contained temporal data and in which we could express outcomes as effect sizes generated from the "rate of change under the intervention" compared with "rate of change under a counterfactual scenario" (*19*), yielding 186 studies that were included in our meta-analysis (table S3). Using a rate of change to calculate the effect size allowed us to assess conservation actions over different time scales, avoiding premature conclusions based on study duration. Where studies measured change in biodiversity using more than one metric (e.g., different species), each was treated as a distinct trial nested within that study, totaling 665 trials across the dataset.

Our comprehensive dataset shows the variable outcomes of conservation action. As such, interventions that generate gains in the state of biodiversity compared with a counterfactual in which biodiversity declines, stays the same, or improves to a lesser degree than the intervention, reveal absolute positive impacts of conservation action (Fig. 1A). Relative positive impacts of conservation action result when biodiversity declines but the intervention slows the decline compared with the counterfactual (Fig. 1B). Conversely, relative negative impacts of conservation action occur when biodiversity improves but the counterfactual reveals greater improvements than the intervention (Fig. 1C). Absolute negative impacts of conservation action result when biodiversity declines following the intervention while it improves, stays the same, or declines to a lesser degree in the counterfactual (Fig. 1D). These four categories are mutually exclusive.




Fig. 1. Schematic representation of different broad categories of conservation impact, with illustrative case studies drawn from our dataset (example reference numbers in superscript). (**A**) Absolute positive impact: intervention outperforms counterfactual and there is an increasing biodiversity trend under the intervention. (**B**) Relative positive impact: intervention outperforms counterfactual but there is a declining biodiversity trend under both intervention and counterfactual. (**C**) Relative negative impact: counterfactual outperforms intervention but there is an increasing biodiversity trend under both intervention and counterfactual. (**D**) Absolute negative impact: counterfactual outperforms intervention and there is a declining biodiversity trend under intervention.

Results

We find that in most cases, biodiversity conservation works. Our meta-analysis shows that the "overall" impact of conservation is positive and significant [mean Hedges' g (± 95% confidence interval) (CI) = 3.24 (2.95 – 3.52), $P < 0.001$], indicating that conservation interventions yield beneficial outcomes for biodiversity compared with the outcome in the absence of an intervention (Fig. 2A). We also show that conservation actions can yield positive impacts in both an absolute and relative sense (Fig. 1). In two-thirds of trials, conservation either improved the state of biodiversity (absolute positive impacts, 45.4%), or at least slowed declines (relative positive impacts, 20.6%). However, in one-fifth of trials, biodiversity under the intervention declined more than no action (absolute negative impacts, 20.6%), whereas in a smaller number of cases biodiversity improved in both the intervention and counterfactual, but the counterfactual revealed greater improvements (relative negative impacts, 11.6%). There was no difference between intervention and counterfactual for 1.8% of trials. Moreover, we find that the effect sizes of some individual interventions are high in magnitude and positive, indicating a substantial positive impact of those actions on the whole.

All types of interventions assessed that had more than five trials showed a significant positive effect compared with a counterfactual (Fig. 2A and fig. S2): eradication and control of invasive alien and problematic native species [7.07 (6.1 to 8.04), $P < 0.001$], sustainable management of ecosystems [5.70 (4.66 to 6.74), $P < 0.001$], habitat loss reduction and restoration [5.58 (4.5 to 6.7), $P < 0.001$], and establishment and management of protected areas [1.41 (1.03 to 1.78), $P < 0.001$]. The impact of efforts toward sustainable use of species is inconclusive [2.07 (−0.71 to 4.84), $P = 0.15$], with large confidence intervals possibly related to the small number of studies [$N_{studies} = 5$, $N_{trials} = 7$]. There were too few studies assessing the impact of interventions of pollution control or climate change adaptation to analyze separately (each had five or fewer trials) but these

Downloaded from https://www.science.org at University of Nevada Reno on September 26, 2026

Corrected 24 July 2024. See full text.




Fig. 2. Effect sizes of conservation interventions. (**A**) Overall and by class of intervention (**B**) for different levels of ecological organization. The number of intervention X counterfactual trials of data (n_{trials}), and unique studies ($n_{studies}$) are reported in parentheses. Mean standardized effect size (Hedges' g) is indicated by the vertical line and 95% confidence intervals (CI) are represented by the bar width. Where the confidence intervals do not overlap zero, the effect size is significant. Vertical dashed lines indicate zero effect and effect sizes to the right indicate that the intervention is more successful than the counterfactual. Interventions with five or fewer trials (pollution control, climate change adaptation, and those classified as "other") are not shown but do contribute to the calculation of the overall effect size. The icons (Copyright BIP/SCBD) in (A) show the primary Aichi target that the intervention classes align with; however as shown in table S1 these interventions also align with a suite of other goals and targets from intergovernmental environmental agreements.

were included in the calculation of the overall effect size. Likewise "other" studies that did not fit into one of the seven key intervention categories [e.g., food supplementation ($N_{studies}$ = 1), culling of diseased individuals ($N_{studies}$ = 1), and captive breeding and release ($N_{studies}$ = 5)] were only included in the calculation of the overall effect size (categorized as "other" in the figures). Studies focused on the impact of controlling problematic native species were combined with studies on IAS given the similarity of the interventions involved.

The impact of actions targeting different levels of ecological organization are positive and significant, with the greatest impact shown for actions targeting species [3.56 (3.14 to 3.97), $P < 0.001$)], followed by ecosystems [2.88 (2.46 to 3.29), $P < 0.001$], and then genetic diversity [3.84 (1.35 to 6.32), $P = 0.002$], which had wide confidence intervals because there were few studies (Fig. 2B). In terms of geographic breakdown, effect sizes were positive and significant on all continents (table S4). Our dataset also shows that effect sizes are positive and significant across each of the differing approaches to measuring the impact of an intervention (table S5).

The timespan of datasets analyzed in the different studies was highly variable, with the shortest being 1 month and the longest 110 years (median = 4.7, mean = 7.4) (Fig. 3). These date back to 1890 and show that studies mainly focused on protected areas until the 1990s but later diversified to a wider range of interventions. Studies with a longer duration were not significantly more likely to show more beneficial or detrimental impacts of conservation actions than shorter studies (fig. S3). Meta-regression of effect sizes against year of publication (Fig. 4) indicated that more recent studies were more likely to show a positive effect of conservation action although the low R^2 shows that other factors influence effect sizes, as would be expected (because many factors in addition to the intervention determine effect sizes).

Half the studies that met our meta-analysis inclusion criteria (95 of 186) were conducted in Western Europe, North America, Australia, and New Zealand (Fig. 3). Of the seven intervention types examined, the largest proportion focused on terrestrial and marine protected areas (38%) and the eradication and control of invasive alien and problematic native species (25%). Fewer studies evaluated other conservation actions for ecosystems (e.g., restoration, habitat conservation policy, sustainable management), for species (e.g., sustainable use, reintroductions), and for genetic diversity (e.g., supplementation, supportive breeding; table S3).

Our results remain largely unchanged in sensitivity analyses designed to test the impact of different methodological considerations (fig. S4), including imputing the rate of change when it was zero in either the intervention or counterfactual, nesting trials within studies, and undertaking a supplemental literature search. Cumulative meta-analysis revealed that effect sizes stabilize after the addition of studies published from approximately 2011 onward (fig. S5) and in assessing publication bias, the symmetrical nature of our funnel plots (fig. S6)—combined with a fail-safe N of 1280—led us to conclude that any publication bias in our dataset is minimal.

Discussion

We have shown that across a full suite of conservation actions and intervention types, multiple levels and metrics of biodiversity, and over a century of action, conservation has improved the state of biodiversity—or at least slowed its

Downloaded from https://www.science.org at University of Nevada Reno on September 26, 2026

Corrected 24 July 2024. See full text.




Fig. 3. Characteristics of the studies (*n* = 186) included in the meta-analysis. Each bar represents a single study included within the meta-analysis and shows (against the *x*-axis) the timescale (start to end years) covered by the dataset in that study (note break in the scale). The color of each bar denotes the intervention type explored by that study. Points show the year in which the associated study was published in the literature. (Inset) World map showing the number of studies carried out in each country represented within the meta-analysis (number of countries is 42).

decline—compared with no conservation action. Our calculated effect sizes are often large and positive, meaning that the outcomes from conservation actions are substantially better than no action at all.

Among the different conservation actions evaluated in our study, the eradication, control, and management of IAS showed the largest impact of conservation action (as highlighted by the largest effect size in our intervention groupings), followed by actions to reduce habitat loss and degradation, sustainable management of ecosystems, and protected areas (Fig. 2A). IAS eradication and control has generated some of the most notable conservation successes reported to date, particularly on islands (*20*, *21*). There were numerous studies evaluating the impact of IAS control and eradication that were excluded from our meta-analysis because they report data for only one time point, but these tend to also show a positive impact.

Although few in number, studies assessing actions to reduce habitat loss and degradation, including restoration, did show positive results (Fig. 2A) consistent with global analyses (*22*, *23*). Efforts to ensure sustainable management of ecosystems, particularly on land, also generally increased native species abundance and habitat cover (Fig. 2A). These results are consistent with global analyses of sustainable management of ecosystems under agriculture and forestry (*24*, *25*).

Protected areas have been shown to be effective in reducing conversion of natural land cover (*10*), terrestrial habitat loss (*15*), coral loss (*26*), tropical forest fires (*27*), species extinction risk (*28*), and in increasing biomass and density of marine organisms (*14*, *29*). Protected area effectiveness varies geographically, in terms of their effectiveness in preventing deforestation (*30*) and reducing anthropogenic pressure (*31*). Poor performance of protected areas often results from shortfalls in human and financial capacities (*32*, *33*), while protected area downgrading, downsizing, and degazettement presents another major challenge in some regions (*34*). However, our results concur that while their effectiveness is not universal (*31*), protected areas are an important tool for achieving conservation outcomes (Fig. 2A). The impacts of efforts to address unsustainable use of species in the ocean and on land were mixed (Fig. 2A), but the number of studies meeting the criteria for inclusion in our meta-analysis was small ($N_{studies}$ = 5).

Across our dataset, 137 trials (21% of cases) provide examples in which conservation interventions

Downloaded from https://www.science.org at University of Nevada Reno on September 26, 2026

Corrected 24 July 2024. See full text.




Fig 4. Publication year versus mean standardized effect sizes for each study, colored by intervention. Blue line and dark gray shaded area show linear line of best fit with 95% CI. Linear regression is significant ($P < 0.001$) but R^2 is small ($R^2 = 0.03$), as would be expected.

were not only associated with a negative rate of change but performed more poorly than counterfactuals for biodiversity state. Eradicating and controlling invasive alien and problematic native species can negatively impact nontarget species through incidental damage or mortality or mesopredator release, and this explains the negative impacts observed for this group of papers. For example, in the US application of herbicides to invasive alien plants harmed native forbs (*35*); in India, physical removal of invasive alien algae caused further spread and establishment elsewhere (*36*) (Fig. 1). Protected areas can show negative impacts if there is poor enforcement and insufficient resourcing, leading to higher rates of resource extraction, poaching, or agricultural expansion compared with counterfactual areas (*37*), or if protection increases the abundance of both predators and the target species in a no-take marine protected area (*38*). These unintended outcomes emphasize the importance of evidence-based interventions alongside effective monitoring, so that negative responses can be quickly detected and conservation measures adjusted accordingly.

More and better counterfactual studies are needed for a wider range of conservation interventions and geographic regions. Particular gaps include assessments of pollution control, climate change adaptation, sustainable use of species, habitat loss reduction (beyond protected areas), actions targeting species and genetic diversity, and conservation actions in the Global South.

The finding that contemporary studies were more likely to show a positive effect of conservation actions (Fig. 4) may be due to increases in funding and project-level resources, more targeted interventions, or that conservation practice is improving over time as lessons are learned from previous failures and methods are improved. These same factors may help explain the larger effect sizes of more rigorous experimental/quasi-experimental study designs (table S5), as their application (particularly quasi-experimental) tends to be more recent. That the relationship between effect size versus study duration (fig. S3) showed no general trend in direction indicates that valuable insights can be gained even for studies that are short in duration, provided the study design is appropriate. This further underscores the necessity of counterfactual evaluation and meta-analysis to assess conservation impact.

Overall, global conservation efforts have helped slow declines in biodiversity and could eventually bend the curve of absolute biodiversity loss (*39*). Quantifying and evidencing the relative biodiversity outcomes (typically gains) of conservation is crucial, to contextualize and explain declining state indicators and increasing response indicator trends. Although the state of biodiversity is declining across the globe in absolute terms, conservation actions work most of the time—the challenge now is to expand these to the scale necessary to reverse the global biodiversity crisis. That is, conservation interventions are working but there are simply not enough conservation actions implemented or in the right places. Realizing the highly ambitious vision of the GBF—not simply to slow declines of biodiversity by the end of the decade but to reverse them (*2*)—will require ongoing assessment of the impact of specific conservation interventions to inform adaptive management with evidence. Importantly, it will require substantially scaled-up funding and commitment for implementation of demonstrably effective conservation interventions—a real transformational change—which in turn depends on increased political will and investment.

Such an increase in conservation action and associated outcomes will require expanded implementation and significant additional investment across many sectors of society, particularly beyond the traditional conservation sector. Meeting global biodiversity conservation targets to reduce the extinction risk of all species and safeguard sites of international biodiversity importance was estimated to cost around $80 billion annually over a decade ago (*40*). A comprehensive global conservation program would require an investment of $178 billion to $524 billion annually (*4*), much of it focused in highly biodiverse countries. Although high, these costs are dwarfed by the value that biodiversity provides to society through the delivery of ecosystem services (*41*). Thus, conservation actions are investments rather than payments—and, as our study demonstrates, they are typically investments that yield genuine, high-magnitude positive impacts.

REFERENCES AND NOTES

1. S. Díaz *et al.*, *Science* **366**, eaax3100 (2019).
2. Convention on Biological Diversity, *Kunming-Montreal Global Biodiversity Framework Draft Decision Submitted by the President*. (UN environment programme, 2022); https://www.cbd.int/doc/c/e6d3/cd1d/daf663719a03902a9b116c34/cop-15-l-25-en.pdf.
3. UN Department of Economic and Social Affairs, Transforming our world: The 2030 Agenda for Sustainable Development. https://sdgs.un.org/publications/transforming-our-world-2030-agenda-sustainable-development-17981 (2015).
4. A. Seidl, K. Mulungu, M. Arlaud, O. van den Heuvel, M. Riva, *Ecosyst. Serv.* **46**, 101216 (2020).
5. J. Terborgh, *Requiem for Nature* (Island Press, 1999).
6. M. Marvier, P. Kareiva, R. Lalasz, Conservation in the Anthropocene. (The Breakthrough Institute, 2012); https://thebreakthrough.org/journal/issue-2/conservation-in-the-anthropocene.
7. P. J. Ferraro, S. K. Pattanayak, *PLOS Biol.* **4**, e105 (2006).
8. F. C. Bolam *et al.*, *Conserv. Lett.* **14**, e12762 (2020).
9. M. Hoffmann *et al.*, *Science* **330**, 1503–1509 (2010).
10. L. N. Joppa, A. Pfaff, *Proc. Biol. Sci.* **278**, 1633–1638 (2011).
11. E. Wiik *et al.*, *Conserv. Sci. Pract.* **1**, e8 (2019).
12. D. P. Armstrong *et al.*, *Conserv. Biol.* **28**, 713–723 (2014).
13. J. M. Holland, B. M. Smith, J. Storkey, P. J. Lutman, N. J. Aebischer, *Biol. Conserv.* **182**, 215–222 (2015).
14. S. E. Lester *et al.*, *Mar. Ecol. Prog. Ser.* **384**, 33–46 (2009).
15. J. Geldmann *et al.*, *Biol. Conserv.* **161**, 230–238 (2013).
16. J. E. Bicknell, M. J. Struebig, D. P. Edwards, Z. G. Davies, *Curr. Biol.* **24**, R1119–R1120 (2014).
17. Conservation Evidence; www.conservationevidence.com [accessed 08 July 2023].
18. H. S. Wauchope *et al.*, *Trends Ecol. Evol.* **36**, 196–205 (2021).
19. I. M. Côté, J. A. Gill, T. A. Gardner, A. R. Watkinson, *Philos. Trans. R. Soc. B* **360**, 385–395 (2005).
20. H. P. Jones *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* **113**, 4033–4038 (2016).
21. D. R. Spatz *et al.*, *Sci. Rep.* **12**, 13391 (2022).
22. J. M. Rey Benayas, A. C. Newton, A. Diaz, J. M. Bullock, *Science* **325**, 1121–1124 (2009).
23. R. Crouzeilles *et al.*, *Nat. Commun.* **7**, 11666 (2016).
24. K. Fedrowitz *et al.*, *J. Appl. Ecol.* **51**, 1669–1679 (2014).

Downloaded from https://www.science.org at University of Nevada Reno on September 26, 2026

Corrected 24 July 2024. See full text.

25. J. Bengtsson, J. Ahnström, A. C. Weibull, . *J. Appl. Ecol.* **42**, 261–269 (2005).
26. E. R. Selig, J. F. Bruno, *PLOS ONE* **5**, e9278 (2010).
27. A. Nelson, K. M. Chomitz, *PLOS ONE* **6**, e22722 (2011).
28. S. H. M. Butchart *et al*., *PLOS ONE* **7**, e32529 (2012).
29. M. Sciberras *et al*., *Fish Fish.* **16**, 58–77 (2015).
30. M. Heino *et al*., *PLOS ONE* **10**, e0138918 (2015).
31. J. Geldmann, A. Manica, N. D. Burgess, L. Coad, A. Balmford, *Proc. Natl. Acad. Sci. U.S.A.* **116**, 23209–23215 (2019).
32. D. A. Gill *et al*., *Nature* **543**, 665–669 (2017).
33. J. Geldmann *et al*., *Conserv. Lett.* **11**, e12434 (2018).
34. R. E. Golden Kroner *et al*., *Science* **364**, 881–886 (2019).
35. E. E. Crone, M. Marler, D. E. Pearson, *J. Appl. Ecol.* **46**, 673–682 (2009).
36. B. Kamalakannan, J. J. J. Jeevamani, N. A. Nagendran, D. Pandiaraja, S. Chandrasekaran, *Curr. Sci.* **106**, 1401–1408 (2014).
37. A. Blackman, A. Pfaff, J. Robalino, *Glob. Environ. Change* **31**, 50–61 (2015).
38. D. Harasti, K. Martin-Smith, W. Gladstone, *PLOS ONE* **9**, e105462 (2014).
39. G. M. Mace *et al*., *Nat. Sustain.* **1**, 448–451 (2018).
40. D. P. McCarthy *et al*., *Science* **338**, 946–949 (2012).
41. A. Balmford *et al*., *Science* **297**, 950–953 (2002).
42. M. O. O'Brien, J. D. Wilson, *Bird Study* **58**, 399–408 (2011).
43. D. Nepstad *et al*., *Conserv. Biol.* **20**, 65–73 (2006).
44. J. Bicknell, Data used in Langhammer et al. 2024., Kent Data Repository (2024); https://doi.org/10.22024/UniKent/01.01.146.

ACKNOWLEDGMENTS

We are grateful to I. Andersen, M. Dalibor, M. Davis, S. Delahaye, A. Hillers, N. Ishii, A. Monard, J.-Y. Pirot, C. Sendashonga, S. N. Stuart, and the late G. Fonseca, for their support in facilitating this project; to J. Koricheva for advice on the meta-analysis; to J. Avanidhar, A. Lamb, and C. Stengel for assistance with the figures; and to E. J. Milner-Gulland, B. W. T. Coetzee, and several anonymous reviewers for their insightful and helpful comments. **Funding:** This work was supported by a Global Environment Facility grant to IUCN on "The Global Environmental Commons: Solutions for a Crowded Planet". **Author contributions:** Conceptualization: P.F.L. and T.M.B. Methodology: P.F.L., J.W.B., J.E.B. and T.M.B. Investigation: P.F.L., J.W.B., J.E.B., J.L.O., M.H.B., M.W.B., S.H.M.B., J.A.C., R.C., S.C., W.F., M.N.F., J.G., P.G., M.H., J.H.-M., T.L., N.M., Z.M., R.M., M.G.M., S.P., B.P., G.S., K.S., J.E.S., K.S., J.W., and S.W. Formal analysis: J.E.B., J.W.B., and P.F.L. Visualization: J.E.B., P.F.L., and J.W.B. Funding acquisition: T.M.B., C.G., N.B.W.M., R.G.M., D.C., and W.S. Writing – original draft: P.F.L., J.W.B., J.E.B., and T.M.B. Writing – review and editing: P.F.L., J.W.B., J.E.B., J.L.O., M.H.B., M.W.B., S.H.M.B., J.A.C., D.C., R.C., S.C., W.F., M.N.F., C.G., J.G., P.G., M.H., J.H.-M., T.L., N.B.W.M., Z.E.M., R.G.M., M.G.M., S.P., B.P., W.S., G.S., K.G.S., J.S., K.S., J.W., S.W., and T.M.B. **Competing interests:** P.F.L., J.W.B., M.H.B., S.H.M.B., D.C., R.C., W.F., M.N.F., C.G., P.G., M.H., J.H.-M., N.B.W.M., R.S.M., W.S., K.G.S., and T.M.B. are employed by organizations engaged in conservation management activities (see author affiliations for details). R.C. serves on the External Scientific Advisory Committee for the Louis Vuitton x People for Wildlife partnership. **Data and materials availability:** The dataset and code generated and analyzed during the current study is available at (*44*). **License information:** Copyright © 2024 the authors, some rights reserved; exclusive licensee American Association for the Advancement of Science. No claim to original US government works. https://www.science.org/about/science-licenses-journal-article-reuse

SUPPLEMENTARY MATERIALS

science.org/doi/10.1126/science.adj6598
Materials and Methods
Figs. S1 to S6
Tables S1 to S5
References (*45–236*)

Submitted 9 July 2023; accepted 14 March 2024
10.1126/science.adj6598

Downloaded from https://www.science.org at University of Nevada Reno on September 26, 2026

Science
AAAS

The positive impact of conservation action

Penny F. Langhammer, Joseph W. Bull, Jake E. Bicknell, Joseph L. Oakley, Mary H. Brown, Michael W. Bruford, Stuart H. M. Butchart, Jamie A. Carr, Don Church, Rosie Cooney, Simone Cutajar, Wendy Foden, Matthew N. Foster, Claude Gascon, Jonas Geldmann, Piero Genovesi, Michael Hoffmann, Jo Howard-McCombe, Tiffany Lewis, Nicholas B. W. Macfarlane, Zoe E. Melvin, Rossana Stoltz Merizalde, Meredith G. Morehouse, Shyama Pagad, Beth Polidoro, Wes Sechrest, Gernot Segelbacher, Kevin G. Smith, Janna Steadman, Kyle Strongin, Jake Williams, Stephen Woodley, and Thomas M. Brooks

Science **384** (6694), . DOI: 10.1126/science.adj6598

Editor's summary

The current biodiversity crisis requires action to protect species from extinction and ecosystems from severe degradation. Billions of dollars are thus invested annually in global conservation. Assessing whether conservation initiatives result in positive outcomes for biodiversity is necessary to guide future efforts. Langhammer *et al*. performed a meta-analysis to determine whether conservation actions, including protected areas and management, provide better outcomes compared with no action. They found that more than half of the time, conservation actions had a net positive effect but did not necessarily stop biodiversity decline. This work shows that multiple types of conservation actions are usually beneficial and are needed to curb the loss of biodiversity. —Bianca Lopez

View the article online

https://www.science.org/doi/10.1126/science.adj6598

Permissions

https://www.science.org/help/reprints-and-permissions

Use of this article is subject to the Terms of service

Science (ISSN 1095-9203) is published by the American Association for the Advancement of Science. 1200 New York Avenue NW, Washington, DC 20005. The title *Science* is a registered trademark of AAAS.

Copyright © 2024 The Authors, some rights reserved; exclusive licensee American Association for the Advancement of Science. No claim to original U.S. Government Works

Downloaded from https://www.science.org at University of Nevada Reno on September 26, 2026